HISKP-TH 03/23

arXiv:hep-ph/0311212v1 17 Nov 2003

Mini-Proceedings of the

Fourth International Workshop on

CHIRAL DYNAMICS : THEORY and EXPERIMENT (CD2003)

Bonn, Germany
September 8 – 13, 2003

edited by

Ulf-G. Meißner, Hans-Werner Hammer and Andreas Wirzba

Helmholtz-Institut für Strahlen- und Kernphysik (Theorie)
Rheinische Friedrich-Wilhelms–Universität Bonn
Nussallee 14-16, D-53115 Bonn, Germany

ABSTRACT

These are the proceedings of the fourth international workshop on "Chiral Dynamics: Theory and Experiment" which was held at the University of Bonn, September 8-13, 2003. The workshop concentrated on the various experimental and theoretical aspects of chiral dynamics, including for the first time in this series detailed discussions of lattice gauge theory. It consisted of an introductory lecture, plenary talks, working group talks and working group summaries. Included is a short contribution per talk.

Fourth International Workshop on
CHIRAL DYNAMICS 2003
Theory and Experiment

Rheinische Friedrich-Wilhelms-Universität Bonn, Germany
September 8-13, 2003
organized by Helmholtz-Institut für Strahlen- und Kernphysik (Theorie)
http://www.itkp.uni-bonn.de/~cd2003

The conference is dedicated to recent progress in theoretical and experimental chiral dynamics and related areas. It consists of invited plenary talks and working group sessions. Working group topics are: *"Goldstone boson dynamics"*, *"meson-baryon dynamics"*, *"few-body dynamics"*, *"chiral lattice dynamics"*, and *"hadronic atoms"*.

Plenary Speakers

S. Beane (UWash Seattle)
V. Bernard (ULP Strasbourg)
J.-W. Chen (Taiwan Nat. Univ.)
H. Calen (TSL Uppsala)
J. Donoghue (UMASS Amherst)
E. Epelbaum (Ruhr-U. Bochum)
J. Goity (Jefferson Lab)
D. Gotta (FZ Jülich)
H.-W. Hammer (Univ. Bonn)
N. Kaiser (TU München)
M. Knecht (CNRS Marseille)
W. Korsch (Univ. Kentucky)
U. Müller (Univ. Mainz)
H. Neufeld (Univ. Wien)
J. Oller (Univ. Murcia)
M. Primavera (INFN Lecce)
J. Schacher (Univ. Bern)
B. Schoch (Univ. Bonn)
J. Schweizer (Univ. Bern)
R. Sommer (DESY Zeuthen)
I. Stewart (MIT)

Working Group Convenors

R. Beck (Univ. Mainz)
P. Bedaque (LBL Berkeley)
J. Bijnens (Univ. Lund)
A. Farilla (INFN Roma III)
T. Hemmert (TU München)
N. Kalantar (KVI Groningen)
E. Pallante (SISSA Trieste)
A. Rusetsky (Univ. Bonn)
M. Savage (UWash Seattle)

Int. Advisory Committee

Aron Bernstein (MIT)
Hans Bijnens (Lund)
Dieter Drechsel (Mainz)
Gerhard Ecker (Wien)
Marc Knecht (Marseille)
Heiri Leutwyler (Bern)
Aneesh Manohar (San Diego)
Giulia Pancheri (Frascati)
Thomas Walcher (Mainz)
Henry Weller (Duke)

Program Committee

Reinhard Beck (Mainz)
Véronique Bernard (Strasbourg)
Ada Farilla (INFN Roma III)
Jürg Gasser (Bern)
Barry Holstein (Amherst)
Norbert Kaiser (München)
Rory Miskimen (Amherst)
Martin Savage (Seattle)
Jan Stern (Orsay)

Local Organisation

Jens Bisplinghoff
Hans-Werner Hammer
Ulf-G. Meißner (chair)
Bernard Metsch
Andreas Wirzba

Ms. C. Schwenk (workshop secr.)
HISKP (Th)
Nussallee 14-16
D-53115 Bonn, Germany
Tel.: +49 - 228 73 2366
Fax: +49 - 228 73 3728
e-mail: cd2003@itkp.uni-bonn.de

Funded by:
BMBF
DFG
FZ Jülich
Jefferson Lab
HISKP(Th)

1 Introduction

This volume presents the mini–proceedings of the workshop on "Chiral Dynamics: Theory and Experiment" which was held at the University of Bonn, September 8-13, 2003. This workshop was the fourth in a series, following MIT (1994), University of Mainz (1997) and Jefferson Laboratory (2000). The fifth one will be organized at TUNL-Duke in 2006.

The unique feature of this series of workshops is the approximately equal mixture of theory and experiment. Due to the rapid developments in the last decade, the range of topics which was discussed has considerably increased over the years. This is reflected in the structure of the working groups, which besides the "classical" topics of Goldstone boson and Goldstone boson-baryon physics were centered around few-body systems and hadronic bound states as well as chiral lattice dynamics.

The workshop consisted of an introductory lecture by John Donoghue, which spurred lively discussions, and 20 plenary talks in the morning sessions. In the afternoons, four working groups took place with about 90 talks in total. The results of these working groups were summarized by the convenors in the plenary session on the last day of the workshop. We had about 140 participants from 20 different countries, with a large amount of younger people, showing that this field is still growing and gaining importance. The outcome of all these talks and the related discussions are summarized in these mini-proceedings. Although the proceedings of the first three workshops were published in conventional proceedings, we believe that the format chosen here better captures the actual status of the field and allows for a quicker access to the presently available literature. Some of the talks are available as PDF or PS file on the conference web page under `www.itkp.uni-bonn.de/~cd2003/`.

Finally, we would like to take this occasion to thank the speakers, the working group convenors, and all the participants for making this workshop a real success.

Acknowledgements

This workshop could not have been done without the dedicated work of many people. First, we would like to thank the conference secretary, Ms. Claudia Schwenk together with Ms. Barbara Mosblech and the staff of the Helmholtz-Institut for all their efforts. Our post–docs and students Paul Büttiker, Matthias Frink, Christian Haupt, Hermann Krebs, Sascha Migura, Lucas Platter, Udit Raha and Akaki Rusetsky performed many different tasks, which we gratefully acknowledge. Special thanks to our co-organizers: Bernard Metsch, who contributed at many different fronts, and to Jens Bisplinghoff for help with the

organization. We are grateful to Herbert Petry for his advice. All members of the advisory and the program committee are warmly thanked for their help and support. Technical assistance in the Wolfgang-Paul lecture hall was provided by Michael Kortmann and we are also grateful to Prof. E. Hilger, the then acting director of the Physikalisches Institut, for support and the allocation of some lecture halls. Financial support was provided by the Deutsche Forschungsgemeinschaft, the Bundesministerium for Bildung und Forschung, Forschungszentrum Jülich, and Jefferson Laboratory, complemented by a contribution from the HISKP.

Bonn, November 2003

Ulf-G. Meißner
Hans-Werner Hammer
Andreas Wirzba

2 Participants and their email

S. Ando	TRIUMF	sando@triumf.ca
V. Anisimovsky	INR Moscow	valera@al20.inr.troitsk.ru
R. A. Arndt	George Washington Univ.	arndtra@said.phys.vt.edu
D. Arndt	Univ. of Washington	arndt@phys.washington.edu
C. Aubin	Washington Univ.	caubin@hbar.wustl.edu
S. R. Beane	Univ. of New Hampshire	silas@physics.unh.edu
D. Bećirević	LPT Orsay	damir.becirevic@th.u-psud.fr
R. Beck	Univ. Mainz	rbeck@kph.uni-mainz.de
V. Bernard	LPT Strasbourg	bernard@lpt6.u-strasbg.fr
A. M. Bernstein	MIT Cambridge	bernstein@lns.mit.edu
J. Bijnens	Univ. Lund	bijnens@thep.lu.se
M. C. Birse	Univ. of Manchester	mike.birse@man.ac.uk
J. Bisplinghoff	Univ. Bonn	jens@iskp.uni-bonn.de
T. Black	UNC Wilmington	blackt@uncw.edu
B. Borasoy	TU München	borasoy@ph.tum.de
A. Braghieri	INFN Pavia	braghieri@pv.infn.it
M. Buechler	Univ. Zürich	buechler@physik.unizh.ch
P. Büttiker	Univ. Bonn	buettike@itkp.uni-bonn.de
D. Cabrera	Univ. Valencia	daniel.cabrera@ific.uv.es
H. Calén	Univ. Uppsala	calen@tsl.uu.se
M. Cargnelli	Austrian Acad. of Sci.	michael.cargnelli@oeaw.ac.at
J.-W. Chen	National Taiwan Univ.	jwc@phys.ntu.edu.tw
T. Chiarappa	DESY Zeuthen	thomas.chiarappa@ifh.de
G. Colangelo	Univ. Bern	gilberto@itp.unibe.ch
M. Davier	LAL Orsay	davier@lal.in2p3.fr
S. Descotes-Genon	LPT Orsay	descotes@th.u-psud.fr
P. Dhonte	Univ. Lund	pierre@thep.lu.se
B. Di Micco	INFN Roma 3	dimicco@fis.uniroma3.it
J. F. Donoghue	Univ. of Massachusetts	donoghue@physics.umass.edu
D. Drechsel	Univ. Mainz	drechsel@kph.uni-mainz.de
S. Dürr	DESY Zeuthen	duerr@ifh.de
G. Ecker	Univ. Wien	gerhard.ecker@univie.ac.at
P. J. Ellis	Univ. of Minnesota	ellis@physics.umn.edu
C. Elster	Ohio Univ.	elster@ohiou.edu
E. Epelbaum	Univ. Bochum	evgeni.epelbaum@tp2.ruhr-uni-bochum.de
A. Farilla	INFN Roma 3	ada.farilla@roma3.infn.it
H. Fearing	TRIUMF	fearing@triumf.ca
L. Fil'kov	Lebedev Institute	filkov@sci.lebedev.ru
H. Fonvieille	CNRS Aubiere	helene@clermont.in2p3.fr
M. Frink	FZ Jülich & Univ. Bonn	mfrink@itkp.uni-bonn.de
E. Frlež	Univ. of Virginia	frlez@virginia.edu

N. Fuchs	Purdue Univ.	nhf@physics.purdue.edu
T. Gail	TU München	tgail@ph.tum.de
A. Gasparian	North Carolina A&T	gasparan@jlab.org
J. Gasser	Univ. Bern	gasser@itp.unibe.ch
J. Gegelia	Univ. Mainz	gegelia@kph.uni-mainz.de
L. Girlanda	ECT* Trento	girlanda@ect.it
S. Giudice	SNS Pisa	sergio.giudice@cern.ch
K.-H. Glander	Univ. Bonn	glander@physik.uni-bonn.de
M. Göckeler	Univ. Leipzig	meinulf.goeckeler@physik.uni-regensburg.de
J. L. Goity	Hampton Univ. & JLab	goity@jlab.org
M. Golterman	San Francisco State Univ.	maarten@stars.sfsu.edu
D. Gotta	FZ Jülich	d.gotta@fz-juelich.de
H.W. Grießhammer	TU München	hgrie@ph.tum.de
C. Haefeli	Univ. Bern	haefeli@itp.unibe.ch
H.-W. Hammer	Univ. Bonn	hammer@itkp.uni-bonn.de
C. Hanhart	FZ Jülich	c.hanhart@fz-juelich.de
C. Haupt	Univ. Bonn	haupt@itkp.uni-bonn.de
T. R. Hemmert	TU München	themmert@physik.tu-muenchen.de
R. P. Hildebrandt	TU München	rhildebr@ph.tum.de
C. Hoelbling	CNRS Marseille	christian.holbling@cern.ch
A. N. Ivanov	Austrian Acad. of Sci.	ivanov@kph.tuwien.ac.at
N. Kaiser	TU München	nkaiser@ph.tum.de
N. Kalantar-N.	KVI Groningen	nasser@kvi.nl
S. Kamalov	JINR Dubna	kamalov@thsun1.jinr.ru
S. Kistryn	Jagellonian Univ.	skistryn@if.uj.edu.pl
M. Knecht	CNRS Marseille	knecht@cpt.univ-mrs.fr
U. van Kolck	Univ. of Arizona	vankolck@physics.arizona.edu
M. Kotulla	Univ. Basel	martin.kotulla@unibas.ch
H. Krebs	Univ. Bonn	hkrebs@itkp.uni-bonn.de
S. Krewald	FZ Jülich	s.krewald@fz-juelich.de
B. Kubis	Univ. Bern	kubis@itp.unibe.ch
R. Lewis	Univ. of Regina	randy.lewis@uregina.ca
R. Lindgren	Univ. of Virginia	ral5q@virginia.edu
M. F. M. Lutz	GSI Darmstadt	m.lutz@gsi.de
A. L'vov	Lebedev Institute	lvov@x4u.lebedev.ru
R. Mawhinney	Columbia Univ.	rdm@phys.columbia.edu
J. McGovern	Univ. of Manchester	judith.mcgovern@man.ac.uk
U.-G. Meißner	Univ. Bonn	meissner@itkp.uni-bonn.de
H. Merkel	Univ. Mainz	merkel@kph.uni-mainz.de
J.G. Messchendorp	KVI Groningen	messchendorp@kvi.nl
B. Metsch	Univ. Bonn	metsch@itkp.uni-bonn.de
S. Migura	Univ. Bonn	migura@itkp.uni-bonn.de
M. A. Moinester	Tel Aviv Univ.	murraym@tauphy.tau.ac.il

I. Montvay	DESY Hamburg	istvan.montvay@desy.de
B. Moussallam	IPN Orsay	moussall@ipno.in2p3.fr
U. Müller	Univ. Mainz	ulm@kph.uni-mainz.de
F. Myhrer	Univ. of South Carolina	myhrer@physics.sc.edu
L. Nemenov	JINR Dubna & CERN	leonid.nemenov@cern.ch
H. Neufeld	Univ. Wien	neufeld@ap.univie.ac.at
F. Nguyen	INFN Roma 3	nguyen@fis.uniroma3.it
R. Nissler	TU München	rnissler@ph.tum.de
A. Nogga	INT Seattle	nogga@phys.washington.edu
V. Obraztsov	IHEP Protvino	obraztsov@mx.ihep.su
J. A. Oller	Univ. of Murcia	oller@um.es
E. Oset	Univ. of Valencia	oset@ific.uv.es
M. Ostrick	Univ. Bonn	ostrick@physik.uni-bonn.de
H. Paetz gen. Schieck	Univ. Köln	schieck@ikp.uni-koeln.de
E. Pallante	SISSA Trieste	pallante@he.sissa.it
V. Pascalutsa	Jefferson Lab.	vlad@jlab.org
S. Peris	Univ. Aut. de Barcelona	peris@ifae.es
H. R. Petry	Univ. Bonn	petry@itkp.uni-bonn.de
H. Pichl	PSI Villigen	hannes.pichl@psi.ch
P. Piirola	Univ. Helsinki	pekko.piirola@helsinki.fi
L. Platter	FZ Jülich & Univ. Bonn	l.platter@fz-juelich.de
A. A. Poblaguev	Yale Univ.	poblaguev@bnl.gov
M. Primavera	INFN Lecce	margherita.primavera@le.infn.it
M. Procura	TU München	massimiliano_procura@ph.tum.de
U. Raha	Univ. Bonn	udit@itkp.uni-bonn.de
F. Rathmann	FZ Jülich	f.rathmann@fz-juelich.de
B. Raue	Florida Int. Univ.	baraue@fiu.edu
L. Roca	Univ. of Valencia	luis.roca@ific.uv.es
H. Rollnik	Univ. Bonn	hrollnik@th.physik.uni-bonn.de
A. Rusetsky	Univ. Bonn	rusetsky@itkp.uni-bonn.de
M. E. Sainio	Univ. Helsinki	sainio@phcu.helsinki.fi
H. Sakai	Univ. of Tokyo	sakai@phys.s.u-tokyo.ac.jp
E. Santos	Univ. Sao Paulo	esantos@fnal.gov
M. Savage	Univ. of Washington	savage@phys.washington.edu
J. Schacher	Univ. Bern	schacher@lhep.unibe.ch
S. Scherer	Univ. Mainz	scherer@kph.uni-mainz.de
M. Schmid	Univ. Bern	schmidm@itp.unibe.ch
B. H. Schoch	Univ. Bonn	b.schoch@uni-bonn.de
J. Schweizer	Univ. Bern	schweizer@itp.unibe.ch
A. Schwenk	Ohio State Univ.	aschwenk@mps.ohio-state.edu
I. Scimemi	Univ. Bern	scimemi@itp.unibe.ch
J. Seely	MIT Cambridge	seely@mit.edu
N. Shoresh	Boston Univ.	shoresh@bu.edu

B. Shwartz	BINP Novosibirsk	shwartz@inp.nsk.ru
S. Sint	Univ. Aut. de Madrid	stefan.sint@uam.es
R. Sommer	DESY Zeuthen & CERN	rainer.sommer@cern.ch
S. Stave	MIT Cambridge	stave@mitlns.mit.edu
E. J. Stephenson	IUCF Bloomington	stephens@iucf.indiana.edu
J. Stern	IPN Orsay	stern@ipno.in2p3.fr
I. W. Stewart	MIT Cambridge	iains@mit.edu
L. Tiator	Univ. Mainz	tiator@kph.uni-mainz.de
W. Tornow	TUNL & Duke Univ.	tornow@tunl.duke.edu
R. Unterdorfer	Univ. Wien	rene.unterdorfer@ap.univie.ac.at
G. Villadoro	Univ. of Rome	giovanni.villadoro@roma1.infn.it
T. Walcher	Univ. Mainz	walcher@kph.uni-mainz.de
A. Wirzba	Univ. Bonn	wirzba@itkp.uni-bonn.de
S. N. Yang	Taiwan National Univ.	snyang@phys.ntu.edu.tw
F. J. Ynduráin	Univ. Aut. de Madrid	fjy@delta.ft.uam.es
P. Zemp	Univ. Bern	zemp@itp.unibe.ch
J. Zmeskal	Austrian Acad. of Sci.	johann.zmeskal@oeaw.ac.at

3 Scientific program

Plenary Talks

Page

J. F. Donoghue
Introductory Lecture: The Physics of the Chiral Effective Field Theory ... 18

J. L. Goity
$\pi^0 \to \gamma\gamma$ *and the PRIMEX* ... 20

M. Primavera
Results from DAΦ*NE* ... 22

V. Bernard
Spin Structure of the Nucleon — Theory ... 24

B. H. Schoch
Polarized Total Photoabsorption Cross Sections ... 26

W. Korsch
Spin Structure of the Nucleon (at low Q^2*) — Experiment* ... 28

U. Müller
Results from the Mainz Microtron ... 30

H. Calén
WASA Physics – Results and Perspectives ... 32

H. Neufeld
Isospin Violation in Semileptonic Kaon Decays ... 34

M. Knecht
The Anomalous Magnetic Moment of the Muon ... 36

J. Schweizer
Hadronic Atoms ... 38

J. Schacher
Results from DIRAC ... 40

D. Gotta
PSI Results from Hadronic Atoms ... 42

J. A. Oller
The Chiral Unitary Approach ... 44

E. Epelbaum
Improving the Convergence of Chiral EFT for Few–Nucleon Systems ... 46

H.-W. Hammer
Limit Cycle Physics ... 48

J.-W. Chen
Nuclear Electroweak Processes in Effective Field Theories ... 50

Page[1]

R. Sommer
Lattice QCD n.r.

S. R. Beane
Nuclear Physics and Lattice QCD 52

I. W. Stewart
The Soft-Collinear Effective Field Theory 54

N. Kaiser
Chiral Dynamics of Nuclear Matter 56

J. Bijnens, A. Farilla
Summary of the Working Group I: Goldstone Boson Dynamics 58

R. Beck, T. R. Hemmert
Summary of the Working Group II: Meson-Baryon Dynamics 60

P. Bedaque, N. Kalantar-Nayestanaki
Summary of the Working Group III: Few-Body Dynamics 62

A. Rusetsky
Summary of the Working Group III: Hadronic Atoms 64

E. Pallante, M. Savage
Summary of the Working Group IV: Chiral Lattice Dynamics 65

[1] n.r.: abstract not received.

Working Group I: Goldstone Boson Dynamics

Convenors: Johan Bijnens and Ada Farilla

Page

S. Giudici
NA48 Results on χPT 68

G. Ecker
$K \to 2\pi$, ϵ'/ϵ and Isospin Violation 69

J. Bijnens
$K \to 3\pi$ in Chiral Perturbation Theory 70

A. A. Poblaguev
E865 Results on K_{e3} 71

V. Obraztsov
High Statistics Study of the $K^- \to e\nu\pi^0$ and $K^- \to \pi^-\pi^0\pi^0$ Decays with ISTRA+ Setup 72

V. Anisimovsky
Measurement of the K_{e3} Form Factors Using Stopped Positive Kaons 73

J. Bijnens
K_{l3} at Two Loops in ChPT 74

E. Frlež
Pion Beta Decay, V_{ud} and $\pi \to l\nu\gamma$ 75

M. A. Moinester
A New Determination of the $\gamma\pi \to \pi\pi$ Anomalous Amplitude Via $\pi^- e \to \pi^- e\, \pi^0$ Data 76

A. Gasparian
The π^0 Lifetime Experiment and Future Plans at JLab 77

M. Büchler
Renormalization Group Equations for Effective Field Theories 78

P. Dhonte
Scalar Form Factors to $O(p^6)$ in SU(3) Chiral Perturbation Theory 79

I. Scimemi
Hadronic Processes and Electromagnetic Corrections 80

R. Unterdorfer
The Generating Functional of Chiral SU(3) 81

G. Colangelo
Hadronic Contributions to a_μ below 1 GeV 82

Page

F. Nguyen
The Measurement of the Hadronic Cross Section with KLOE Via the Radiative Return 83

M. Davier
Hadronic Vacuum Polarization: the μ Magnetic Moment 84

B. Shwartz
CMD-2 Results on $e^+e^- \to$ hadrons 85

P. Büttiker
Roy–Steiner Equations for πK Scattering 86

F. J. Ynduráin
Comments on some Chiral-Dispersive Calculations of $\pi\pi$ Scattering 87

G. Colangelo
On the Precision of the Theoretical Calculations for $\pi\pi$ Scattering 88

J. Stern
From Two to Three Light Flavours 89

S. Descotes-Genon
Bayesian Approach to the Determination of N_f=3 Chiral Order Parameters 90

B. Borasoy
Hadronic Decays of η and η' 91

E. Oset
Φ Radiative Decay to Two Pseudoscalars 92

B. Di Micco
η Decays Studies with KLOE 93

L. Roca
$\eta \to \pi^0\gamma\gamma$ Decay within a Chiral Unitary Approach 94

Working Group II: Meson-Baryon Dynamics

Convenors: Reinhard Beck and Thomas R. Hemmert

Page

Nucleon Spin Structure / GDH

D. Drechsel
GDH, Nucleon Spin Structure and Chiral Dynamics: Theory 96

A. Braghieri
Helicity Dependence of Pion Photo-Production and the GDH Sum Rule .. 97

Real Compton Scattering / VCS

H. Fonvieille
Virtual Compton Scattering at Low Energy and the Generalized Polarizabilities of the Nucleon 98

R. P. Hildebrandt
Dynamical Polarizabilities from (Polarized) Nucleon Compton Scattering 99

L. Fil'kov
Radiative Pion Photoproduction and Pion Polarizability 100

Strangeness Production/SU(3)

K.-H. Glander
Strangeness Production from ELSA 101

B. Raue
Strangeness Production at Jefferson Lab 102

E. Oset
Unitarized Chiral Dynamics: SU(3) and Resonances 103

Pion (Electro) Production and Chiral Dynamics

H. Merkel
Threshold Pion Electroproduction: Update on Experiments 104

M. Kotulla
Two π^0 Production at Threshold 105

S. N. Yang
Threshold π^0 Photo- and Electro-Production 106

Hadronic Reactions / Sigma Term

Page

R. A. Arndt
Partial Wave Analysis of Pion-Nucleon Scattering below $T_{\rm LAB}$ = 2.1 GeV 107

P. Piirola
Towards the Pion-Nucleon PWA 108

M. F. M. Lutz
Meson-Baryon Scattering 109

M. Frink
Order Parameters of Chiral Symmetry Breaking from Meson–Baryon Dynamics 110

Chiral Dynamics and Nucleon Resonances

A. M. Bernstein
Hadron Deformation and Chiral Dynamics 111

V. Pascalutsa
Chiral Dynamics in the $\Delta(1232)$ Region 112

Recent Formal Developments

R. Lewis
Lattice Regularized Chiral Perturbation Theory 113

S. Scherer
Renormalization and Power Counting in Manifestly Lorentz-invariant Baryon Chiral Perturbation Theory 114

J. Gegelia
Baryon ChPT with Vector Mesons 115

P. J. Ellis
Baryon ChPT with Infrared Regularization: Update 116

U.-G. Meißner
Cutoff Schemes in Chiral Perturbation Theory and the Quark Mass Expansion of the Nucleon Mass 117

Working Group III: Few-Body Dynamics and Hadronic Atoms

Convenors: Paulo Bedaque, Nasser Kalantar-Nayestanaki and Akaki Rusetsky

Few-Body Dynamics I

Page

U. van Kolck
Charge Symmetry Breaking in Pion Production 119

E. J. Stephenson
Observation of the Charge Symmetry Breaking $d+d \rightarrow {}^4He+\pi^0$
Reaction near Threshold 120

A. Nogga
Probing Chiral Interactions in Light Nuclei 121

F. Myhrer
Solar Neutrino Reactions and Effective Field Theory 122

S. Ando
Solar-Neutrino Reactions on Deuteron in EFT and*
Radiative Corrections of Neutron Beta Decay 123

M. C. Birse
EFT's for the Inverse-Square Potential and Three-Body Problem 124

Hadronic Atoms

M. Cargnelli
DEAR - Kaonic Hydrogen: First Results 125

J. Gasser
Comments on $\bar{K}N$ *Scattering* 126

A. N. Ivanov
On Pionic and Kaonic Hydrogen 127

P. Zemp
Deser-type Formula for Pionic Hydrogen 128

L. Nemenov
Atoms Consisting of $\pi^+\pi^-$ *and* πK *Mesons* 129

L. Girlanda
Deeply Bound Pionic Atoms: Optical Potential at $O(p^5)$ *in ChPT* 130

Few-Body Dynamics II

Page

H. W. Grießhammer
All-Order Low-Energy Expansion in the 3-Body System 131

T. Black
New Results in the Three and Four Nucleon Systems 132

H. Paetz gen. Schieck
Nd Scattering at Low Energies 133

W. Tornow
Nucleon-Deuteron Scattering at Low Energies 134

S. Kistryn
Three-Nucleon System Dynamics Studied in the dp-Breakup at 130 MeV 135

J. G. Messchendorp
Few-Body Studies at KVI 136

H. Sakai
dp Elastic Scattering Experiments 137

Few-Body Dynamics III

J. F. Donoghue
Nuclear Binding Energies and Quark Masses 138

F. Rathmann
Proton-Proton Scattering Experiments 139

C. Hanhart
Systematic Approaches to Meson Production in NN and dd Collisions .. 140

D. Cabrera
Vector Meson Properties in Nuclear Matter 141

A. Schwenk
A Renormalization Group Method for Ground State Properties of Finite Nuclei 142

L. Platter
The Four-Body System in EFT 143

H. Krebs
Electroproduction of Neutral Pions Off the Deuteron Near Threshold 144

Working Group IV: Chiral Lattice Dynamics

Convenors: Elisabetta Pallante and Martin Savage

Formal Lattice Issues

Page

I. Montvay
Partially Quenched Chiral Perturbation Theory and Numerical Simulations 146

C. Aubin
Pion and Kaon Properties in Staggered Chiral Perturbation Theory 147

R. Mawhinney
Recent Developments in Domain Wall Fermions n.r.

S. Sint
Twisted Mass QCD and Lattice Approaches to the $\Delta I = 1/2$ Rule 148

C. Hoelbling
Overlap Phenomenology 149

Mesons Properties from the Lattice

N. Shoresh
Lattice QCD and Chiral Effective Lagrangians 150

D. Bećirević
Chiral Perturbation Theory, Heavy-light Mesons and Lattice QCD 151

S. Dürr
The Pion Mass in a Finite Volume 152

G. Villadoro
The Lattice Determination of $K \to \pi\pi$ in the $I = 0$ Channel 153

T. Chiarappa
Investigation of the ϵ Regime of ChPT: Meson Correlation Functions ... 154

S. Peris
ϵ'/ϵ and the ΔI=1/2 Rule in the $1/N_c$ Expansion 155

Nucleon Properties from the Lattice

Page

M. Göckeler
Nucleon Form Factors from the QCDSF Collaboration 156

S. R. Beane
N and NN Properties in PQQCD 157

D. Arndt
Partially Quenched Chiral Perturbation Theory for the Baryon and Meson Charge Radii 158

R. Lewis
Strange Matrix Elements in the Nucleon 159

M. Procura
Nucleon Mass and Nucleon Axial Charge from Lattice QCD 160

Future Directions and Focus

M. Golterman
What should be addressed in the next five years 161

The Physics of the Chiral Effective Field Theory

John F. Donoghue

Dept. of Physics, University of Massachusetts,
Amherst, MA 01003, USA

In this talk I looked at the effective field theory aspects of chiral perturbation theory. We have known for a long time about the physics that determines the low energy constants that appear in the chiral lagrangian - these are very often dominated by the effects of the nearest resonances. However, effective field theory goes beyond the lagrangian by including the quantum effects that come from loop corrections. The physics of chiral loops is less understood and the talk focussed on this aspect. Some of the conclusions are perhaps controversial, but I presented the experimental evidence that supports them. A copy of the full talk will be available on my web site, http://www.physics.umass.edu/jdonoghue/ , for the moderate future.

Effective field theory is a procedure for extracting the low energy predictions of a theory. An effective field theory will contain an energy scale that defines the separation of low energy (where the effective field theory is valid) from high energy (where a more complete theory is needed). Loop diagrams always involve the integration over all momentum, including momenta that occur beyond this separations scale. These high energy components of loops in the effective field theory will not be correct because the effective field theory is no longer valid/accurate at these high energies. However, since this inaccuracy arises from short distance physics, it can be corrected for by an adjustment of the local terms in the chiral lagrangian. Nevertheless, it is interesting to explore where this separation occurs in loop processes and to see when the loops are accurate.

I first explored loops as a function of energy. In very many low energy processes, loops are relatively unimportant - the main ingredients to the processes are the low energy constants of the chiral lagrangian. Indeed, in the classic meson sector, the only situation where loops are important is when the two intermediate pions can be in an S-wave. The clearest probe of this physics that I know of is the reaction $\gamma\gamma \to \pi^0\pi^0$, which to order E^4 occurs purely through loops. In this case, the two photon initial state provides a variable energy to probe the loop diagram. Here we have the one-loop and two-loop chiral analysis, plus data, as well as a dispersive analysis based on the Omnes function. (References can be found in [1].) We see clearly that the one-loop analysis fails at $\sqrt{s} = 600$ MeV. The reason for this is clear - loop diagrams are made of tree level vertices and the lowest order $\pi\pi$ S-wave scattering amplitude fails at this energy. A two loop calculation or the use of the Omnes function can improve results as one approaches this energy, but still 600 MeV appears as the scale where the effective field theory breaks down in S-wave pion loops.

I then explored loops as a function of mass. For this exercise, the dispersive

analysis of the $V-A$ vacuum polarization functions is useful because the Weinberg and DMO sum rules allow one to probe the chiral logarithms using data (A portion of this analysis has been published in [2].). The point here is that chiral logarithms arise from the threshold behavior of the dispersive spectral functions, yet in the data one can also see the onset of the high energy regime (which occurs at the rho mass in this channel) beyond which the effective theory is no longer effective. The region between the threshold and the onset of the high energy regime is the region where the effective theory can be applied. We can probe this region as a function of the meson mass by using the chiral prediction for the spectral function in this region. It is easy to see that kaon loops fall outside the region of validity of the effective field theory. Their threshold starts at $\sqrt{s} = 1$ GeV, which is well beyond the separation scale. This and other evidence points to the conclusion that kaon loops are not valid parts of the chiral effective field theory. Moreover, one can compare the chiral parameterization for the loop process (in terms of m^2 and $m^2 \ln m^2$) and the low energy part of the dispersive result. Doing this one learns that the chiral representation is excellent near the pion mass, but fails near meson masses of $m = 250$ MeV. This is lower than most of us would initially expect. However part of the difference is just a factor of two which arises from the fact that the threshold occurs at $\sqrt{s} = 2m$. Thus for this mass, the kinematic threshold suppression is not yet overcome before passing out of the low energy regime.

Finally I discussed regularization schemes, in particular the difference between scale-invariant schemes (variants of dimensional regularization) and scale-dependent schemes (such as various cutoff methods). In the region where the effective theory is valid, there cannot be a difference due to this scheme dependence. However, scale dependent schemes have the ability to probe how much of the loop physics arises from short distances - this is a useful diagnostic. Various examples of this procedure reinforce the conclusion that kaon loops lie outside the range of validity of the effective field theory.

My conclusions were: Pion loops are only important in the $\pi\pi$ S wave; The transition from low-energy to high-energy happens around E = 600 MeV and m = 250 MeV; Pion loops are firmly in EFT region; If you are doing kaon loops, you are not doing effective field theory; Nature uses a cutoff regularization.

References

[1] S. Bellucci, J. Gasser and M. E. Sainio, Nucl. Phys. B **423** (1994) 80 [Erratum-ibid. B **431** (1994) 413] [arXiv:hep-ph/9401206].

[2] J. Sanz-Cillero, J. F. Donoghue and A. Ross, arXiv:hep-ph/0305181 (to be published in Physics Letters).

$\pi^0 \to \gamma\gamma$ and the PRIMEX

Jose L. Goity

Department of Physics, Hampton University,
Hampton, VA 23668, USA, and
Jefferson Lab, Newport News, VA 23606, USA.

In the chiral limit, the amplitude of the decay $\pi^0 \to \gamma\gamma$ is determined by the anomaly induced on the neutral iso-triplet axial current by the EM field. The amplitude is thus given in terms of the only available quantities in that limit, namely, the fine structure and the pion-decay constants. The prediction for the π^0 width using that decay amplitude is in good agreement with the current experimental determination, which has a rather poor precision of about 8%. The deviations from the chiral limit prediction can be theoretically pinned down with rather good precision, as several analyses have shown [1,2,3]. The chief correction is induced by state mixing: due to isospin breaking the π^0 is not a pure isospin state, but it has an admixture with the two iso-singlet states, identified in the chiral limit with the η and η' mesons. The mixing with the first is driven by the ratio $(m_u - m_d)/(m_s - \hat{m})$, and that with the second by the ratio $N_c(m_u - m_d)/\Lambda_\chi$. It turns out that these two mixing effects give each an enhancement of about 1% to the π^0 amplitude. Such an enhancement had been pointed out long ago [4], and the more recent analyses have put it on solid ground, with calculations of strong interaction [2] and electromagnetic [3] corrections at NLO in the chiral and the fine structure constant respectively.

The NLO strong interaction corrections were analyzed in the framework of ChPT combined with the $1/N_c$ expansion [5] (to allow for a consistent inclusion of the η'). In this approach the mixing corrections mentioned above are LO. The NLO corrections to the π^0 amplitude stem from order p^2 and order $1/N_c$ corrections to the decay constants and order p^4, p^2/N_c and $1/N_c^2$ to the masses of the π^0, η and η', as well as a correction from the order p^6 Lagrangian of unnatural parity. These corrections are dominated by effects proportional to m_s/Λ_χ and $1/N_c$, which are expected to correct the enhancement that results from the LO result by an amount of the order of 30%. The actual analysis, based on a global fit involving as inputs the meson masses and decay constants and the η and η' partial widths to two photons, shows that the NLO correction to the π^0 amplitude is within that expectation. The strong interaction effects are then determined to give an increase of the $\pi^0 \to \gamma\gamma$ width (with respect to the result obtained using the chiral limit amplitude) of $4.7 \pm 0.8\%$. The chief uncertainty resides here in the input required for the ratio $R = (m_s - \hat{m})/(m_u - m_d)$. The uncertainty in the EM piece of the $K^+ - K^0$ mass difference leads to the dominant uncertainty in the determination of R. The other sub leading corrections to the π^0 amplitude are EM. It was shown that at order α^2 the amplitude is only affected via EM corrections to the masses and decay constants. The analysis entails the estimate

of a few low energy constants, and the result is that the EM effects tend to reduce the π^0 width by roughly 0.03 eV (or 0.4 %).

In summary, the theoretical prediction that results from the analyses just described is

$$\Gamma_{\pi^0\to\gamma\gamma} = 8.11 \pm 0.07 \text{ eV}.$$

The upcoming Jefferson Lab experiment PRIMEX [6] will determine the π^0 width by means of the Primakoff effect. The coherent photo-production of the π^0 off the Coulomb field of a nucleus (^{12}C, ^{120}Sn and ^{208}Pb) is dominated at small angles by that effect. The Primakoff cross section gives direct access to the $\pi^0 \to \gamma\gamma$ width because it is proportional to it. The measurement of this cross section to high accuracy is therefore the goal of this experiment. The PRIMEX represents a new generation of such experiments (three previous experiments of this kind have a very important weight in the present average value of the width). It has a tagged bremmstrahlung photon beam with uncertainties of less than 1% in photon flux and 10^{-3} in energy, which represents a big improvement over previous experiments. Also, a high resolution hybrid Pb glass and crystal calorimeter provides for a precise determination of the π^0 production angle (uncertainty of the order of 0.01^o). With this the maping the Primakoff peak is possible and in this way the determination of other production mechanisms can be also determined, such as the coherent production via strong interactions whose interference with the Primakoff effect gives the dominating deviation to be taken into account. With this setup, the PRIMEX aims at a determination of the π^0 width at the 1.4% level of precision. This experimental accuracy implies that the enhancement effect of over 4% predicted by theory can be put to the test and in this way uncover experimentally chiral symmetry breaking effects in the π^0 lifetime. The PRIMEX is scheduled to run in early to mid 2004.

References

[1] B. Moussallam, Phys. Rev. **D51** (1995) 4939.

[2] J. L. Goity, A. M. Bernstein and B. R. Holstein, Phys. Rev. **D66** (2002) 076014.

[3] B. Ananthanarayan and B. Moussallam, JHEP **0205** (2002) 052.

[4] Y. Kitazawa, Phys. Lett. **B151** (1985) 165.

[5] P. Herrera-Siklódy, J. I. Latorre, P. Pascual and J. Taron, Nucl. Phys. **B497** (1997) 345, and Phys. Lett. **B419** (1998) 326.
R. Kaiser and H. Leutwyler, Eur. Phys. J. **C17** (2000) 623.

[6] A. Gasparian, these proceedings, and http://www.jlab.org/primex.

Results from DAΦNE

Margherita Primavera[1] on behalf of the **KLOE Collaboration**[1]

[1]INFN Lecce, Via Arnesano, 73100 Lecce, Italy

DAΦNE[2], the Frascati ϕ factory, is an e^+e^- accelerator operating at $W \approx m_\phi \approx 1.02\ GeV/c^2$. The design luminosity is $\approx 5 \cdot 10^{32}\ cm^{-2}s^{-1}$. One of the two machine interaction regions is occupied by the KLOE experiment and the other one is shared in time between the two experiments DEAR and FINUDA.

After the 2002 data taking campaign, in which data of kaonic nitrogen ($\sim 17\ pb^{-1}$) and kaonic hydrogen ($\sim 60\ pb^{-1}$) has been collected, DEAR [3] has measured the three transitions $7 \rightarrow 6, 6 \rightarrow 5, 5 \rightarrow 4$ in kaonic nitrogen and performed a feasibility test of a precision measurement of K^- mass, and has obtained a preliminary measurement of the shift ε and the width Γ of the K_α line of kaonic hydrogen.

FINUDA [4] physics program is concentrated on Λ hypernuclei spectroscopy and decays, especially the non mesonic decays, which are only possible in a nucleus. The detector has been installed during this year and is now ready to collect data.

KLOE [5] is a general purpose detector, optimized to study CP simmetry violation in the neutral kaon system. It consists of a large drift chamber [6] surrounded by a lead/scintillating fiber electromagnetic calorimeter [7]. The detector is immersed in a 0.52 T solenoidal magnetic field, provided by a superconducting coil. The momentum resolution of the drift chamber is $\sigma_p/p \leq 0.4\%$ on tracks with $45° < \theta < 135°$, while calorimeter energy and time resolution are, respectively, $\sigma_E/E = 5.7\%/\sqrt{E(GeV)}$ and $\sigma_t = 54\ ps/\sqrt{E(GeV)} \oplus 50\ ps$. The whole data set collected by KLOE during 2001-2002 amounts to about $450\ pb^{-1}$. Much of this data set has been analized and has yielded improved results on K_S and radiative decays, as well as studies concerning a wide range of topics in kaon and hadronic physics.

Kaon physics at KLOE. At DAΦNE, a tagging tecnique can be applied in order to signal the presence of $K_S(K_L)$ or $K^+(K^-)$ in an event with a $K_L(K_S)$ or $K^-(K^+)$ in the opposite hemisphere. KLOE has measured the ratio $\Gamma(K_S \rightarrow \pi^+\pi^-(\gamma))/\Gamma(K_S \rightarrow \pi^0\pi^0) = 2.236 \pm 0.003 \pm 0.015$ [8], from which the difference of $\pi\pi$ phase shifts in $K \rightarrow \pi\pi$ transitions with I=0 and 2 has been determined: $\chi_0 - \chi_2 = (48 \pm 3)°$, thus improving the agreement with present predictions. KLOE has improved its previous measurement of $\mathrm{BR}(K_S \rightarrow \pi^\pm e^\mp \nu)$[9] obtaining $(6.81 \pm 0.12 \pm 0.10) \cdot 10^{-4}$ and the first-ever measurement of the K_S semileptonic asymmetry $A_S = (19 \pm 17 \pm 6) \cdot 10^{-3}$. For the ratio $\Gamma(K_L \rightarrow \gamma\gamma)/\Gamma(K_L \rightarrow 3\pi^0)$ KLOE obtains the value $(2.793 \pm 0.022 \pm 0.024) \cdot 10^{-3}$[10] and is currently evaluating the BR of the decay channels of K_L to charged particles. On 187 pb^{-1} of 2001-2002 data, KLOE measures $BR(K^\pm \rightarrow \pi^\pm\pi^0\pi^0) = 1.781 \pm 0.013 \pm 0.016\%$ [11]. KLOE should be able to measure all K_{l3} BR's, thus improving the

experimental error on the CKM matrix element $\mid V_{us} \mid$.

ϕ decays and continuum physics at KLOE. KLOE has recently published [12] an analysis of the decay $\phi \to \pi^+\pi^-\pi^0$, mainly proceeding through $\rho\pi$ final states. The fit to the Dalitz plot provides values for the ρ mass and width: $m_\rho = 775.8 \pm 0.5 \pm 0.3\ MeV$ and $\Gamma_\rho = 143.9 \pm 1.3 \pm 1.1\ MeV$. From the ratio of BR's for the decays $\phi \to \eta'\gamma$ and $\phi \to \eta\gamma$ KLOE has obtained a value of the pseudoscalar mixing angle, $\varphi_P = (41.8^{+1.9}_{-1.6})^\circ$ and a limit on the gluonium content of the η' [13]. KLOE has analized, using 17 pb^{-1} of 2000 data, [14,15] the decays $\phi \to \pi^0\pi^0\gamma$ and $\phi \to \eta\pi^0\gamma$, where the dominant contributions come from production and decay of the scalar mesons f_0 and a_0, respectively. Finally, KLOE is concluding the measurement of $\sigma(e^+e^- \to \pi^+\pi^-)$ for $0.3 < s_\pi < 1\ GeV^2$ [16] by using initial state radiation and a MC generator to relate $\sigma(e^+e^- \to \pi^+\pi^-\gamma)$ to $\sigma(e^+e^- \to \pi^+\pi^-)$. The KLOE data provide $a_\mu^{hadr} \times 10^{10}(0.37 < s_\pi < 0.95\ GeV^2) = 374.1 \pm 1.1 \pm 5.2 \pm 2.6_{th}$ plus a $0-2\%$ correction for loss of ISR+FSR events in the simulation.

References

[1] The KLOE author list can be found on http://www.lnf.infn.it/kloe.

[2] S. Guiducci et al., Proc. of PAC99, New York, march 1999.

[3] The DEAR Collaboration, Riv. Nuovo Cimento 22(11), 1 (1999).

[4] The FINUDA Collaboration, LNF-93/021(IR) (1993).

[5] The KLOE Collaboration, LNF-92/019 (IR) (1992).

[6] The KLOE Collaboration, Nucl. Instr. Meth. A488, 51 (2002).

[7] The KLOE Collaboration, Nucl. Instr. Meth. A482, 364 (2002).

[8] The KLOE Collaboration, Phys. Lett. B538, 21 (2002).

[9] The KLOE Collaboration, Phys. Lett. B535, 37 (2002).

[10] The KLOE Collaboration, Phys. Lett. 566, 61 (2003).

[11] The KLOE Collaboration, hep-ex/0307054 (2003).

[12] The KLOE Collaboration, Phys. Lett. B561,55 (2003).

[13] The KLOE Collaboration, Phys. Lett. B541, 45 (2002).

[14] The KLOE Collaboration, Phys. Lett. B537, 21 (2002).

[15] The KLOE Collaboration, Phys. Lett. B536, 209 (2002).

[16] The KLOE Collaboration, hep-ex/0307051 (2003).

Spin Structure of the Nucleon — Theory

V. Bernard[1], T.R. Hemmert[2] and Ulf-G. Meißner [3]

[1]Laboratoire de Physique Théorique, F67084 Strasbourg, France

[2]Technische Universität München, D-85747 Garching, Germany

[3]Universität Bonn, D-53115 Bonn, Germany.

Understanding the spin structure of the nucleon is a central topic of present nuclear and particle physics interest. In the low energy region it allows to test QCD chiral dynamics with processes involving two photons. It is also of particular interest to obtain an understanding of how in QCD the transition from the non-perturbative to the perturbative regime takes place, guided by the precise experimental mapping of spin-dependent observables from low momentum transfer to the multi-GeV region, as it is one of the main thrusts of the research carried out at Jefferson Laboratory (see W. Korsch, these mini-proceedings and [1,2]). The theoretical investigation of the nucleon's spin structure in the non-perturbative regime of QCD can be done using a dispersive analysis (see D. Drechsel, these mini-proceedings). We are utilizing chiral perturbation theory (CHPT) which is based on the spontaneous and explicit chiral symmetry breaking QCD is supposed to undergo. However, baryon CHPT is complicated by the fact that the nucleon mass m does not vanish in the chiral limit and thus introduces a new mass scale apart from the one set by the quark masses. Therefore, any power of the quark masses can be generated by chiral loops, spoiling the one-to-one correspondence between the loop expansion and the one in the small parameter q denoting the pion mass or a small external momenta. One method to overcome this is heavy baryon chiral perturbation theory, (HBCHPT) where the nucleon mass is transformed from the propagator into a string of vertices with increasing powers of $1/m$. However, this method has the disadvantage that certain types of diagrams are at odds with strictures from analyticity. In a fully relativistic treatment, such constraints from analyticity are automatically fulfilled. It was recently argued [3] that relativistic one–loop integrals can be separated into "soft" and "hard" parts. While for the former the power counting as in HBCHPT applies, the contributions from the latter can be absorbed in certain LECs. The underlying method is called "infrared regularization" [4]. It was shown that in this scheme one actually resums graphs with n kinetic insertions on the nucleon line which in the case of HBCHPT would be $1/m^n$ suppressed. This of course leads to better convergence of the chiral serie and should allow to increase the range of applicability of the theory. However this range is limited since one introduces in the calculation of the loop functions an *infinite* Feynman parameter which might result in divergences. In the case of forward double virtual scattering (V^2CS) which is the process under investigation here, they appear at $Q^2 = 1\text{GeV}^2$ in principle well beyond the range of validity of CHPT. However dealing with higher derivatives of the loop functions the influence of the singularity is felt at smaller values of Q^2,

restricting our range to $Q^2_{\rm max} = 0.2 \cdots 0.4 {\rm GeV}^2$. We have performed an $\mathcal{O}(q^4)$ parameter free calculation of the V^{2}CS spin structure functions in the infrared regularization scheme [5]. These are intimately connected to the ones probed in inelastic electroproduction experiments because of unitarity. The $\Delta(1232)$ being well known to play a significant role in the spin sector of the nucleon, we have calculated relativistic Born graphs to get an estimate of its contribution to the spin structure functions.

A typical result for the case of the proton is depicted in the figure which shows the so called generalized Gerasimov-Drell-Hearn sum rule. The zero crossing around 0.25 GeV^2 nicely reproduces the data [1]. The inclusion of the Δ and the vector mesons improves the comparison between theory and experiment as confirmed by the very preliminary data obtained by the CLAS collaboration. However the calculation of the Δ is very conservative and thus leads to large theoretical uncertainties.



A fully systematic treatment is underway, a consistent extension of the Lorentz-invariant method to spin-3/2 fields having been formulated [6]. We have performed a chiral expansion of the forward and longitudinal-transverse spin polarizabilities and found bad convergence in the case of the forward one. The reason rests entirely on the slow convergence of the pole (Born) graphs. A resummation of these is needed, work in this direction is in progress. An understanding of the spin structure of the nucleon in the non perturbative regime of QCD and of the transition to the perturbative one is clearly underway.

References

[1] CLAS collaboration, R. Fatemi et al., nucl-ex/0306019.

[2] M. Amarian et al., Phys. Rev. Lett. 89, 242301 (2002), nucl-ex/0205020

[3] P. J. Ellis and H. B. Tang, Phys. Rev. C 57 (1998) 3356, hep-ph/9709354.

[4] T. Becher and H. Leutwyler, Eur. Phys. J. C 9 (1999) 643, hep-ph/9901384.

[5] V. Bernard et al., Phys. Rev D 67 (2003) 076008, hep-ph/0212033 .

[6] V. Bernard et al., Phys. Lett. B565 (2003) 137, hep-ph/0303198.

Polarized Total Photoabsorption Cross Sections

Berthold H. Schoch

Physikalisches Institut, Universität Bonn,
Nussallee 12, D53115 Bonn, Germany

Fundamental static properties of the nucleons are connected via sum rules with the total photon absorption cross sections of the nucleons. For the case of the unpolarized cross section σ_T the sum of the electric $\alpha(E_\gamma)$ and magnetic $\beta(E_\gamma)$ polarizabilities can be expressed by Baldins sum rule:

$$\alpha + \beta = \frac{1}{2\pi^2} \int_{E_0}^{\infty} \frac{\sigma_T(E_\gamma^{'})}{E_\gamma^{'2}} dE_\gamma^{'} \qquad (1)$$

The energy dependence of the energy weighted total absorption cross section exhibits already the most relevant nucleon excitations for these static properties.Contributions from non resonant single pion production and the first and second resonance region dominate the integral. For the energies above the second resonance region the total absorption cross section can be described by a Regge parametrization which reflects the dominance of diffractive processes. The contributions to the integral from these processes are small due to the inverse quadratic energy weighting of the cross section.

By using a circulary polarized photon beam and a longitudinal – in beam direction – polarized target, helicity 3/2 ($\sigma_{3/2}$) and helicity 1/2 ($\sigma_{1/2}$) cross sections can be measured. The knowledge of these cross sections provides again insight into the origin of static responses for low energy photon scattering. The GDH sum rule connects the anomalous magnetic moment, κ_N , with the difference of the helicity cross sections:

$$\frac{\pi \cdot e^2 \cdot \kappa_N^2}{2m^2} = \int_{E_{threshold}}^{\infty} \frac{\sigma_{3/2}(E_\gamma^{'}) - \sigma_{1/2}(E_\gamma^{'})}{E_\gamma^{'}} dE_\gamma^{'} = I_{GDH} \qquad (2)$$

Due to the only inverse weighting with the gamma energy possible contributions from above the second resonance region of the nucleon may contribute. The value of κ_N is well known. However, it is of great interest where the strength in the excitation spectrum resides in building up κ_N.The helicity cross sections have been measured at MAMI ($200 \leq E_\gamma/MeV \leq 800$) [1] and ELSA ($680 \leq E_\gamma/MeV \leq 2900$) [2]. The cross section, $\sigma_{3/2}(E_\gamma^{'}) - \sigma_{1/2}(E_\gamma^{'})$, differs fom the total unpolarized cross section $(\sigma_{3/2}(E_\gamma^{'}) + \sigma_{1/2}(E_\gamma^{'}))/2$ by an almost complete suppression of the diffractive processes dominating the total unpolarized cross section for $E_\gamma \succ 1200 MeV$. Thus the nucleon resonances stand up more clearly, up to a visible fourth resonance region. This behaviour of the cross sections could have been expected by realizing that the anomalous magnetic moment is a C (charge conjugation) $= -1$ quantity whereas the polarizabilities

transform like C=1 [3]. This makes a difference concerning the coupling of the photon to the constituent quarks and the sea quark/meson cloud, respectively. For the GDH sum rule the following contributions have been extracted from the experiments: MAMI($200 \leq E_\gamma/MeV \leq 800$), $I_{GDH} = 226 \pm 5 \pm 12\mu b$, ELSA ($800 \leq E_\gamma/MeV \leq 2800$), $I_{GDH} = 28.0 \pm 2.1(overall)$. A multipole analysis [4] yields for the energy range, $E^\gamma_{\pi-threshold} \leq E_\gamma \leq 200MeV$, $I_{GDH} = -27.5\mu b$. Based on Regge extrapolations and fits to deep inelastic electron scattering the asymtotic high energy contribution has been estimated to be $I_{GDH} = -15\mu b$[6]. The sum of these contributions adds up to $I_{GDH} = 211.5\mu b$. The overall uncertainty amounts to $\pm 6\%$. Thus, I_{GDH} agrees quite well with the expected value of $I^{\exp ected}_{GDH} = 205\mu b$. Of great interest are the measurements on the neutron. Measurements on the deuteron have been carried out on ELSA and MAMI up to $E_\gamma = 1900MeV$. It remains to be seen, how reliable information on the neutron can be extracted from these data. With the knowledge of $\sigma_{3/2} - \sigma_{1/2}$ the forward spin polarizability γ^p_0 can be extracted via the sum rule:

$$\gamma^p_0 = -\frac{1}{4\pi^2} \int^\infty_{E_{threshold}} \frac{\sigma_{3/2}(E'_\gamma) - \sigma_{1/2}(E'_\gamma)}{E'^3_\gamma} dE'_\gamma \qquad (3)$$

The different contributions yield:
MAMI($200 \leq E_\gamma/MeV \leq 800$), $\gamma^p_0 = -\left[1.87 \pm 0.08(stat) \pm 0.1(syst)\right] \cdot 10^{-4} fm^4$, ELSA ($800 \leq E_\gamma/MeV \leq 2800$), $\gamma^p_0 = -\left[-2.7 \pm 0.2(overall)\right] \cdot 10^{-6} fm^4$. A multipole analysis [4] yields for the energy range, $E^\gamma_{\pi-threshold} \leq E_\gamma \leq 200MeV$, $\gamma^p_0 = [0.9] \cdot 10^{-4} fm^4$. The sum amounts to : $\gamma^p_0 = -\left[1 \pm 0.1\right] \cdot 10^{-6} fm^4$. This value poses a challenge for theory. Various recent calculations are at variance with this result [5].

References

[1] J. Ahrens et al.,Phys.Rev.Lett. 87(2) (2001) 022003.

[2] G. Zeitler, PHD-thesis, University Erlangen, (2002).

[3] S.D. Bass, private communication.

[4] L. Tiator, Proceedings of the 2nd International Symposion on the Grasimov-Drell-Heran sum rule and the spin structure of the nucleon (GDH2002), World Scientific, Singapore, (2003).

[5] D. Drechsel, B. Pasquini, M. Vanderhaeghen, Phys.Rep.378(2-3) (2003) 99.

[6] N. Bianchi and E. Thomas , Phys.Lett.B450 (1999) 439.

Spin Structure of the Nucleon (at low Q^2) — Experiment

Wolfgang Korsch[1]

[1] Dept. of Physics and Astronomy, University of Kentucky,
177 Chemistry - Physics Bldg.,
Lexington, KY 40506, USA

Recent progress in the development of polarized few-nucleon targets and electron beams allows us to perform high precision measurements on the Q^2 evolution of fundamental sum rules related to the spin of the nucleon. During the last 15 years the main focus of nucleon spin structure studies has been the decomposition of the nucleon spin in terms of quarks, gluons, and their orbital angular momenta. However, already in the 1960's Drell, Hearn, and Gerasimov derived a very fundamental sum rule which related the difference of the total photon absorption cross sections, $\sigma_{1/2} - \sigma_{3/2}$, to the square of the anomalous magnetic moments of the nucleon citegdh. Here $\sigma_{3/2}$ and $\sigma_{1/2}$ refer to the cross sections with the nucleon spin aligned parallel or anti-parallel to the spin of the photon. This result can be extended using the low energy expansion of the forward scattering amplitudes in the long wavelength limit of the photon. The first two orders of this expansion for the transverse-transverse and longitudinal-transverse interference amplitudes are given in Eqs. (1) and (2):

$$Re\ \tilde{g}_{TT}(\nu, Q^2) = \frac{2\alpha}{M^2} I_1(Q^2)\nu + \gamma_0(Q^2)\nu^3 + \mathcal{O}(\nu^5) \quad (1)$$

$$Re\ \tilde{g}_{LT}(\nu, Q^2) = \frac{2\alpha}{M^2} Q I_3(Q^2) + Q\delta_{LT}(Q^2)\nu^2 + \mathcal{O}(\nu^4) \quad (2)$$

In the $Q^2 \to 0$ limit I_1 reduces to the GDH sum rule ($\propto \kappa_N^2$). $\gamma_0(0)$ describes the forward spin polarizability of the nucleon and $I_3(0)$ is proportional to $e_N\kappa_N$ (e_N being the charge of the nucleon). The quantity $\delta_{LT}(0)$ is the longitudinal-transverse polarizability. By means of dispersion relations and the optical theorem the above amplitudes can be related to spin dependent structure functions and therefore cross sections [2]. The Q^2 evolution of these static properties of the nucleon is of fundamental theoretical interest, since the low Q^2 behavior can be calculated within the framework of chiral perturbation theory (Ch.P.T.) [3].

The first experiments using a polarized electron beam and polarized targets at Jefferson Lab focused on the Q^2 evolutions of the GDH integral for the proton, deuteron, and the neutron (^{3}He). CEBAF routinely delivers a polarized beam with polarization values of 75% or higher and energies up to 5.7 GeV. In Hall B the large acceptance spectrometer, CLAS, was used in combination with dynamically longitudinally polarized NH3 and ND3 targets to evaluate these integrals from π production threshold to invariant masses, W, up to $\approx$ 3 GeV. Longitudinal asymmetries were measured in inclusive electron − proton (deuteron) scattering. Fits to world data and models were used for the unpolarized structure functions

and transverse spin contributions in order to extract the spin structure functions $g_1^{p,d}(x,Q^2)$. The first moments of $g_1^{p,d}(x,Q^2)$ were extracted in a Q^2 range from about 0.1 GeV^2 to about 3.5 GeV^2 [4]. It was found that both, g_1^p and g_1^d depend strongly on Q^2. g_1^p changes sign at a Q^2 value around 0.3 GeV^2 whereas the sign change for g_1^d appears to occur between 0.5 GeV^2 and 0.6 GeV^2. The zero crossings of the structure functions are an indication of the transition where non-partonic degrees of freedom such as resonances become dominant. Therefore, resonance contributions seem to be slightly less important for the proton.

The high precision data are in nice agreement with chiral perturbation theory calculations for Q^2 values less than 0.2 GeV^2.

Hall A used a dense polarized ^{3}He gas target to measure the generalized GDH integral [5], the generalized spin polarizability, and the longitudinal−transverse polarizability for the neutron. The spin dependent cross sections were measured directly by polarizing the target longitudinally and transversely to the direction of the incident beam. Nuclear corrections were applied according to Ref. [6]. Results in a Q^2 range from 0.10 GeV^2 to 0.90 GeV^2 were presented [7]. The GDH integral for the neutron exhibits a strong Q^2 dependence as well. The MAID model [8] underestimates the GDH integral at low values of Q^2 but describes both, $\gamma_0(Q^2)$ and $\delta_{LT}(Q^2)$, quite well. Ch.P.T. calculations are in good agreement with the data for the GDH integral and $\gamma_0(Q^2)$ for $Q^2 < 0.2$ GeV^2 [3].

The combination of the Hall A and Hall B data will allow us to evaluate the difference of the first moments $\Gamma_1^p(Q^2) - \Gamma_1^n(Q^2)$. In the high Q^2 regime this difference is known as the Bjorken sum rule. It has practically no sensitivity to sea quarks in the asymptotic limit and is also less sensitive to the Δ excitation in the low Q^2 regime. Therefore, the comparison of this difference can serve as a crucial test for theoretical calculations.

References

[1] S.D. Drell and A.C. Hearn, Phys. Rev. Lett. **16** (1966) 908;
S.B. Gerasimov, Sov. J. Nucl. Phys. **2** (1966) 430.

[2] D. Drechsel *et al.*, Phys. Rept. **378** (2003) 99.

[3] V. Bernard, *these proceedings.*

[4] R. Fatemi *et al.*, nucl-ex/0306019; V. Burkert, private communication.

[5] M. Amarian *et al.*, Phys. Rev. Lett. **89** (2002) 242301.

[6] C. Ciofi degli Atti, S. Scopetta; Phys. Lett. B **404** (1997) 223.

[7] M. Amarian *et al.*, hep-ex/0310003v2.

[8] D. Drechsel *et al.*, Phys. Rev. D **63** (2001) 114010.

Results from the Mainz Microtron

Ulrich Müller[1]
representing the A1, A2, and A4 collaborations

[1]Institut für Kernphysik, Universität Mainz,
Johann-Joachim-Becher-Weg 45, 55099 Mainz, Germany

Some highlights of recent experimental results at MAMI-B [1] are presented.

The neutron electric form factor $G_{E,n}$ has been measured at momentum transfers of $Q^2 = 0.30$, 0.59, and $0.79\,(\mathrm{GeV}/c)^2$ in a $\mathrm{D}(\vec{e}, e'\vec{n})p$ experiment of the A1 collaboration, using a spin precession method and neutron recoil polarimetry. Preliminary values of $G_{E,n} = 0.0552 \pm 0.0061\,(\mathrm{stat}) \pm 0.0055\,(\mathrm{syst})$, $0.0469 \pm 0.0073\,(\mathrm{stat}) \pm 0.0054\,(\mathrm{syst})$, and $0.047 \pm 0.009\,(\mathrm{stat})\,^{+0.001}_{-0.002}\,(\mathrm{syst})$, have been obtained, respectively [2].

An extensive programme of charged pion electroproduction off the proton has been performed by the A1 collaboration. The reaction $\mathrm{H}(e, e'\pi^+)n$ has been measured at a constant invariant mass of $W = 1125\,\mathrm{MeV}$ and four-momentum transfers of $Q^2 = 0.058$, 0.117, 0.195, and $0.273\,(\mathrm{GeV}/c)^2$. For each Q^2 value, measurements at different values of ϵ, the virtual photon polarisation parameter, allow for a Rosenbluth separation of longitudinal and transverse cross sections. Analysis of the data taken in 2000 and 2002 is in progress [3].

Threshold production of neutral pions has been measured in an $\mathrm{H}(e, e'p)\pi^0$ experiment by the A1 collaboration at a momentum transfer of $Q^2 = 0.05\,(\mathrm{GeV}/c)^2$ and $\epsilon = 0.933$. Cross sections of the structure functions σ_0, σ_{LT}, σ_{TT}, and $\sigma_{\mathrm{LT}'}$ and the helicity asymmetry $A_{\mathrm{LT}'}$ have been determined [4]. The helicity-independent cross sections are described by the MAID 2000 model [5] within their errors. The magnitude of the transverse-transverse part is significantly overestimated by ChPTh [6].

The A2 collaboration used the DAPHNE detector to measure the helicity dependent exclusive cross sections for the absorption of polarised photons by longitudinally polarised nucleons. The cross section integral of the Gerasimov-Drell-Hearn sum rule from 200 to 800 MeV was calculated from the data as $I_p^{\mathrm{GDH}} = (-226 \pm 5\,(\mathrm{stat}) \pm 12\,(\mathrm{syst}))\,\mu\mathrm{b}$ [7]. After a correction for the missing region below 200 MeV and by including ELSA data above 800 MeV [8], a value of $I_p^{\mathrm{GDH}} = (-227.5 \pm 5\,(\mathrm{stat}) \pm 12\,(\mathrm{syst}) \pm 3\,(\mathrm{theo}))\,\mu\mathrm{b}$ is obtained.

The A2/TAPS collaboration determined the E2 amplitude of the $N \to \Delta$ transition in an $\mathrm{H}(\vec{\gamma}, \pi^0)p$ experiment using linear polarised photons in the energy range from 200 to 790 MeV. At the position of the Δ resonance, $E_\gamma = 340\,\mathrm{MeV}$, a preliminary value of $R_{EM} = (-2.40 \pm 0.16 \pm 0.24) \cdot 10^{-2}$ is obtained [9]. This is in agreement with a previous A2/DAPHNE measurement [10].

The reaction $\mathrm{H}(\gamma, p\pi^0\gamma')$ has been measured by the A2 collaboration with the TAPS calorimeter for energies between $\sqrt{s} = 1221$ and 1331 MeV. Cross sections differential in angle and energy have been determined for all particles in the final

state in three bins of the excitation energy. A value for the magnetic dipole moment of the $\Delta^+(1232)$ resonance of $2.7^{+1.0}_{-1.3}\,(\text{stat}) \pm 1.5\,(\text{syst}) \pm 3\,(\text{theo})$ nuclear magnetons has been extracted for the first time [11].

Parity violating electron scattering by the A4 collaboration using counting techniques gives access to the strange vector form factors of the nucleon. A preliminary value of $A_{\text{phys}} = (-5.6 \pm 0.6 \pm 0.2) \cdot 10^{-6}$ corresponding to a strange quark contribution of $A_{\text{phys}} - A_0 = (0.9 \pm 0.7) \cdot 10^{-6}$ has been extracted at a beam energy of 854.3 MeV, based on $4.6 \cdot 10^{12}$ events [12]. Analysis of data taken at 570.1 MeV is in progress.

References

[1] H. Herminghaus et al., Proceedings 1990 Linear Accelerator Conference, Albuquerque, NM, LA-12004-C (1990) 362.

[2] M. Seimetz, doctoral thesis, Univ. Mainz, 2003; D. Glazier, PhD thesis, Univ. Glasgow, 2003; to be published.

[3] D. Baumann, doctoral thesis, Univ. Mainz, in preparation.

[4] M. Weis, doctoral thesis, Univ. Mainz, 2003; to be published.

[5] D. Drechsel, O. Hanstein, S. S. Kamalov, L. Tiator, Nucl. Phys. A 645 (1999) 145.

[6] V. Bernard, N. Kaiser, U.-G. Meißner, Nucl. Phys. A 607 (1996) 379; 633 (1998) 695 (Erratum).

[7] J. Ahrens et al., Phys. Rev. Lett. 87 (2001) 022003 and references therein.

[8] H. Dutz et al., Phys. Rev. Lett. 91 (2003) 192001.

[9] R. Leukel, doctoral thesis, Univ. Mainz, 2001; to be published.

[10] R. Beck et al., Phys. Rev. Lett. 78 (1997) 606.

[11] M. Kotulla et al., Phys. Rev. Lett. 89 (2002) 272001.

[12] F. Maas, private communication; to be published.

WASA Physics - Results and Perspectives

Hans Calén[1]
for the CELSIUS/WASA collaboration

[1]The Svedberg Laboratory, Uppsala University,
P.O. Box 533, SE-751 21 Uppsala, Sweden

The CELSIUS accelerator and storage ring at the The Svedberg Laboratory in Uppsala is used for hadron physics research in the GeV region. The programme includes production and decay of π and η mesons in light ion collisions at the WASA 4π detector facility [1].

The CELSIUS ring provides protons of kinetic energies up to 1360 MeV. For proton energies up to 500 MeV, electron cooling can be used, and this gives a beam size around 1 mm and a relative momentum spread down to the 10^{-5} region.

The recently developed pellet-target system provides small spheres of frozen hydrogen as internal targets. This allows high luminosity and high detection coverage for meson decay products like photons, electrons and charged pions. The target system is now operating close to design performance with hydrogen pellets having a diameter of about 25 micron, a speed of 80 m/s and occuring at a rate of up to 15000 pellets/s. The pellets are guided 2.5 m in a thin tube to the CELSIUS beam and then further to a dump. The pellet beam diameter at the interaction region is 2-3 mm.

Deuterium pellet generation has been tested successfully at different occasions and now a useful deuterium pellet beam will be developed using a separate test station that is being prepared.

The pellet target is integrated in the WASA 4π detector facility. WASA consists of a forward part for measurements of charged target-recoil particles and scattered projectiles, and a central part designed for measurements of the meson decay products. The forward part consists of eleven planes of plastic scintillators and of proportional counter drift tubes. The central part consists of an electromagnetic calorimeter of CsI(Na) crystals surrounding a superconducting solenoid. Inside of the solenoid are placed a cylindrical drift chamber and a barrel of plastic scintillators.

Initially, production of ηs and multiple πs in pp collisions have been measured. The two-pion production studies started in the mid nineties at the CELSIUS cluster-jet target [2] have been repeated and extended to higher energies exploiting the much larger acceptance at WASA, in particular for the $2\pi^0$ channel. Data from threshold up to excess energies, Q, around 100 MeV are well described by model calculations [3] involving heavy meson exchange and excitation of the $N^*(1440)$ Roper resonance. At the highest CELSIUS energies, with Q above 200 MeV, a preliminary analysis shows that data rather follow flat phase space than simple expectations for a dominant mechanism based on N^* or $\Delta\Delta$ excitations.

At the highest energies, data on $\eta \to \gamma\gamma$ and $\eta \to 3\pi^0$ are regularly collected for control and development of detector performance. These data are also being analysed for detailed studies of η production mechanism. The $\eta \to 3\pi^0$ channel turned out to be cleaner than expected for the pp reaction. The $pp \to pp3\pi^0$ prompt production has not been measured in this energy region. By assumimg flat phase space distributions we obtain an upper limit of 1 μbarn for the total cross section at 1360 MeV.

The experiment is being developed for studies of η charged-particle decays with a final goal to measure rare decays. Deuterium pellets as a target will allow very clean tagging of ηs in the $pd \to {}^3\mathrm{He}\eta$ reaction at threshold. It will also allow studies of meson production in pd, dd and other reactions, one of special interest being the quasi-free pn reaction.

References

[1] http://www.tsl.uu.se/wasa

[2] W. Brodowski et al., Phys. Rev. Lett. **88** (2002) 192301;
J. Johanson et al., Nucl. Phys. **A712** (2002) 75-94;
J. Pätzold et al., Phys. Rev. **C67** (2003) 052202.

[3] L. Alvares-Ruso et al., Nucl. Phys. **A633** (1998) 519-546.

Isospin Violation in Semileptonic Kaon Decays

Helmut Neufeld

Institut für Theoretische Physik der Universität Wien,
Boltzmanngasse 5, A-1090 Wien, Austria.

Current and future experiments on the semileptonic decays of kaons and pions are already sensitive to the following subleading effects:

- strong contributions of order p^6
- isospin-violation of order $(m_d - m_u)p^2, e^2p^2$ generated by the mass difference of the light quarks and by electromagnetism

All these contributions are of comparable size and have to be included in an up-to-data analysis. The appropriate theoretical framework is provided by chiral perturbation theory with virtual photons and leptons [1]. This effective quantum field theory describes the interactions of the pseudoscalar octet, the photon and the light leptons at low energies and allows a comparison of experimental data with the predictions of the standard model. So far, this machinery has been applied to the following problems:

- $K_{\ell 3}$ form factors including electromagnetic contributions to order e^2p^2 [2]
- numerics of K^+_{e3} [2] and K^0_{e3} to $\mathcal{O}(p^6, (m_d - m_u)p^2, e^2p^2)$ [3]
- pionic beta decay [4]

The extraction of the CKM matrix element $|V_{us}|$ from K^+_{e3} decay data may serve as an illustration. The relevant formula is given by [2]

$$|V_{us}| = \frac{16\pi^{3/2}\Gamma(K^+_{e3(\gamma)})^{1/2}}{G_F M_{K^+}^{5/2} S_{\rm EW}^{1/2} |f_+^{K^+\pi^0}(0)| I(\lambda_+)^{1/2}}. \tag{1}$$

Theory has to provide the form factor at zero momentum transfer including all contributions up to $\mathcal{O}(p^6, (m_d - m_u)p^2, e^2p^2)$ [3]:

$$f_+^{K^+\pi^0}(0) = 1.002 \pm 0.010. \tag{2}$$

The error is largely dominated by the uncertainty of the $\mathcal{O}(p^6)$ contribution [5]. The application of Eq. (1) requires a consistent treatment of rleal photon emission in $\Gamma(K^+_{e3(\gamma)})$ (experimental input) and in the phase space integral $I(\lambda_+)$. A possible choice is to accept all pion and positron energies in the whole K^+_{e3} Dalitz plot without any further cut on the photon energy. In this case, radiative corrections reduce $I(\lambda_+)$ by 1.27%.

Recently, a new high statistics measurement of the K^+_{e3} branching ratio has been performed by the E865 Collaboration at Brookhaven [6]. Their analysis of more than 70,000 K^+_{e3} events yielded a branching ratio which was about 2.3σ larger than the current value of the Particle Data Group (PDG 2002). If the E865 result can be confirmed, the CKM unitarity puzzle would be resolved. Inserting the E865 rate in Eq. (1) gives [3]

$$|V_{us}| = 0.2238 \pm 0.0033 \tag{3}$$

being in good agreement with the unitarity of the first row of the CKM matrix:

$$|V_{ud}|^2 + |V_{us}|^2 + |V_{ub}|^2 - 1 = -0.0024 \pm 0.0032. \tag{4}$$

It should be noted, however, that the E865 result does not only differ from older K^+_{e3} data but is also inconsistent with the present K^0_{e3} rate given by PDG 2002 (based again on very old and imprecise data). We propose [3] a rather powerful consistency check of present and future high-precision K^+_{e3} and K^0_{e3} data by considering the ratio

$$r_{+0} = f_+^{K^+\pi^0}(0) \Big/ f_+^{K^0\pi^-}(0). \tag{5}$$

This quantity is largely insensitive to the dominating theoretical uncertainties, in particular the contributions of order p^6. The theoretical prediction [3]

$$r_{+0}^{\rm th} = 1.022 \pm 0.002 - 16\pi\alpha X_1 \tag{6}$$

depends only on the (unknown) electromagnetic low-energy coupling X_1 [1]. Already simple dimensional analysis ($|X_1| \leq 1/(4\pi)^2$) confines r_{+0} to a rather narrow band leading to a stringent test for the observable quantity

$$r_{+0}^{\rm exp} = \left(\frac{2\Gamma(K^+_{e3(\gamma)}) M^5_{K^0} I_{K^0}}{\Gamma(K^0_{e3(\gamma)}) M^5_{K^+} I_{K^+}} \right)^{1/2}. \tag{7}$$

References

[1] M. Knecht, H. Neufeld, H. Rupertsberger and P. Talavera, Eur. Phys. J. C 12 (2000) 469.

[2] V. Cirigliano, M. Knecht, H. Neufeld, H. Rupertsberger and P. Talavera, Eur. Phys. J. C 23 (2002) 121.

[3] V. Cirigliano, H. Neufeld and H. Pichl, in preparation.

[4] l V. Cirigliano, M. Knecht, H. Neufeld and H. Pichl, Eur. Phys. J. C 27 (2003) 255.

[5] P. Post and K. Schilcher, Eur. Phys. J. C 25 (2002) 427;
J. Bijnens and P. Talavera, hep-ph/0303103 v2 (2003).

[6] A. Sher et al., hep-ex/0305042 (2003).

The Anomalous Magnetic Moment of the Muon [1]

Marc Knecht

Centre de Physique Théorique,
CNRS Luminy, Case 907, F-13288 Marseille Cedex 9, France

Over the last few years, the experimental situation concerning the measurement of the anomalous magnetic moment of the muon has witnessed tremendous improvements. The present world average

$$a_{\mu^+}^{\rm exp} \;=\; 11\,659\,203(8)\;[0.7\,{\rm ppm}], \tag{1}$$

is dominated by the latest result from the Brookhaven E821 experiment [1]. This situation has triggered a lot of theoretical activity devoted to the evaluation of the muon g-2 in the standard model, as well as to the contributions of possible degrees of freedom beyond the standard model. Within the standard model, the contributions to a_ℓ, $\ell = e^\pm, \mu^\pm, \tau^\pm$, are conveniently decomposed as $a_\ell^{\rm SM} = a_\ell^{\rm QED} + a_\ell^{\rm had} + a_\ell^{\rm weak}$. The first contribution arises from loops containing virtual leptons and photons only. Including also quark and gluon loops induces the hadronic part $a_\mu^{\rm had}$. Finally, the remaining degrees of freedom of the standard model, neutrinos, electroweak gauge bosons, are responsible for the last piece, $a_\mu^{\rm weak}$.

The contributions to $a_\ell^{\rm QED}$ have been computed analytically up to three loops (for references, and for an introduction to the subject, see [2]). The four loop contribution has been evaluated numerically, while the five loop contribution in the muon case is an estimate only. The results

$$\begin{aligned} a_e^{\rm QED} &= 0.5\left(\tfrac{\alpha}{\pi}\right) - 0.328\,478\,444\,00\left(\tfrac{\alpha}{\pi}\right)^2 + 1.181\,234\,017\left(\tfrac{\alpha}{\pi}\right)^3 - 1.750\,2(38\,4)\left(\tfrac{\alpha}{\pi}\right)^4 \\ a_\mu^{\rm QED} &= 0.5\left(\tfrac{\alpha}{\pi}\right) + 0.765\,857\,399(45)\left(\tfrac{\alpha}{\pi}\right)^2 + 24.050\,509\,5(2\,3)\left(\tfrac{\alpha}{\pi}\right)^3 \\ &\quad +125.08(41)\left(\tfrac{\alpha}{\pi}\right)^4 + 930(170)\left(\tfrac{\alpha}{\pi}\right)^5 \end{aligned} \tag{2}$$

include the recent reevaluation of the four loop calculation [3], which affects $a_e^{\rm QED}$, but only marginally $a_\mu^{\rm QED}$. Using the comparison between the calculation of $a_e^{\rm SM}$ and the experimental value $a_{e^-}^{\rm exp} = 11\,596\,521\,88.3(4.2)$ [3.7 ppb] in order to obtain a determination of the fine structure constant α, then yields $a_\mu^{\rm QED} = 11\,658\,470.35(28) \times 10^{-10}$.

Present limitations in the standard model evaluation of a_μ come from the presence of hadronic contributions which involve the low energy region. These can be decomposed into leading and next-to-leading hadronic vacuum polarization, denoted as $a_\mu^{\rm hvp;1}$ and $a_\mu^{\rm hvp;2}$, respectively, and hadronic light-by-light scattering, $a_\mu^{\rm h\ L\times L}$. The evaluation of the latter relies on specific models, and the present

[1] Work supported in part by TMR, EC contract No. HPRN-CT-2002-00311 (EURIDICE).

situation is summarized by $a_\mu^{\mathrm{h\,L\times L}} = +8(4)\times 10^{-10}$, where the error reflects a conservative estimate of the model dependence. In the case of $a_\mu^{\mathrm{hvp;1}}$, a data based evaluation is possible, using measurements of the cross section for $e^+e^- \to$ hadrons and/or of the hadronic decays of the τ. A new analysis [4] of the CMD-2 data for $e^+e^- \to \pi^+\pi^-$, which give the most important contribution, has resulted in several recent reevaluations, $a_\mu^{\mathrm{hvp;1}}(e^+e^-) = 696.3(7.2)\times 10^{-10}$ [5], $691.8(6.1)\times 10^{-10}$ [6], and $694.8(8.6)\times 10^{-10}$ [7]. While these values are quite compatible, their average gives $a_\mu^{\mathrm{hvp;1}}(e^+e^-) = 694.3(7.5)\times 10^{-10}$, they still show a noticeable and puzzling discrepancy with the evaluation obtained using the τ data, $a_\mu^{\mathrm{hvp;1}}(\tau) = 711.0(5.8)\times 10^{-10}$ [5]. Adding the remaining pieces (see references in [2]), $a_\mu^{\mathrm{hvp;2}} = -10.0(0.6)\times 10^{-10}$, $a_\mu^{\mathrm{weak}} = +15.4(0.3)\times 10^{-10}$, one arrives at the standard model value (the errors are associated to vacuum polarization, hadronic light-by-light scattering, and to QED and weak contributions, respectively)

$$a_\mu^{\mathrm{SM}}(e^+e^-) = 11\,159\,178.1 \pm 7.5 \pm 4.0 \pm 0.4 \tag{3}$$

if only e^+e^- data are used, or

$$a_\mu^{\mathrm{SM}}(\tau) = 11\,159\,194.8 \pm 5.9 \pm 4.0 \pm 0.4 \tag{4}$$

if the τ data are used at low energies. The deviation with respect to the experimental value (1) amounts to 2.2σ or 0.8σ, respectively. Excluding further problems with the data analyses, the only possibility to explain the difference between the e^+e^- and the τ based evaluations seems to be unaccounted for isospin violating corrections. The last paper of [7] presents an interesting discussion in this respect. Future data from KLOE or BABAR will also be helpful.

References

[1] G. W. Bennett *et al.* [Muon g-2 Collaboration], Phys. Rev. Lett. **89**, 101804 (2002) [Erratum-ibid. **89**, 129903 (2002)].

[2] M. Knecht, Lectures given at IUTP41: Flavor Physics, Schladming, Styria, Austria, 22-28 Feb. 2003, arXiv:hep-ph/0307239.

[3] T. Kinoshita and M. Nio, Phys. Rev. Lett. **90**, 021803 (2003).

[4] R. R. Akhmetshin *et al.* [CMD-2 Collaboration], arXiv:hep-ex/0308008.

[5] M. Davier, S. Eidelman, A. Höcker and Z. Zhang, arXiv:hep-ph/0308213.

[6] K. Hagiwara, A. D. Martin, D. Nomura and T. Teubner, talk by T. Teubner at the HEP3002 Conference, Aachen, 17-23 July 2003.

[7] F. Jegerlehner, arXiv:hep-ph/0308117 and arXiv:hep-ph/0310234;
S. Ghozzi and F. Jegerlehner, arXiv:hep-ph/0310181.

Hadronic Atoms

Julia Schweizer

Institute for Theoretical Physics, University of Bern,
Sidlerstr. 5, CH-3012 Bern, Switzerland

Nearly fifty years ago, Deser et al. [1] derived the formula for the width of pionic hydrogen at leading order in isospin symmetry breaking. Similar formulas also hold for pionium and the $\pi^- K^+$ atom, which decay predominantly into $2\pi^l 0$ and $\pi^0 K^0$, respectively. These Deser-type relations allow to extract the scattering lengths from measurements of the decay width and the strong energy shift. The DIRAC collaboration at CERN [2] aims to measure the pionium lifetime to 10% accuracy which allows to determine the S-wave $\pi\pi$ scattering lengths difference $|a_0^0 - a_0^2|$ at 5% precision. The experimental result can then be confronted with the very precise theory prediction $a_0^0 - a_0^2 = 0.265 \pm 0.004$ [3]. Particularly interesting is that one may determine in this manner the nature of the SU(2)× SU(2) chiral symmetry breaking in QCD by experiment. In a new experiment on pionic hydrogen at PSI [4], the pionic hydrogen collaboration plans to measure the strong energy shift of the ground state at 0.2% and the decay width at 1% accuracy. In principle, the πN scattering lengths can then be extracted from data on pionic hydrogen alone. Finally, the DEAR collaboration [5] will measure the strong energy shift and decay width of the $1s$ state in kaonic hydrogen and kaonic deuterium.

In order to determine the scattering lengths from these precision measurements, the theoretical expressions for the decay width and the strong energy shift must be known to an accuracy that matches the experimental precision. The non-relativistic effective Lagrangian framework has proven to be the most efficient method to investigate bound-state characteristics [6]. In particular, this technique allows one to evaluate the higher order corrections to the Deser-type formulae. In the following, we present the result for the decay width of the $\pi^- K^+$ atom at next-to-leading order in isospin symmetry breaking. The non-relativistic πK Lagrangian provides a systematic expansion in powers of the isospin breaking parameter δ, where both the fine-structure constant α and the quark mass difference $m_u - m_d$ count as order δ. The result for the width of the ground state of the $\pi^- K^+$ atom at order $\delta^{9/2}$ yields,

$$\Gamma_{\pi^0 K^0} = 8\alpha^3 \mu_+^2 p^* \mathcal{A}^2 \left(1 + K\right), \quad \mathcal{A} = -\frac{1}{8\sqrt{2}\pi} \frac{1}{M_{\pi^+} + M_{K^+}} \mathrm{Re}\, A_{\mathrm{thr}}^{00;\pm} + o(\delta), \quad (1)$$

with

$$\begin{aligned} p^* &= \left[\frac{M_{\pi^+}\Delta_K + M_{K^+}\Delta_\pi}{M_{\pi^+} + M_{K^+}} - \alpha^2 \mu_+^2 + \frac{(\Delta_K - \Delta_\pi)^2}{4(M_{\pi^+} + M_{K^+})^2} \right]^{\frac{1}{2}}, \\ K &= \frac{M_{\pi^+}\Delta_K + M_{K^+}\Delta_\pi}{M_{\pi^+} + M_{K^+}} {a_0^+}^2 - 4\alpha\mu_+ \left[\ln\alpha - 1\right] (a_0^+ + a_0^-) + o(\delta). \end{aligned} \quad (2)$$

Here μ_+ denotes the charged reduced mass and the meson mass differences read $\Delta_\pi = M_{\pi^+}^2 - M_{\pi^0}^2$, $\Delta_K = M_{K^+}^2 - M_{K^0}^2$. The quantity $\mathrm{Re}\, A_{\mathrm{thr}}^{00;\pm}$ is determined as follows. One evaluates the relativistic $\pi^- K^+ \to \pi^0 K^0$ amplitude at order δ near threshold, see Refs. [7,8] and removes the infrared-divergent Coulomb phase. The real part of this matrix element contains a singularity $\sim 1/|\mathbf{p}|$ at threshold ($\mathbf{p}$ denotes the center of mass momentum of the charged pion and kaon). The constant term in the threshold expansion corresponds to $\mathrm{Re}\, A_{\mathrm{thr}}^{00;\pm}$. Further, the normalization of $\mathcal{A}$ is chosen such that in the isospin symmetry limit it coincides with the isospin odd scattering length a_0^-. The isospin even and odd πK scattering lengths[1] a_0^+ and a_0^- are defined in QCD at $m_u = m_d$ and $M_\pi \doteq M_{\pi^+}$, $M_K \doteq M_{K^+}$.

References

[1] S. Deser, M. L. Goldberger, K. Baumann and W. Thirring, Phys. Rev. **96** (1954) 774.

[2] B. Adeva *et al.*, CERN-SPSLC-95-1; see also J. Schacher, this workshop.

[3] G. Colangelo, J. Gasser and H. Leutwyler, Phys. Lett. B **488** (2000) 261 [arXiv:hep-ph/0007112].

[4] PSI experiment R-98.01, http://pihydrogen.web.psi.ch; see also D. Gotta, this workshop.

[5] S. Bianco *et al.* [DEAR Collaboration], Riv. Nuovo Cim. **22N11** (1999) 1; see also M. Cargnelli, this workshop.

[6] W. E. Caswell and G. P. Lepage, Phys. Lett. B **167** (1986) 437; P. Labelle and K. Buckley, arXiv:hep-ph/9804201; X. Kong and F. Ravndal, Phys. Rev. D **59** (1999) 014031; B. R. Holstein, Phys. Rev. D **60** (1999) 114030 [arXiv:nucl-th/9901041]; D. Eiras and J. Soto, Phys. Rev. D **61** (2000) 114027 [arXiv:hep-ph/9905543]; V. Antonelli, A. Gall, J. Gasser and A. Rusetsky, Annals Phys. **286** (2001) 108 [arXiv:hep-ph/0003118]; J. Gasser, V. E. Lyubovitskij, A. Rusetsky and A. Gall, Phys. Rev. D **64** (2001) 016008 [arXiv:hep-ph/0103157].

[7] B. Kubis and U. G. Meißner, Nucl. Phys. A **699** (2002) 709 [arXiv:hep-ph/0107199].

[8] A. Nehme and P. Talavera, Phys. Rev. D **65** (2002) 054023 [arXiv:hep-ph/0107299].

[1] We use the same notation as in Ref. [7].

Results from DIRAC

Jürg Schacher[1]
for the DIRAC Collaboration

[1]Lab for High Energy Physics, Bern University,
Sidlerstrasse 5, CH-3012 Bern, Switzerland

The $\pi^+\pi^-$ atom or $A_{2\pi}$ is a hydrogen-like atom consisting of π^+ and π^- mesons. This atom decays predominantly by strong interaction into $\pi^0\pi^0$. Hence the $A_{2\pi}$ lifetime is in good approximation inversely proportional to the squared difference between the S-wave $\pi\pi$ scattering lengths for isospin 0 and 2, $|a_0 - a_2|$. This value is predicted in chiral perturbation theory (ChPT). An experimental determination of the $A_{2\pi}$ atom lifetime provides a possibility to check predictions ofl ChPT in a model-independent way.

Using ChPT at next-to-leading order in isospin breaking the $A_{2\pi}$ lifetime is predicted to be [1]: $\tau = (2.9 \pm 0.1) \cdot 10^{-15} s$. A lifetime measurement with 10% accuracy allows to determine the above $\pi\pi$ scattering length difference at 5%.

The $\pi^+\pi^-$ atoms are produced by Coulomb interaction in the final state of $\pi^+\pi^-$ pairs generated in proton–target interactions [2,3]. After production these atoms travel through the target and some of them are broken up due to their interaction with matter: "atomic pairs" are produced, characterized by small pair c.m. relative momenta $Q < 3$ MeV/c. These pairs will be detected in the DIRAC setup. Other atoms annihilate into $\pi^0\pi^0$. The amount of broken up atoms n_A depends on the lifetime which defines the decay rate. Therefore, the breakup probability is a function of the $A_{2\pi}$ lifetime.

In addition, the proton–target interaction produces $\pi^+\pi^-$ pairs with Coulomb ("Coulomb pairs") and without Coulomb final state interaction. The latter category includes pion pairs with one pion from the decay of long-lived resonances ("non-Coulomb pairs") as well as two pions from different interactions ("accidental pairs"). "Coulomb" and "non-Coulomb pairs" together are the so-called "free pairs". The total number of produced atoms (N_A) is proportional to the number of "Coulomb pairs" (N^C) with low relative momenta ($N_A = K \cdot N^C$). The coefficient K is precisely calculable.

DIRAC aims to measure the $A_{2\pi}$ breakup probability: $P_{\rm br}(\tau)$ is defined as ratio of the observed number n_A of "atomic pairs" to the number N_A of produced $\pi^+\pi^-$ atoms calculated from the measured number of "Coulomb pairs".

The purpose of the DIRAC setup at the CERN PS is to record $\pi^+\pi^-$ pairs with small relative momenta [4]. The 24 GeV proton beam hits a thin target (typically 100 μm thick Ni foil). Emerging $\pi^+\pi^-$ pairs travel in vacuum through the upstream spectrometer part with coordinate and ionisation detectors, before they are split by the 2.3 Tm bending magnet into the "positive" and "negative" arm. Both arms are equipped with high precision drift chambers, time of flight

detectors, cherenkov, preshower and muon counters. The relative time resolution between the two arms is around 180 ps.

The momentum reconstruction makes use of the drift chamber information of the two arms as well as of the measured hits in the upstream coordinate detectors. The relative momentum resolution for Q_X and Q_Y is estimated to be 0.4 MeV/c, whereas the Q_L resolution is 0.6 MeV/c. A system of fast trigger processors selects low Q events.

In the analysis the signal extraction of "atomic pairs" from $A_{2\pi}$ ionisation takes advantage of the specific signature of such events, namely time-correlation and very small relative momenta. After rejection of electrons and muons and subtraction of "accidental pairs", the reduced sample of time-correlated low Q events consists of "atomic" and "free pairs" only. The number of "atomic pairs" is found directly from the excess of $\pi^+\pi^-$ pairs with very low relative momenta ($Q < 3$ MeV/c or $|Q_L| < 1.5$ MeV/c) relative to the expected number of "free pairs" simulated using "accidental pairs". Such a $|Q_L|$-distribution of $\pi^+\pi^-$ pairs from nickel data in 2001 is shown (Fig. left). A preliminary analysis of this data sample leads to a breakup probability of $P_{\rm br}(\tau) = 0.46$ with a statistical error of 9%. There are also systematic errors due to the accuracy in describing multiple scattering in the target, membranes and detector planes and also in describing detector responses. Exploiting the dependence of the breakup probability on lifetime (Fig. right) a preliminary estimate for the $\pi^+\pi^-$ atom lifetime yields $\tau = [3.1^{+0.9}_{-0.7}(stat) \pm 1(syst)] \cdot 10^{-15} s$.

Up to now DIRAC has extracted more than 12000 "atomic pairs" and hence will be able to improve substantially the lifetime measurement.





References

[1] J. Gasser, V. E. Lyubovitskij, A. Rusetsky and A. Gall, Phys. Rev. D **64** (2001) 016008.

[2] L. Nemenov, Sov. J. Nucl. Phys. **41** (1985) 629.

[3] L. Afanasyev *et al.*, Phys. Lett. B **308** (1993) 200.

[4] B. Adeva *et al.* [DIRAC Collaboration], hep-ex/0305022 (2003).

PSI Results from Hadronic Atoms

Detlev Gotta

Forschungszentrum Jülich, D-52425 Jülich

In pionic hydrogen (πH) the hadronic interaction manifests itself by a change of the energies and of the natural line width of K X–rays as compared to a purely electromagnetically bound atomic system. Measurement of ground–state transitions allows the determination of the πN s–wave interaction, which is described by the isoscalar and isovector scattering lengths a^+ and a^- [1,2,3]. In addition, from the line broadening Γ_{1s}, which depends only on a^-, the πN coupling constant can be extracted by the Goldhaber–Miyazawa–Ohme sum rule [4].

To improve on the accuracy achieved by previous measurements [5], a thorough study of the atomic cascade is essential. For that purpose, a first series of measurements has been completed by the new pionic–hydrogen experiment at the Paul–Scherrer–Institut (PSI R–98.01 [6]), using the new cyclotron trap, a cryogenic target, a Bragg spectrometer equipped with spherically bent silicon and quartz crystals and a large–area CCD array. Data analysis is in progress.

Except radiative decay, de–excitation during the atomic cascade is due to collisions of the πH system with other atoms of the target, which leads to density effects because of different collision probabilities. In order to identify radiative de–excitation of the πH atom – when bound into complex molecules formed during collisions $\pi^- p + H_2 \rightarrow [(pp\pi^-)p]ee$ [7] – the energy of the πH(3p-1s) transition was measured at various target densities corresponding to a pressure range from 3.5 bar to liquid. X–ray transitions from molecular states should then show up as low–energy satellites with density dependent intensity. The new data do not show any density effect and, consequently, the measured line shift ϵ_{1s} can be attributed exclusively to the strong interaction. The value of $\epsilon_{1s} = 7.120 \pm 0.013$ eV was found to be in good agreement with the result of the previous experiment. Precision was improved by more than a factor of two [8,9].

At present, the accuracy for Γ_{1s} (7%) is limited by the not sufficiently well known correction for the Doppler broadening of the X–ray lines [5]. The broadening is caused by conversion of de–excitation energy into kinetic energy during collisions (Coulomb de–excitation) [10]. For that reason the precisely measured 1s–level shift in pionic deuterium was used together with the shift of hydrogen in the determination of the πN scattering lengths [11]. This procedure, however, requires a sophisticated treatment of the 3–body system πD. In addition, up to now it cannot be excluded that the radiative decay channel after molecule formation is strongly enhanced in deuterium compared to hydrogen.

To study the influence of Coulomb de–excitation on the cascade, the three transitions πH(2p-1s) (2.4 keV), πH(3p-1s) (2.9 keV) and πH(4p-1s) (3.0 keV) were studied at a target density equivalent to 10 bar. An increase of the line width was found for the (2p-1s) line compared to the (3p-1s) transition, which

is attributed to the higher energy release available for the acceleration of the πH system. This result is corroborated by a reduced line width of the (4p-1s) transitionl [8,9]. The response of the crystal spectrometer was obtained here from the π^{12}C(5g-4f) line (3.0 keV), which is negligibly narrow compared to the experimental resolution. From the (4p-1s) line width, a safe upper limit for the 1s–level broadening of $\Gamma_{1s} < 850$ meV is extracted, which is smaller than the result of [5] but still consistent within the errors.

For further improvement, Coulomb de–excitation must be studied in more detail. For this reason a second series of measurements is foreseen starting with a high statistics study of ground–state transitions from muonic hydrogen (μH), where no strong–interaction effects occur. Beforehand, the crystal resolution function, which has to be known with better accuracy than available from a πC spectrum, will be determined with X–rays emitted from helium–like argon, ionised by means of an electron–cyclotron resonance ion trap (ECRIT) presently set up at PSI [13]. With that, studies of the Bragg crystals can be performed with the neccessary statistics within a reasonable time scale. The μH experiment will then be followed by a remeasurement of πH with high statistics.

With the detailed knowledge of the crystal response together with a newly developed cascade code [12], which includes the velocity development during the atomic cascade, a sufficiently accurate correction for the Doppler broadening in pionic hydrogen should be achievable to extract Γ_{1s} at the level of about 1%.

References

[1] V. E. Lyubovitzkij and A. Rusetski, Phys. Lett. B 494 (2000) 9.

[2] J. Gasser et al., Eur. Phys. J. C 26 (2003) 13.

[3] P. Zemp, contribution to this workshop.

[4] M. L. Goldberger, H. Miyazawa and R. Oehme, Phys. Rev. 99 (1955) 986.

[5] H.–Ch. Schröder et al., Eur. Phys. J. C 21 (2001) 433.

[6] PSI experiment R–98.01, http://pihydrogen.web.psi.ch.

[7] S. Jonsell, J. Wallenius, and P. Froelich, Phys. Rev. A 59 (1999) 3440.

[8] D. F. Anagnostopoulos et al., Proc. of the Int. Workshop on Exotic Atoms (EXA 2002), Vienna, Austria, 28-30 November, 2002, (Austrian Academy of Science Press, Vienna, 2003), p. 71.

[9] M. Hennebach, thesis Universität zu Köln, 2003.

[10] A. Badertscher et al., Europhys. Lett. 54 (2001) 313.

[11] T. E. O. Ericson et al., Phys. Rev. C 66 (2002) 014005.

[12] T. Jensen and V. M. Markushin, πN newsletter 16 (2002) 358.

[13] D. F. Anagnostopoulos et al., Nucl. Instr. Meth. B 205 (2003) 9.

The Chiral Unitary Approach

José A. Oller

Departamento de Física, Universidad de Murcia, E-30071 Murcia, Spain.

The effective field theory of low energy QCD is Chiral Perturbation Theory (CHPT). It is a perturbative expansion in powers of the external fourmomenta of the pseudo-Goldstone bosons (the canonical ones being the pions) over a typical hadronic scale, $M_\rho \simeq 4\pi f_\pi \simeq 1$ GeV. Nevertheless, even at low energies there are several examples where such perturbative procedure cannot be applied or the convergence is very slow. The most important ones are the S-waves corresponding to low energy isosclar pion-pion interactions, $\bar{K}N$ scattering close to threshold or nucleon-nucleon interactions. In all the three cases, its non-perturbative nature originates from large unitarity corrections that, indeed, give rise to the appearance of poles in the physical amplitudes: the broad σ resonance with a mass around 450 MeV, the $\Lambda(1405)$ in $\bar{K}N$, the deuteron in ${}^3S_1$ and an antibound state in ${}^1S_0$ just below threshold in NN. Such large unitarity corrections are due to numerical enhancements in $\pi\pi$ scattering [1] and to the large kaon and nucleon masses in $\bar{K}N$ and NN, see ref.[2]. In the last case, there is also a problem associated with the unnatural values of the S-wave NN scattering lengths, much larger than the inverse of the pion mass. On the other hand, CHPT can only be used for rather low energies, although the chiral constraints are beyond the pure perturbative regime since they constitute Ward identities that must be satisfied by the Green functions at any energy. It should be then interesting to dispose of parameterizations valid for higher energies that could apply the chiral constraints in a large energy window, e.g. from threshold up to around 2 GeV in meson-meson, where one could try to match with perturbative+OPE of QCD. When going to higher energies, apart from the right hand cut (unitarity corrections) one should also face the appearance of preexisting resonances (that is, resonances that are not due to the meson-meson self interactions), e.g. the $\rho(770)$ in P-wave $\pi\pi$ scattering, the $\Delta(1232)$ in πN P-waves, etc. Of course, these resonances also contribute to other partial waves but not through their direct exchanges in the s-channel but because of their exchanges in the crossing ones. These facts can be included systematically through the so called Chiral Unitary Approach (UCHPT) [3,4,1]. The general idea is to match *algebraically* the general expression of a partial wave amplitude, or the one of a production process (e.g. a form factor), that takes into account the right hand cut to all orders, with the perturbative expression from CHPT (or alike chiral effective field theories like Heavy Baryon CHPT, Kaplan-Savage-Wise counting, etc). The simplest way to proceed is by performing a dispersion relation for the inverse of the T matrix, since unitarity requires above threshold that $\mathrm{Im}T_{ij}^{-1} = -\rho_i\delta_{ij}$ with $\rho_i = \theta(s-s_{th})\, q/8\pi\sqrt{s}$. Then, performing a dispersion integral along this cut one has,

$$T_{ij}^{-1} = R_{ij}^{-1} + \delta_{ij} \underbrace{\left(g(s_0)_i - \frac{s-s_0}{\pi}\int_{s_{th}}^{\infty} \frac{\rho(s')}{(s'-s-i0^+)(s'-s_0)} ds' \right)}_{g(s)_i\text{: single unitarity bubble}} , \qquad (1)$$

thus $T = [1/R + g]^{-1}$, where $g(s_0)_i$ is a subtraction constant and $1/R$ is the remnant not fixed by the right hand cut. The next step is to fix the unknown input R, or interaction kernel, by matching algebraically with the effective field theory calculation of T. E.g. if we know $T = T_2 + T_4 + \mathcal{O}(p^6)$ for meson-meson scattering then taking g naturally as order p^0 one has, $R_2 = T_2$, $R_4 = T_4 - T_2 g T_2$, and so on. In this way, the interaction kernel R is fixed order by order with increasing precision, with the advantage that eq.(1) can be also applied when the interactions between the hadrons are not perturbative. The perturbation is now performed on R, instead of directly on T, because the latter is affected by the large unitarity corrections. In addition, since many of the effects that limit a straightforward application of CHPT to higher energies are overcome with this procedure, because unitarity is fulfilled to all orders and explicit resonances can also be included as part of R, then this scheme is also applicable to increase the energy range where chiral Lagrangians can be applied. An analogous expression to eq.(1) can also be derived for production mechanisms [5] Many applications of the UCHPT can be found in the literature in meson-meson, meson-baryon [6] and baryon-baryon [3] scattering. Its repercussion in the strong interacting S-waves for all the previous scattering processes has been particularly important and clarifying, providing a reliable and systematic way of procedure for these particularly problematic waves, out of the range of standard vector meson like models for scalars, naive quark models and straightforward applications of large N_c QCD insights. Furthermore, it has also provided the necessary final(initial) state corrections to production mechanisms, e.g. ϕ decays, $\gamma\gamma$ fusion into two mesons, production of exotic resonances, J/Ψ decays, electromagnetic interactions of meson and baryons [7].

References

[1] J. A. Oller and E. Oset, Phys. Rev. D **60** (1999) 074023.

[2] S. Weinberg, Nucl. Phys. B **363** (1991) 3.

[3] J. A. Oller, Nucl. Phys. A **725** (2003) 85.

[4] J. A. Oller and U. G. Meißner, Phys. Lett. B **500** (2001) 263; J. A. Oller and E. Oset, Nucl. Phys. A **620** (1997) 438 (E)-ibid. A **652** (1999) 407.

[5] U. G. Meißner and J. A. Oller, Nucl. Phys. A **679** (2001) 671. J. A. Oller, E. Oset and J. E. Palomar, Phys. Rev. D **63** (2001) 114009.

[6] D. Jido, J. A. Oller, E. Oset, A. Ramos and U. G. Meißner, Nucl. Phys. A **725** (2003) 181 and references therein.

[7] J. A. Oller, Nucl. Phys. A **714** (2003) 161; J. E. Palomar, L. Roca, E. Oset and M. J. Vicente Vacas, arXiv:hep-ph/0306249; J. A. Oller and E. Oset, Nucl. Phys. A **629** (1998) 739; A. P. Szczepaniak, Phys. Lett. B **567** (2003) 23; U. G. Meißner and J. A. Oller, Nucl. Phys. A **679** (2001) 671; E. Oset, E. Marco, J. C. Nacher, J. A. Oller, J. R. Pelaez, A. Ramos and H. Toki, Prog. Part. Nucl. Phys. **44** (2000) 213 and references therein.

Improving the Convergence of Chiral EFT for Few–Nucleon Systems

Evgeny Epelbaum[1], Walter Glöckle[2] and Ulf-G. Meißner[3,4]

[1]Jefferson Laboratory,
Newport News, VA 23606, USA

[2]Institut für Theoretische Physik II, Ruhr–Universitaet Bochum,
D-44780 Bochum, Germany

[3]Universität Bonn, Helmholtz–Institut für Strahlen– und Kernphysik (Theorie),
Nußallee 14 - 16, D-53115 Bonn, Germany

[4]Forschungszentrum Jülich, Institut für Kernphysik (Theorie),
D-52425 Jülich, Germany

In the past decade chiral EFT has been applied by several groups to study the properties of few–nucleon systems based on the original idea of Weinberg [1], see [2] for a recent review. It is of utmost importance for all these applications to verify the convergence of the low–momentum expansion. As pointed out by Kaiser et al. [3], nucleon–nucleon peripheral scattering provides an excellent ground to test the convergence of the chiral expansion. Indeed, the high angular momentum scattering states are dominated by the long–range part of the nuclear force, where chiral dynamics is important. For example, no short–range contact interactions (with, in general, unknown coupling constants) contribute to D– and higher partial waves at next–next–to–leading order (NNLO), which allows for parameter–free calculations. In addition, the interaction between two nucleons becomes weak due to the centrifugal barrier, so that the S–matrix can be obtained using perturbative methods. It was found in [3] that the subleading two–pion exchange (TPE) contribution, which arises at NNLO and depends on the low–energy constants (LECs) $c_{1,3,4}$ shows unphysically strong attraction already at intermediate distances, if the values for these LECs are taken from pion–nucleon scattering. This results in strong disagreement between theoretical predictions and the data for $E_{\rm lab} > 50$ MeV ($E_{\rm lab} > 150$ MeV) for D– (F–) waves and might therefore indicate problems with the convergence of the chiral expansion, see also [4] for more discussion.

In [5] we have investigated the origin of the unphysical attraction of chiral TPE and proposed a new regularization scheme for calculating pion loop integrals, which allows to improve the convergence and is closely related to the long–distance regularization method applied in baryon SU(3) chiral perturbation theory [6]. The proposed scheme makes use of the spectral function representation of the TPE potential

$$V(q) = \frac{2}{\pi} \int_{2M_\pi}^{\infty} d\mu \, \mu \, \frac{\rho(\mu)}{\mu^2 + q^2}, \qquad (1)$$

where the spectral function $\rho(\mu)$ can be obtained via $\rho(\mu) = \Im\,[V(0^+ - i\mu)]$. In [5] we have demonstrated that the short–range high–μ components of the TPE spectral function calculated using dimensional regularization (DR) or infinite momentum cut–off regularization start to dominate the potential already at intermediate distances. For example, the dominant contributions to the isoscalar central part at distances $r \sim M_\pi^{-1}$ arise from the region $\mu \sim 600$ MeV. Chiral EFT predictions for the spectral function can hardly be expected to converge for such and larger values of μ. These spurious short–range components picked up in DR or equivalent regularization schemes have been shown to be responsible for the unphysical behavior of the subleading chiral TPE potential.

Following the lines of [6], the convergence of the chiral expansion can be greatly improved by keeping only truly long–distance portion of the pion loop integrals. This has been achieved introducing the finite cut–off Λ in the spectral function representation eq. (1). The results for nuclear forces obtained using the new regularization scheme are equivalent to the DR ones modulo an infinite set of higher–order contact interactions (this holds true if one uses a sharp cut–off). Both regularization schemes thus lead to exactly the same result for observables provided one keeps terms in all orders in the EFT expansion.

The new regularization scheme has been successfully applied to the two–nucleon scattering problem at NNLO [5,7]. Extensions to N^{3}LO [8] and to systems with more than two nucleons are underway.

References

[1] S. Weinberg, Phys. Lett. B 251 (1990) 288; Nucl. Phys. B 363 (1991) 3.

[2] P.F.Bedaque, U. van Kolck, Ann. Rev. Nucl. Part. Sci. 52 (2002) 339.

[3] N. Kaiser, R. Brockmann, W. Weise, Nucl. Phys. A625 (1997) 758.

[4] E. Epelbaum et al., Eur. Phys. J. A15 (2002) 543.

[5] E. Epelbaum, W. Glöckle, and U.-G. Meißner, Eur. Phys. J. A (2003) in press.

[6] J.F. Donoghue, B.R. Holstein, and B. Borasoy, Phys. Rev. D59 (1999) 036002.

[7] E. Epelbaum, W. Glöckle, and U.-G. Meißner, nucl–th/0308010, to be published in Eur. Phys. J. A.

[8] E. Epelbaum, W. Glöckle, and U.-G. Meißner, in preparation.

Limit Cycle Physics

H.-W. Hammer[1]

[1]Universität Bonn, Helmholtz-Institut für Strahlen- und Kernphysik (Theorie), Nussallee 14-16, 53115 Bonn, Germany

The development of the renormalization group (RG) has had a profound effect on many branches of physics. Its successes range from explaining the universality of critical phenomena in condensed matter physics to the nonperturbative formulation of quantum field theories that describe elementary particles. The RG can be reduced to a set of differential equations that define a flow in the space of coupling constants. Scale-invariant behavior at long distances, as in critical phenomena, can be explained by RG flow to an infrared fixed point. Scale-invariant behavior at short distances, as in asymptotically-free field theories, can be explained by RG flow to an ultraviolet fixed point. However, a fixed point is only the simplest topological feature that can be exhibited by a RG flow. Another possibility is a limit cycle, which is a 1-dimensional orbit that is closed under the RG flow. The possibility of RG flow to a limit cycle was proposed by Wilson in 1971 [1]. However, few physical applications of RG limit cycles have emerged.

In 1970, Efimov discovered a remarkable effect in the 3-body sector for nonrelativistic particles with short-ranged S-wave 2-body interactions [2]. The strength of the interaction is governed by the scattering length a. Efimov showed that if $|a|$ is much larger than the range r_0 of the interaction, there are shallow 3-body bound states whose number increases logarithmically with $|a|/r_0$. In the limit $a \to \pm\infty$, there are infinitely many shallow 3-body bound states with an accumulation point at the 3-body scattering threshold. The ratio of the binding energies of successive states rapidly approaches the universal constant $\lambda_0^2 \approx (22.7)^2$.

The Efimov effect was revisited by Bedaque, Hammer, and van Kolck within the framework of effective field theory (EFT) [3]. The nonperturbative solution of the EFT in the 3-body sector can be obtained by solving integral equations numerically. These integral equations have unique solutions only in the presence of an ultraviolet cutoff Λ. The resulting predictions for 3-body observables, although finite, depend on the cutoff and are periodic functions of $\ln(\Lambda)$. In Ref. [3], it was shown that the EFT could be fully renormalized to remove the residual dependence on Λ in the 3-body sector by adding a 3-body interaction term $g_3(\Lambda)(\psi^*\psi)^3$ to the effective Lagrangian [3]. The dependence of 3-body observables on the cutoff decreases like $1/\Lambda^2$ if $g_3(\Lambda) \propto H(\Lambda)/\Lambda^2$, where $H(\Lambda) = \cos[s_0 \ln(\Lambda/\Lambda_*) + \arctan(s_0)]/\cos[s_0 \ln(\Lambda/\Lambda_*) - \arctan(s_0)]$ with $s_0 \approx 1.006$. With this renormalization, 3-body observables have well-defined limits as $\Lambda \to \infty$, but they depend on the parameter Λ_* introduced by dimensional transmutation. Since $H(\Lambda)$ is a periodic function of $\ln(\Lambda)$, the renormalization of the field theory involves a limit cycle. This contact EFT has also been applied to the 3-nucleon problem [3]. The success of this program suggests that physical

QCD is close to an infrared limit cycle.

We conjecture that QCD can be tuned to the critical RG trajectory for an infrared limit cycle by adjusting the up and down quark masses m_u and m_d [4]. Our argument is based on recent work in which an EFT with explicit pions was used to extrapolate nuclear forces to the chiral limit of QCD [5,6]. In this limit, the masses m_u and m_d of the up and down quarks are zero and the pion is an exactly massless Goldstone boson associated with spontaneous breaking of the chiral symmetry of QCD. According to these chiral extrapolations, the small binding energy 2.2 MeV of the deuteron is a fortuitous consequence of the physical values of m_u and m_d. When extrapolated farther from the chiral limit to slightly larger pion masses, the deuteron's binding energy decreases to zero and then it becomes unbound. In this region, the inverses of both the spin-singlet and spin-triplet scattering lengths $1/a_s$ and $1/a_t$ are close to zero [5,6].

In the next-to-leading order chiral extrapolation of Ref. [6], only quark mass dependent operators proportional to $m_u + m_d$ enter. The extrapolation in m_π can be interpreted as an extrapolation in the sum $m_u + m_d$, with $m_u - m_d$ held fixed. Changing $m_u - m_d$ changes the degree of isospin-symmetry breaking. Since a_s and a_t correspond to different isospin channels, they respond differently to changes in $m_u - m_d$. Therefore it may be possible by tuning both m_u and m_d to make $1/a_t$ and $1/a_s$ vanish simultaneously. This point corresponds to a critical RG trajectory for an infrared limit cycle. At this critical point, the triton has infinitely-many increasingly-shallow excited states with an accumulation point at the 3-nucleon threshold. The ratio of the binding energies of successively shallower states rapidly approaches a constant λ_0^2 close to 515. This opens the exciting possibility of observing an RG limit cycle in QCD using a combination of lattice QCD and chiral EFT.

This talk is based on work done in collaboration with P.F. Bedaque, Eric Braaten, and U. van Kolck.

References

[1] K. G. Wilson, Phys. Rev. D3 (1971) 1818.

[2] V. N. Efimov, Sov. J. Nucl. Phys. 12 (1971) 589; 29 (1979) 546.

[3] P. F. Bedaque, H.-W. Hammer, and U. van Kolck, Phys. Rev. Lett. 82 (1999) 463; Nucl. Phys. A646 (1999) 444; A676 (2000) 357.

[4] E. Braaten and H. W. Hammer, Phys. Rev. Lett. 91 (2003) 102002.

[5] S. R. Beane and M. J. Savage, Nucl. Phys. A717 (2003) 91; A713 (2003) 148.

[6] E. Epelbaum, U.-G. Meißner, and W. Glöckle, Nucl. Phys. A714 (2003) 535.

Nuclear Electroweak Processes in Effective Field Theories

Jiunn-Wei Chen

Department of Physics, National Taiwan University, Taipei 10617, Taiwan

In this talk, I review some selected topics in nuclear electroweak processes using effective field theories (EFT), including applications in astrophysics, nucleon property extractions from nuclear systems and some applications in parton physics.

In nuclear astrophysics, EFT was applied to several very low energy processes such that pions can be treated as heavy particles and integrated out. In Big Bang Nucleosynthesis (BBN), the $np \to d\gamma$ cross section is an important input contributing the largest uncertainty to the primordial abundance rate of ^{7}Li. Using EFT, the theoretical uncertainty is reduced to 1% [1] in the energy range relevant to BBN. The EFT predictions also agree with a very recent measurement of the inverse process in the same energy region [2].

The same theory is also applied to neutrino-deuteron breakup processes which are important inputs for the measurement of solar neutrino flux at Sudbury Neutrino Observatory (SNO). However, there was no precise, direct measurement for those cross sections and two state of the art potential model calculations differ by 5-10%. Using EFT, this difference was successfully described by a parameter $L_{1,A}$, which is an axial two-body current encoding the unknown short distance physics [3]. Since $L_{1,A}$ is the *only* unknown parameter that appears in all the low energy weak interaction deuteron breakup process at the next-to-next-to-leading order, any weak deuteron breakup measurement will constrain $L_{1,A}$. By now, several independent constraints on $L_{1,A}$ have been obtained with consistent results (see [4] for a summary).

Another recent development is the model independent calculations on $NN \to NNX$ in supernova (SN) 1987a [6], where X is some undetected light particle. From the missing energy upper bound of SN1987a, several improved constraints have been obtained with X taken as graviton and dilaton—to probe the size of extra dimensions; as axion, saxion and light neutralinos, and as $\nu\bar{\nu}$ for neutron star cooling [5].

In the extractions of the nucleon polarizabilities from deuteron Compton scattering, ref.[7] used EFT with pion integrated out to obtain the isoscalar electric polarizability $\alpha_N = 7.2 \pm 2.1 \pm 1.6$ and magnetic polarizability $\beta_N = 6.9 \mp 2.1 \mp 1.6$, in units of $10^{-4} fm^3$. This is consistent with $\alpha_N = 9.0 \pm 1.5^{+3.6}_{-0.8}$ and $\beta_N = 1.7 \pm 1.5^{+1.4}_{-0.6}$ found in ref. [8] using EFT with pions plus some higher energy data. However, higher precision Compton scattering data are clearly needed to better determine α_N and β_N.

Recently chiral perturbation theory has been applied to study parton physics through computations of hadronic twist-2 matrix elements [9]. The computed quark mass dependence of twist-2 matrix elements are useful to guide the chiral

extrapolations of lattice data. The method can also be applied to generalized parton distributions, SU(3) flavor structure, large N_c behavior and nuclear modifications (EMC effects) [10]. A recent study [11] in deeply virtual Compton scattering (DVCS) indicates that $\gamma^* N \to \gamma N \pi$ involves twist-2 matrix elements which cannot be determined from $\gamma^* N \to \gamma N$ alone, contrary to previous claims [12].

References

[1] J.W. Chen and M.J. Savage, *Phys. Rev.* C**60** (1999) 065205; G. Rupak, *Nucl. Phys.* A**678** (2000) 405.

[2] W. Tornow et al., *Phys. Lett.* B**574** (2003) 8.

[3] M.N. Butler and J.W. Chen, *Nucl. Phys.* **A675** (2000) 575; M.N. Butler, J.W. Chen, and X. Kong, *Phys. Rev.* C**63** (2001) 035501.

[4] J.W. Chen, K.M. Heeger, and R.G.H. Robertson, *Phys. Rev.* C**67** (2003) 025801.

[5] C. Hanhart, D.R. Phillips, S. Reddy, and M.J. Savage, *Nucl. Phys.* B**595** (2001) 335; D. Arndt and P.J. Fox, *JHEP* 0302 (2003) 036; H.K. Dreiner, C. Hanhart, U. Langenfeld, and D.R. Phillips, *Phys. Rev.* D**68** (2003) 055004.

[6] C.Hanhart, D.R. Phillips, and S. Reddy, *Phys. Lett.* B**499** (2001) 9.

[7] H.W. Griesshammer and G. Rupak, *Phys. Lett.* B**529** (2002) 57; S.R. Beane and M.J. Savage, *Nucl. Phys.* A**694**(2001) 511.

[8] S.R. Beane, M. Malheiro, J.A. McGovern, D.R. Phillips, and U. van Kolck, *Phys. Lett.* B**567** (2003) 200.

[9] D. Arndt and M.J. Savage, *Nucl. Phys.* A**697** (2002) 429; J.W. Chen and X. Ji, *Phys. Lett.* B**523** (2001) 107.

[10] J.W. Chen and X. Ji, *Phys. Rev. Lett.* 87 (2001) 152002; *Phys. Rev. Lett.* 88 (2002) 052003; *Phys. Lett.* B**523** (2001) 73; A.V. Belitsky and X. Ji, *Phys. Lett.* B**538** (2002) 289.

[11] J.W. Chen and M.J. Savage, nucl-th/0308033.

[12] P.A.M. Guichon, L. Mosse, M. Vanderhaeghen, *Phys. Rev.* D**68** (2003) 034018.

Nuclear Physics and Lattice QCD

Silas R. Beane[1,2]

[1]Department of Physics, University of New Hampshire, Durham, NH 03824

[2]Jefferson Laboratory, 12000 Jefferson Avenue, Newport News, VA 23606

Impressive progress is currently being made in computing properties and interactions of the low-lying hadrons using lattice QCD. However, cost limitations will, for the foreseeable future, necessitate the use of quark masses, M_q, that are significantly larger than those of nature, lattice spacings, a, that are not significantly smaller than the physical scale of interest, and lattice sizes, L, that are not significantly larger than the physical scale of interest. Extrapolations in the quark masses, lattice spacing and lattice volume are therefore required. The hierarchy of mass scales is: $L^{-1} \ll M_q \ll \Lambda_\chi \ll a^{-1}$. The appropriate EFT for incorporating the light quark masses, the finite lattice spacing and the lattice size into hadronic observables is χ-PT, which provides systematic expansions in the small parameters $e^{-m_\pi L}$, $1/\ L\Lambda_\chi$, p/Λ_χ, M_q/Λ_χ and $a\Lambda_\chi$. The lattice introduces other unphysical scales as well. Lattice QCD quarks will increasingly be artificially separated into non-degenerate "sea" and "valence" quarks, a procedure known as partial-quenching. Continuum EFT methods are being developed which properly account for finite-a [1,2], finite-L [3,4] and partial-quenching effects [5] and which will allow for a systematic extrapolation from unphysical simulations of hadron properties to nature. Recently, partially-quenched χ-PT (PQχ-PT) has been formulated for baryons [6,7] with finite lattice spacing corrections [8], and many quantities have been computed. (See Ref. [9] and references therein.)

Progress in few-nucleon effective field theory has reached the point where one may meaningfully ask how nuclear energy levels change as we vary the quark masses in the QCD Lagrangian. A practical motivation for understanding the quark mass dependence of nuclear physics is that future lattice QCD results will require extrapolation. Fortunately, the quark-mass dependence of few-nucleon systems can be studied in effective field theory [10,11,12], and amusingly, with existing results, the successful predictions of Big-Bang nucleosynthesis (BBN) can be used to constrain the time dependence of the Higgs vev and Λ_{QCD}, and thereby search for physics beyond the standard model [13][14]. There has been one attempt to compute nucleon-nucleon (NN) scattering parameters in lattice QCD; Ref. [15] computes the ${}^1S_0$ and ${}^3S_1$ scattering lengths in quenched QCD (QQCD) using Lüscher's method [16]. The NN potential is modified in an unphysical way in QQCD and PQQCD [17], however it is straightforward to develop the partially-quenched EFT which matches to a partially-quenched lattice simulation [18]. A second approach to the NN system on the lattice is to study the simplified problem of two interacting heavy-light particles [19]. It has been suggested that lattice QCD simulations of the potential between hadrons containing

a heavy quark will provide insight into the nature of the intermediate-range force between two nucleons [19]. While the NN potential is not itself an observable, one may instead consider heavy systems. The $\Lambda_Q\Lambda_Q$ interaction is an ideal system to investigate [20]. Since the Λ_Q is an isosinglet, there is no OPE, and the leading large-distance behavior is governed by two-pion exchange (TPE), which is physics analogous to the intermediate-range attraction in the NN potential.

References

[1] S.R. Sharpe and R.J. Singleton, Phys. Rev. **D58**, 074501 (1998).

[2] G. Rupak and N. Shoresh, Phys. Rev. **D66**, 054503 (2002).

[3] J. Gasser and H. Leutwyler, Nucl. Phys. **B307**, 763 (1988).

[4] P. Colangelo, S. Dürr and R. Sommer, `hep-lat/0209110`.

[5] S.R. Sharpe and N. Shoresh, Phys. Rev. **D62**, 094503 (2000).

[6] J.-W. Chen and M.J. Savage, Phys. Rev. **D65**, 094001 (2002).

[7] S.R. Beane and M.J. Savage, Nucl. Phys. **A709**, 319 (2002).

[8] S.R. Beane and M.J. Savage, `hep-lat/0306036`.

[9] D. Arndt and B.C. Tiburzi, `hep-lat/0309013`.

[10] S.R. Beane *et al.*, Nucl.Phys. A700 (2002) 377.

[11] S.R. Beane and M.J. Savage, Nucl. Phys. **A713**, 148 (2003); Nucl. Phys. **A717**, 91 (2003).

[12] E. Epelbaum, W. Glöckle and U.-G. Meißner, Nucl. Phys. **A714**, 535 (2003).

[13] J.J. Yoo and R.J. Scherrer, Phys. Rev. **D67**, 043517 (2003).

[14] J.P. Kneller and G.C. McLaughlin, `nucl-th/0305017`.

[15] M. Fukugita *et al.*, Phys. Rev. **D52**, 3003 (1995).

[16] M. Lüscher, Commun. Math. Phys. **105**, 153 (1986).

[17] S.R. Beane and M.J. Savage, Phys. Lett. **B535**, 177 (2002).

[18] S.R. Beane and M.J. Savage, Phys. Rev. **D67**, 054502 (2003).

[19] http://www.jlab.org/~dgr/lhpc/march00.pdf.

[20] D. Arndt, S.R. Beane and M.J. Savage, `nucl-th/0304004`.

The Soft-Collinear Effective Field Theory

Iain W. Stewart

Massachusetts Institute for Technology, Cambridge, MA 02139, USA

The soft-collinear effective theory [1] (SCET) provides a formalism for systematically investigating processes with both energetic and soft hadrons based solely on the underlying structure of QCD. Most effective theories that we are familiar with are designed to separate the physics for hard $p_h^\mu \sim Q$ and soft p_s^μ momenta ($Q^2 \gg p_s^2$). Examples include chiral perturbation theory (ChPT), few-nucleon effective theory, heavy quark effective theory, or the high density effective theory. In SCET we incorporate an additional possibility, namely energetic hadrons where the constituents have momenta p_c^μ nearly collinear to a light-like direction n^μ. Both the energetic hadron and its collinear constituents have $\bar{n} \cdot p_c \sim Q$, where we have made use of light-cone coordinates $(p_c^+, p_c^-, p_c^\perp) = (n\cdot p_c, \bar{n}\cdot p_c, p_c^\perp)$. However, the collinear constituents still have small offshellness $p_c^2 \sim p_s^2$. The process of disentangling the interactions of hard-collinear-soft particles is known as factorization, and is simplified by the SCET framework. I compare some of the basic features of SCET with ChPT in Table 1.

Much like any effective theory the basic ingredients of SCET are its field content, power counting, and symmetries. The Lagrangian and operators,

$$\mathcal{L} = \mathcal{L}^{(0)} + \mathcal{L}^{(1)} + \dots\,, \qquad \mathcal{O} = \mathcal{O}^{(0)} + \mathcal{O}^{(1)} + \dots\,, \tag{1}$$

are organized in a series where only $\mathcal{L}^{(0)}$ and $\mathcal{O}^{(0)}$ are relevant at LO, an additional $\mathcal{L}^{(1)}$ or $\mathcal{O}^{(1)}$ is needed at NLO, etc. The expansion parameter will be $\lambda = \sqrt{\Lambda_{\rm QCD}/Q}$ or $\eta = \Lambda_{\rm QCD}/Q$ depending on whether the collinear fields describe an energetic jet of hadrons or an individual energetic hadron. The effective theory with an expansion in λ is called $\mathrm{SCET_I}$, while the one with an expansion in η is called $\mathrm{SCET_{II}}$. For a review of the field content, power counting and symmetries see [2], for more detail see [4]. A useful advantage of using the symmetries of SCET rather than full QCD is that transformations like P, C, T, baryon number, isospin, etc., can be implemented separately for the soft and collinear sectors.

In its simplest form SCET is applied to hadrons built of light quarks, for detailed examples see [3]. Progress has been made for processes such as exclusive hadronic form factors and deeply virtual Compton scattering [3], as well as deep inelastic scattering and jet production [6]. Many new results have also been obtained for B-decays, where we combine SCET with heavy quark effective theory. Here the large mass of the B meson provides plenty of energy for light hadrons in the final state (cf. [5] for a short review and further references to the literature). Interesting results are also obtained by combining SCET with non-relativistic QCD, an effective field theory for heavy $Q\bar{Q}$ systems [7]. Very recently progress has been made on combining SCET with ChPT [8] to describe light quark mass dynamics in energetic hadrons with an effective field theory. MIT-CTP 3444.

Table 1: A comparison of two effective field theories, chiral perturbation theory (ChPT) and the soft-collinear effective theory (SCET). For notation see [1].

ChPT	SCET
Soft hadrons, pions, nucleons	Soft & Energetic hadrons, quarks, gluons
Nonlinear fields: $\Sigma = \xi^2$ $\Sigma = \exp(2iM/f)$	Nonlinear fields: W, S, Y (Wilson Lines) $W = P\exp(-g/\bar{\mathcal{P}}\,\bar{n}\cdot A_n)$
Chiral transformations: $\Sigma \to L\Sigma R^\dagger$	Gauge transformations: $W \to U_c W$, $W \to U_u W U_u^\dagger$
Labels on fields: N_v	Labels on fields: $h_v^{(b)}$, ξ_n, A_n^μ
Calculable matrix elements	Calculable coefficients
Perturbative loops	Non-perturbative & perturbative loops
Unknown coefficients: L_i, ...	Unknown matrix elements: $\phi_\pi(x)$, $f_{i/p}(\xi)$, ...
Power counting: $\frac{E}{\Lambda}$, $\frac{m_\pi}{\Lambda}$	Power counting: $\frac{\Lambda}{E}$, $\frac{m_\rho}{E}$

References

[1] C.W. Bauer, S. Fleming and M.E. Luke, Phys. Rev. D **63**, 014006 (2001); C.W. Bauer, S. Fleming, D. Pirjol and I.W. Stewart, Phys. Rev. D **63**, 114020 (2001); C.W. Bauer and I.W. Stewart, Phys. Lett. B **516**, 134 (2001); C.W. Bauer, D. Pirjol and I.W. Stewart, Phys. Rev. D **65**, 054022 (2002).

[2] I.W. Stewart, Nucl. Phys. Proc. Suppl. **115**, 107 (2003) [hep-ph/0209159].

[3] C.W. Bauer, S. Fleming, D. Pirjol, I.Z. Rothstein and I.W. Stewart, Phys. Rev. D **66**, 014017 (2002) [arXiv:hep-ph/0202088];

[4] C.W. Bauer, D. Pirjol and I.W. Stewart, Phys. Rev. D **66**, 054005 (2002); Phys. Rev. D **68**, 034021 (2003); J. Chay and C. Kim, Phys. Rev. D **65**, 114016 (2002); M. Beneke, A.P. Chapovsky, M. Diehl and T. Feldmann, Nucl. Phys. B **643**, 431 (2002); R.J. Hill and M. Neubert, Nucl. Phys. B **657**, 229 (2003); A. Hardmeier, E. Lunghi, D. Pirjol and D. Wyler, arXiv:hep-ph/0307171; T. Becher, R.J. Hill and M. Neubert, arXiv:hep-ph/0308122.

[5] I.W. Stewart, arXiv:hep-ph/0308185.

[6] C. W. Bauer, C. Lee, A. V. Manohar and M. B. Wise, arXiv:hep-ph/0309278; A. V. Manohar, arXiv:hep-ph/0309176.

[7] S. Fleming, A. K. Leibovich and T. Mehen, arXiv:hep-ph/0306139.

[8] J.W. Chen and I.W. Stewart, to appear, MIT-CTP 3445.

Chiral Dynamics of Nuclear Matter

S. Fritsch[1], **N. Kaiser**[1] and W. Weise[1,2]

[1]Physik-Dept. T39, TU München, D-85747 Garching, Germany

[2]ECT*, Villa Tambosi, I-38050 Villazzano (Trento), Italy.

In recent years a novel approach to the nuclear matter many-body problem based on effective field theory (in particular chiral perturbation theory) has emerged [1,2]. The key element there is a separation of long- and short-distance dynamics and an ordering scheme in powers of small momenta. At nuclear matter saturation density, $\rho_0 \simeq 0.17\,\mathrm{fm}^{-3}$, the Fermi-momentum k_{f0} and the pion mass m_π are comparable scales ($k_{f0} \simeq 2m_\pi$), and therefore pions must be included as explicit degrees of freedom in the description of the nuclear many-body dynamics. In ref.[2] we have calculated the equation of state of isospin-symmetric nuclear matter in the three-loop approximation of chiral perturbation theory. The contributions to the energy per particle $\bar{E}(k_f)$ as they arise from 1π- and 2π-exchange (closed vacuum) diagrams are ordered in powers of the Fermi-momentum k_f as

$$\bar{E}(k_f) = \sum_{\nu=2}^{5} k_f^\nu \mathcal{F}_\nu(k_f/m_\pi)\,. \tag{1}$$

Note that each expansion coefficient $\mathcal{F}_\nu(k_f/m_\pi)$ for $\nu \geq 3$ is itself a function of the ratio of the two small scales, k_f and m_π, inherent to this calculation. It has been demonstrated that the empirical saturation point, $\bar{E}(k_{f0}) \simeq -15.3\,\mathrm{MeV}$, $\rho_0 \simeq 0.17\,\mathrm{fm}^{-3}$, and the nuclear matter compressibility $K = k_{f0}^2 \bar{E}''(k_{f0}) \simeq 255\,\mathrm{MeV}$ can well reproduced at order $\mathcal{O}(k_f^5)$ in the small momentum expansion with just one single momentum cut-off of $\Lambda = 7f_\pi \simeq 0.65\,\mathrm{GeV}$ which parametrizes the necessary short-range NN-dynamics. The underlying saturation mechanism is surprisingly simple (in the chiral limit $m_\pi = 0$), namely the proper combination of an attractive k_f^3-term and a repulsive k_f^4-term. In calculation, the attractive k_f^3-term comes mainly from iterated 1π-exchange (regulated by a momentum cut-off Λ) whereas the stabilizing k_f^4-term is induced by Pauli-blocking in the medium.

In the same framework, we calculate the density-dependent asymmetry energy $A(k_f)$ and obtain at nuclear matter saturation density ρ_0 the value $A(k_{f0}) \simeq 34\,\mathrm{MeV}$. This prediction is in good agreement with the empirical value of the asymmetry energy $A(k_{f0}) = (33 \pm 4)\,\mathrm{MeV}$. The equation of state of pure neutron matter is also in fair agreement with sophisticated many-body calculations and a resummation result [3] but only for low neutron densities $\rho_n < 0.25\,\mathrm{fm}^{-3}$. The mere fact that pure neutron matter comes out unbound is already non-trivial.

Furthermore, we evaluate the momentum and density dependent complex single-particle potential $U(p, k_f) + i\,W(p, k_f)$ [4] (i.e. the average nuclear mean field). Chiral 1π- and 2π-exchange give rise to a potential depth for a nucleon

at the bottom of the Fermi sea of $U(0,k_{f0}) = -53.2\,\mathrm{MeV}$. This value is in good agreement with the depth of the empirical optical model potential and the nuclear shell model potential. The imaginary single-particle potential $W(p,k_f)$ is generated entirely by iterated 1π-exchange. The half-width of a nucleon-hole state at the bottom of the Fermi sea comes out as $W(0,k_{f0}) = 29.7\,\mathrm{MeV}$. This number is comparable to results of many-body calculations based on the Paris-potential or effective Gogny forces. The basic theorems of Hugenholtz-Van-Hove and Luttinger are satisfied in our perturbative calculation.

Next, we extend this calculation of nuclear matter to finite temperatures [5]. The free energy per particle $\bar{F}(\rho,T)$ is constructed from the interaction kernels by convoluting them with Fermi-Dirac distributions. The calculated pressure isotherms $P = \rho^2\,\partial\bar{F}/\partial\rho$ display the familiar first-order liquid-gas phase transition of isospin-symmetric nuclear matter. The predicted critical point, $T_c \simeq 25.5\,\mathrm{MeV}$ and $\rho_c \simeq 0.09\,\mathrm{fm}^{-3}$, lies however somewhat too high in temperature. This feature originates from the too strong momentum dependence of the real single-particle potential $U(p,k_f)$ near the Fermi-surface $p = k_f$. We consider also pure neutron matter at $T > 0$ and find fair agreement with sophisticated many-body calculations for low neutron densities $\rho_n < 0.2\,\mathrm{fm}^{-3}$.

Finally, chiral 1π- and 2π-exchange determine a nuclear energy density functional of the form [6]:

$$\begin{aligned}\mathcal{E}[\rho,\tau,\vec{J}\,] &= \rho\bar{E}(k_f) + \Big[\tau - \frac{3}{5}\rho k_f^2\Big]\frac{1}{2\widetilde{M}^*(\rho)} \\ &\quad +(\vec{\nabla}\rho)^2 F_\nabla(k_f) + \vec{\nabla}\rho\cdot\vec{J}\,F_{so}(k_f) + \vec{J}^{\,2}\,F_J(k_f)\,, \qquad (2)\end{aligned}$$

with $\rho(\vec{r}\,) = 2k_f^3(\vec{r}\,)/3\pi^2$ the local nucleon density, $\tau(\vec{r}\,)$ the local kinetic energy density and $\vec{J}(\vec{r}\,)$ the local spin-orbit density. This functional (with explicitly density dependent strength functions $F_{\nabla,so,J}(k_f)$) may open a novel approach to nuclear structure calculations which truly start from the long-range pion-induced NN-interaction.

References

[1] M. Lutz, B. Friman and Ch. Appel, Phys. Lett. **B474** (2000) 7.

[2] N. Kaiser, S. Fritsch and W. Weise, Nucl. Phys. **A697** (2002) 255.

[3] J.V. Steele, nucl-th/0010066.

[4] N. Kaiser, S. Fritsch and W. Weise, Nucl. Phys. **A700** (2002) 343.

[5] S. Fritsch, N. Kaiser and W. Weise, Phys. Lett. **B545** (2002) 73.

[6] N. Kaiser, S. Fritsch and W. Weise, Nucl. Phys. **A724** (2003) 47.

Summary of the Working Group I: Goldstone Boson Dynamics

Johan Bijnens[1] and **Ada Farilla**[2]

[1]Dep. Theor. Phys. 2, Lund University,
Sölvegatan 14A, S22362 Lund, Sweden

[2]Dipartimento di Fisica dell'Università "Roma Tre" e Sezione INFN,
Via della Vasca Navale 84, I-00146 Roma, Italy.

The Goldstone Boson working group had a rather extensive program. We had planned 29 presentations of which 13 were experimental and 16 theoretical. Two of the experimental presentations were canceled just before the meeting. The presentations were divided in a series of rather overlapping sessions in particular Kaons, V_{us}-V_{ud}, the Anomaly, Chiral Perturbation Theory (ChPT), hadronic contributions to the muon $g-2$, Roy Equations, the transition from 2 to 3 flavours as well as eta and phi physics. Each section contained experimental and theoretical talks. For more details and references we refer to the speakers own summaries.

Kaons: Here we had two talks on nonleptonic Kaon decays in the framework of ChPT. G. Ecker reported on ongoing work in isospin breaking effects in $K \to 2\pi$ and CP-violation in that system. J. Bijnens discussed the calculation of $K \to 3\pi$ decays in the isospin limit and first results on local isospin breaking. S. Giudici described the new NA48 rare decay data with an emphasis on those testing ChPT predictions.

***V_{us}-V_{ud}*:** This topic was covered in the plenary talk by Neufeld. Here we concentrated on a series of experimental talks regarding both $K_{\ell 3}$ decays and pion beta decay. The new results on $K_{\ell 3}$ were from ISTRA+ (V. Obrazstov) and KEP-PS E246 (V. Anisimovsky) both presenting new measurements of the slopes. The most surprising new result presented was the new BNL E865 measurement of the branching ratio for K^+_{e3} which differs by several sigma from the PDG average and if confirmed would solve the unitarity problem in this sector. E. Frlez showed the contribution from the PIBETA experiment to this puzzle, the branching ratio of pionic beta decay leading to a value of V_{ud} compatible with the PDG one. J. Bijnens presented ChPT results at two loops for $K_{\ell 3}$ in the isospin limit with two main conclusions. The curvature of the $f_+(t)$ form factor might be important and the needed p^6 low energy constants for $f_+(0)$ can be obtained from the slope and curvature of $f_0(t)$.

Anomaly: The anomaly plays an important role in field theory. Its role in the decay $\pi^0 \to \gamma\gamma$ was discussed in the plenary talk by Goity. In the WG M. Moinester described uncertainties on the result for $F_{3\pi}(0)$ resulting from the choice of different form factors in the experimental subtraction towards a possible future COMPASS measurement. The π^0 lifetime experiments and the present status of the JLab experiment PRIMEX were discussed by A. Gasparian.

ChPT issues: M. Büchler presented work on the renormalization group in ChPT at all orders. It was shown how the leading logarithms can in principle be obtained by calculating only one loop diagrams. The scalar form factor of pions and kaons were presented at two loops in ChPT by P. Dhonte, several large corrections were found and a comparison with dispersive results was made. I. Scimemi discussed in detail the problem of defining precisely what are electromagnetic corrections in an effective theory while R. Unterdorfer described how the generating functional method of calculating processes has been extended to the case of three propagators.

$(g-2)_\mu$ hadronic contributions: A major part of the WG was the session devoted to hadronic corrections. G. Colangelo presented the constraints from the $\pi\pi$ phase shift analysis on the ρ contribution. The updated CMD-2 results on e^+e^- →hadrons at low energy were discussed by B. Shwartz while M. Davier gave a general overview of all the hadronic input used and their final results. He also compared the tau decay spectral functions with those from e^+e^-. Finally, F. Nguyen showed the new KLOE analysis for e^+e^- →hadrons using the radiative return method.

Roy Equations: Using dispersion relations and crossing leads to a series of integro-differential equations for amplitudes generically referred to as Roy Equations. P. Büttiker presented a reanalysis of this system of equations in the πK sector. The Roy equations for the $\pi\pi$ system were reanalyzed in great detail by Colangelo et al. (CGL) The precision claimed in this calculation has been challenged by Yndurain and Pelaez. This criticism was presented by F. Yndurain and the reply to these by G. Colangelo. The original precision claimed is very high and thus needs to be checked independently. The situation at present is that the CGL analysis seems to have survived this challenge but the debate is still ongoing.

From two to three Flavours: J. Stern and S. Descotes both discussed aspects of the difference between ChPT for two and three flavours concentrating on the issue of whether disconnected $\bar{s}s$ loops have a large effect. They tested this question using a Bayesian approach in dealing with the input as well as proposed a possible solution to this problem by not rewriting the quark masses at higher order in terms of the meson masses.

η, η' and ϕ decays: On the theory side three talks presented results here. B. Borasoy treated mainly η' decays using a Bethe-Salpeter based resummation scheme and found good agreement with many decays. E. Oset and L. Roca discussed several decays using the chiral unitary approach presented by L. Oller in his plenary talk. E. Oset talked about $\phi \to \pi^0\pi^0\gamma, \pi^0\eta\gamma$ and was very happy with the new KLOE data which agreed much better with his predictions. L. Roca showed how this method describes $\eta \to \pi^0\gamma\gamma$ and the medium dependence of the σ mass. The KLOE eta decay results and the status of many ongoing analysises was presented by B. Di Micco.

Summary of Working Group II: Meson-Baryon Dynamics

R. Beck[1] and T.R. Hemmert[2]

[1]Institut für Kernphysik, Universität Mainz, Becherweg 45, D-55099 Mainz

[2] Theoretische Physik T39, TU München, James-Frank-Str., D-85747 Garching

With 22 presentations from 10 different countries (of which 14 had a theoretical and 8 an experimental focus) this working group on chiral dynamics in the single nucleon sector covered a wide range of topics, attesting to the fact that 9 years after the first Chiral Dynamics conference this continues to be a very active area of research, driven by the close interaction between theory and experiment.

For individual summaries of these 22 contributions we refer to the following pages of these proceedings. Here we only want to highlight a few developments which we expect generate plenty of interesting results in the next few years and are likely to concern this working group at the time of next Chiral Dynamics conference in 2006:

- SU(3) meson-baryon dynamics: New $K^+\Lambda$ and $K^+\Sigma$ data have been shown in this working group for the differential cross section and the hyperon polarization. In the future precise polarization observables (polarized photons and recoil polarization) will become available to separate the different contributions (background, nucleon resonances, ...). Encouraging theoretical progress has been made by different groups utilizing various forms of non-perturbative/unitarization methods. As we expect new precision data in Kaon-photoproduction near threshold in the next few years, it will be a challenge for theory to find out whether there exists a (small?) region of applicability for purely perturbative SU(3) BChPT and at which energy such a framework could be matched onto the existing unitarized/non-perturbative theories of SU(3) meson-baryon dynamics.

- $\Delta(1232)$ resonance region: Over the last few years double polarization experiments have been performed to test the GDH sum rule and the forward spin polarizability. Very precise data were shown for the proton and for the future the experimental focus will be on the neutron. New data on $2\pi^0$ photoproduction off the proton were presented, which allowed for the first time a test of ChPT in the $2\pi^0$ threshold region. In addition a preliminary value for the pion polarizability has been extracted from radiative pion-photoproduction experiment. Compton scattering and pion-photoproduction have to be addressed simultaneously in upcoming theoretical calculations to obtain results in ChEFT which are competitive with existing models. Two different ChEFT frameworks (SSE and δ-expansion) exist and should both be further explored/developed. The non-perturbative

treatment of the width of the resonance in this region does not pose a problem for either framework. Improved calculations of these two scattering processes up into the resonance region will serve as the basis for better theoretical predictions of nucleon dynamical polarizabilities, $N\Delta$-transition form factors etc.

- Pion threshold region: The large discrepancy between ChPT and pion-electroproduction data reported at the Chiral Dynamics workshop 2000 at JLab seems to be resolved. A new measurement has been performed at $Q^2 < 0.1$ GeV^2, which gives significant higher cross sections and a better agreement with ChPT. Existing non-relativistic calculations of pion photo- and electroproduction near threshold are being extended to $\mathcal{O}(p^4)$ in relativistic BChPT. From a theoretical point of view this should shed new light on the still puzzling experimental situation in pion-electroproduction for $Q^2 < 0.1$ GeV^2. From the comparison between relativistic and non-relativistic calculations one also hopes to learn more about the (radius of) convergence of BChPT.

- $\mathcal{O}(p^4)$ BChPT: The role of the large "pole-contributions" identified in forward virtual Compton scattering needs further study. The hope is that a resummation leads to a better-behaved perturbative expansion. A satisfying solution of this problem will be mandatory for any further progress in $\mathcal{O}(p^4)$ ChEFT calculations pertaining to the low-energy spin structure of the nucleon, VCS on the proton, etc.

- Regularization Schemes: A host of schemes and ideas pertaining to regularization in ChEFT has been presented in the working group. We expect that the "effectiveness" in the sense of "simplification of large calculations at high chiral orders" will decide which ideas/proposals are taken up by the community. "Cutoff methods" are likely to play a prominent role in the future as they can serve as a microscope to detect problems of convergence in a chiral series.

New experimental capabilities, polarized beams, polarized targets and full acceptance instruments at the different laboratories will open the possibilities to isolate small components of the cross section via measurements of interference terms. We hope this will result in exciting new data for the topics of Chiral Dynamics.

Finally, we would like to thank all the participants of this working group for their active participation and all speakers for the high quality of their presentations resulted in a stimulating and productive working group. We also thank the organizer of the workshop, Ulf Meißner, as well as the hospitality of the Bonn university. We hope to see all of you again at Duke University in 1000 days!

Summary of Working Group III: Few-Body Dynamics

Paulo Bedaque[1] and Nasser Kalantar-Nayestanaki[2]

[1]LBL,
One Cyclotron Road, Berkeley, California, U.S.A.
[2]KVI,
Zernikelaan 25, Groningen, The Netherlands.

In the theory part of the working group we had the presentation of 12 talks:

1. U. van Kolck, Charge Symmetry Breaking in Pion Production
2. A. Nogga, Probing chiral interactions in light nuclei
3. F. Myhrer, EFT and low energy neutrino deuteron reactions
4. S. Ando, Solar-neutrino reactions on deuteron in EFT* and radiative corrections of neutron beta decay
5. M. Birse, EFT's for the inverse-square potential and three-body problem
6. H. Griesshammer, All orders power counting for the three-nucleon system
7. J. Donoghue, Nuclear Binding Energies and Quark Masses
8. C. Hanhart, Effective field theory treatment for meson production in NN collisions
9. D. Cabrera, Vector meson properties in nuclear matter
10. A. Schwenk, A renormalization group method for ground state properties of finite nuclei
11. L. Platter, The 4-body system in effective field theory
12. H. Krebs, Electroproduction of neutral pions off deuteron near the threshold

The talks aimed at complementing the topics discussed in the plenary sessions. Griesshammer, Birse and Platter talked about the effective theory obtained by integrating out pions and, in particular, the subtle issues arising in three or more nucleon systems. Birse presented a configuration space analysis of the renormalization in the thre-body problem that agrees with the standard power counting. Griesshammer showed how to extend this power counting to arbitrary orders in the low energy expansion and exemplify it with NNLO calculations of neutron-deuteron phase shifts. van Kolck and Hanhart discussed charge symmetry breaking and how the recent and future experiments can help to determine the low energy constants involved. Myhrer, Ando and Krebs used effective field theories and

related methods to make prediction on innelastic processes involving weak current (Myhrer, Ando) and pion production (Krebs). Myhrer advocated that well established models of exchange currents make a prediction for the two-nucleon weak currents with very small uncertanties. Nogga showed the first many-nucleon calculations (through the no-core shell model method) using a potential derived from chiral perturbation theory while Schwenk discussed a renormalization group inspired method to deal with many-body problems. Donoghue presented a way of estimating two-nucleon low energy constants using dispersion relations. Cabrera used a chiral model to discuss vector meson properties in a medium.

For the experimental work, there were a total of 8 experimental talks in the working group. They were given by:

1. T. Black, Precision scattering length measurements
2. St. Kistryn, pd break-up scattering experiments
3. J. Messchendorp, Few-body studies at KVI
4. F. Rathmann, pp scattering experiments
5. H. Sakai, pd elastic scattering experiments
6. H. Schieck, Nd scattering at low energies
7. E. Stephenson, CSB experiment at IUCF
8. W. Tornow, Nd scattering at low energies

A summary of all these talks can be found in these proceedings. The bulk of the presentations concentrated on studying the three-body systems with the aim of understanding possible effects of the three-nucleon forces. It is clear from these measurements that calculations relying only on the two-body forces fail to predict almost all the physical observables at all energies and the need to add a three-nucleon force is present. However, more work needs to be done to formulate three-body forces which can explain the bulk of the data measured with relatively high precision. In order to understand the three-body systems fully, more spin observables should be measured but also the break-up channel in nucleon-deuteron scattering seems to be very promising due to its much richer phase space compared to the elastic channel. Although there is a large data-base for the two-body system, attention should be paid to some spin observables. Bremsstrahlung results in the two-body sector also lack a good theoretical understanding of the process. Measurements at low energies are promising since they could deal with more fundamental quantities which are needed in all the theories. More experimental efforts are underway. The relatively high-precision results from IUCF show very clearly the charge-symmetry breaking. More theoretical work is being performed to describe this experimental observation.

Summary of Working Group III: Hadronic Atoms

Akaki Rusetsky

Universität Bonn, Helmholtz-Institut für Strahlen- und Kernphysik (Theorie), Nußallee 14-16, D-53115 Bonn, Germany

The discussions in the working group were focused on the following questions:

1. Atoms containing strange hadrons. Nemenov (DIRAC coll.) has presented a new proposal to measure the energy-level shifts and the decay widths of both $\pi\pi$ and πK systems in the external magnetic field [1]. The preliminary result for the $\pi\pi$ atom lifetime measurement $\tau = (3.1^{+0.9}_{-0.7}\ (stat) \pm 1.\ (syst)) \cdot 10^{-15}\ s$ has been also reported. The preliminary results for the strong energy shift and the width of the kaonic hydrogen $\epsilon_{1s} = -175 \pm 26 \pm 15\ eV$ and $\Gamma_{1s} = 120 \pm 100 \pm 15\ eV$ were reported by Cargnelli (DEAR coll. [2]). However, as stressed by Gasser, the precise extraction of the KN σ-term and strangeness content of the nucleon by using the DEAR data still poses a big challenge for the theory [3].

2. Isospin-breaking corrections to the energy levels and decay widths of the hydrogen-like hadronic atoms. Zemp has discussed a systematic use of the non-relativistic effective Lagrangian approach to study the decay of the $\pi^- p$ atom in the presence of the electromagnetic $n\gamma$ decay channel. Full calculation of the $O(\alpha, m_d - m_u)$ corrections to the known leading-order result is forthcoming [4]. Selected isospin-breaking corrections in the pionic and kaonic hydrogen have been addressed within the relativistic framework by Ivanov [5].

3. ChPT for heavy nuclei. Using ChPT to systematically study the properties of the deeply bound states of pions has been discussed by Girlanda [6]. Calculation of the pion-nucleus optical potential at $O(p^5)$, with the inclusion of all electromagnetic and strong isospin-breaking effects, has been presented.

References

[1] L.L. Nemenov and V.D. Ovsyannikov, Phys. Lett. B514 (2001) 247.

[2] S. Bianco *et al.* [DEAR Collaboration], Riv. Nuovo Cim. 22N11 (1999) 1.

[3] J. Gasser and M.E. Sainio, arXiv:hep-ph/0002283.

[4] P. Zemp, in progress.

[5] A.N. Ivanov *et al.*, and N.I. Troitskaya, arXiv:nucl-th/0306047.

[6] L. Girlanda, A. Rusetsky, and W. Weise, in preparation.

Summary of Working Group IV: Chiral Lattice Dynamics

or...

Why we need the lattice and why the lattice needs chiral EFT

Elisabetta Pallante[1] and Martin Savage[2]

[1]S.I.S.S.A. and INFN, Sezione di Trieste, Via Beirut 2-4, 34013 Trieste, Italy

[2]Department of Physics, University of Washington Seattle, WA 98195, USA

Lattice QCD is presently the only known rigorous method of computing strong and electroweak matrix elements between hadronic states. This involves formulating QCD on a euclidean lattice with $1/L \ll m_\pi \ll \Lambda_\chi \ll 1/a$ (L is the lattice size, a is the lattice spacing, and Λ_χ the scale of chiral symmetry breaking). Presently, typical parameters for lattice simulations are $L \sim 3$ fm, $a \sim 0.1$ fm and light quark masses $m_q \sim m_{strange}/2$, still quite far from the ideal regime. The goal of many near future simulations is to move toward the $L \to \infty$, $m_q \to m_{u,d}^{phys}$ and $a \to 0$ limits. Deviations from each of the three limits can be perturbatively described in terms of a systematic expansion within an effective field theory (EFT). Recent progress in the optimization of lattice fermion formulations was described by R. Mawhinney, with emphasis on domain wall (DW) fermions. Nice tests of the chiral behavior of overlap fermions were presented by C. Höbling, who reported on recent results for the pseudoscalar mass, the chiral quark condensate, $B_K^{\overline{MS}}(2\,\mathrm{GeV})$, and $B_{7,8}^{3/2}$. Some problems due to operator mixing for particular fermions can be mitigated by a *twisted mass transformation* (Sint). We were all excited to learn that simulations on $\geq$ Tflop machines with largely reduced lattice masses will be feasible during the next year or so. Questions concerning tests of locality for overlap fermions were asked.

For different size lattices two alternative expansions can be implemented: the p-expansion, when $mL \gg 1$ and the ϵ-expansion, for $mL \ll 1$, with the product $m\Sigma V$ kept fixed. As discussed by Chiarappa and Dürr, the two regimes offer alternative ways of constructing hadronic matrix elements and extracting the low-energy constants (LEC) of χPT.

Shoresh stressed that, as partially-quenched QCD (PQQCD) contains QCD when $N_{sea} = 3$ and $m_{sea} = m_{valence}$, one can use $N_{sea} = 3$ simulations with unphysical masses to determine the LEC of QCD. Shoresh also presented results for the Wilson PQχPT lagrangian in the meson sector up to $\mathcal{O}(a^2)$ in the lattice spacing. Fits to PQQCD lattice data (with $N = 2$) using Wilson PQχPT are under study, as presented by Montvay. Very nice results were reported by C. Aubin, using an EFT formulation for staggered fermions. The fit to lattice data with $N = 3$ provides preliminary values of the strong LEC, $L_5 = 1.9(3)\left(^{+6}_{-2}\right)$, $L_4 = 0.3(3)\left(^{+7}_{-2}\right)$, $2L_8 - L_5 = -0.1(1)\left(^{+1}_{-3}\right)$, $2L_6 - L_4 = 0.5(2)\left(^{+1}_{-3}\right)$ and the ratio $f_K/f_\pi = 1.201(8)(15)$, in good agreement with nature.

Kaon weak matrix elements, in particular $K \to \pi\pi$ decays, are a hot topic

for lattice QCD, as ε'/ε provides a crucial test of the Standard Model and its extensions. G. Villadoro explained how (partial) quenching affects in a crucial way $\Delta I = 1/2$ amplitudes. The size of non-factorizable vs factorizable contributions to penguin matrix elements was discussed (S. Peris). They find a value of $\langle Q_8 \rangle^{3/2}$ that is much larger than the lattice value, making the situation still controversial.

Another exciting topic studied in lattice QCD is B meson physics. Quenched lattice data of the $B \to \pi$ transition that is necessary for the extraction of $|V_{ub}|$ was analyzed using QχPT in an effort to estimate systematic errors in the chiral extrapolation (Becirevic). It appears that partially quenched simulations at smaller quark masses are necessary for a more quantitative understanding.

The next few years promise to be a very exciting time for strong interaction physics as we are starting to see lattice calculations of the nucleon properties, such as the magnetic moments, matrix elements of the $n \leq 3$ twist-2 operators, masses, and also the NN scattering lengths. Nucleon form factors were discussed in our working group, with presentations of quenched calculations of the isovector anomalous magnetic moment and charge radii (Göckeler) and the strange form factors (Lewis). Significant work has been accomplished in examining the properties of nucleons in quenched and partially-quenched chiral perturbation theory, QχPT and PQχPT respectively (Beane and Arndt). In addition, first calculations of the $\mathcal{O}(a)$ corrections to nucleon properties arising from both the Sheikholeslami-Wohlert term in the Symanzik action and also from $\mathcal{O}(a)$ corrections to the inserted operators themselves, which are relevant for actions using unimproved Wilson fermions, were presented. The importance of studying the NN scattering lengths on the lattice was stressed by Beane.

Chiral extrapolations have received attention recently. A presentation by Procura on the extrapolation of the nucleon mass and g_A was interesting. One would like to understand much better why some of the forms used in chiral extrapolations can recover the experimental value, despite extrapolations from pion masses that are clearly outside the range of convergence. It was reiterated that results must be regulator independent up to higher order terms in the expansion.

Maarten Golterman lead the working group in a very lively and interesting discussion concerning what one can expect in the next five years and what we should be thinking about. Five areas emerged for scrutiny: finite volume effects, finite lattice spacing effects, staggered fermions, (partial-) quenching and chiral perturbation theory.

In conclusion, the chiral lattice dynamics working group was very stimulating and a good addition to the chiral dynamics meeting. The next five years will see significant progress in numerical lattice simulations, algorithm development and in the development of the appropriate effective field theories. We are entering a time period where rigorous statements about strong interaction observables can be made directly from first principles.

Working Group I: Goldstone Boson Dynamics

Convenors: Johan Bijnens and Ada Farilla

S. Giudici	*NA48 Results on χPT*	68
G. Ecker	*$K \to 2\pi$, ϵ'/ϵ and Isospin Violation*	69
J. Bijnens	*$K \to 3\pi$ in Chiral Perturbation Theory*	70
A.A. Poblaguev	*E865 Results on K_{e3}*	71
V. Obraztsov	*High Statistics Study of the $K^- \to e\nu\pi^0$ and $K^- \to \pi^-\pi^0\pi^0$ Decays with ISTRA+ Setup*	72
V. Anisimovsky	*Measurement of the K_{e3} Form Factors Using Stopped Positive Kaons*	73
J. Bijnens	*K_{l3} at Two Loops in ChPT*	74
E. Frlez	*Pion Beta Decay, V_{ud} and $\pi \to l\nu\gamma$*	75
M.A. Moinester	*A New Determination of the $\gamma\pi \to \pi\pi$ Anomalous Amplitude Via $\pi^- e \to \pi^- e\,\pi^0$ Data*	76
A. Gasparian	*The π^0 Lifetime Experiment and Future Plans at JLab*	77
M. Büchler	*Renormalization Group Equations for Effective Field Theories*	78
P. Dhonte	*Scalar Form Factors to $O(p^6)$ in SU(3) Chiral Perturbation Theory*	79
I. Scimemi	*Hadronic Processes and Electromagnetic Corrections*	80
R. Unterdorfer	*The Generating Functional of Chiral SU(3)*	81
G. Colangelo	*Hadronic Contributions to a_μ below 1 GeV*	82
F. Nguyen	*The Measurement of the Hadronic Cross Section with KLOE Via the Radiative Return*	83
M. Davier	*Hadronic Vacuum Polarization: the μ Magnetic Moment*	84
B. Shwartz	*CMD-2 Results on $e^+e^- \to$ hadrons*	85
P. Büttiker	*Roy–Steiner Equations for πK Scattering*	86
F.J. Ynduráin	*Comments on some Chiral-Dispersive Calculations of $\pi\pi$ Scattering*	87
G. Colangelo	*On the Precision of the Theoretical Calculations for $\pi\pi$ Scattering*	88
J. Stern	*From Two to Three Light Flavours*	89
S. Descotes-G.	*Bayesian Approach to the Determination of N_f=3 Chiral Order Parameters*	90
B. Borasoy	*Hadronic Decays of η and η'*	91
E. Oset	*Φ Radiative Decay to Two Pseudoscalars*	92
B. Di Micco	*η Decays Studies with KLOE*	93
L. Roca	*$\eta \to \pi^0\gamma\gamma$ Decay within a Chiral Unitary Approach*	94

NA48 Results on χPT

Sergio Giudici for Na48 coll.

Scuola Normale Superiore
7, P.za dei Cavalieri, Pisa, Italy

The experiment Na48, located at the CERN SPS accelerator, has originally been designed to precisely measure direct CP violation in the decays of neutral kaon to pions pair. Apart from the important result concerning the CP violation parameter $Re(\epsilon'/\epsilon)$ [1], the experiment investigated some neutral kaon rare decays as well.

- $K_{L,S} \to e^+e^-\pi^+\pi^-$: the branching ratios are measured to be: $BR(K_L \to e^+e^-\pi^+\pi^-) = (3.08 \pm 0.2) \times 10^{-7}$ and $BR(K_S \to e^+e^-\pi^+\pi^-) = (4.69 \pm 0.3) \times 10^{-5}$. The expected adronic-leptonic angle asymmetry for K_L has been measured to be $A_\phi(K_L) = (14.2 \pm 3.6)\%$ [2]

- $K_S \to \gamma\gamma$: the branching ratio is measured to be $BR(K_S \to \gamma\gamma = (2.78 \pm 0.07)$ revealing a 30% discrepancy with respect to χPT at O(p^4) calculation [3].

- $K_{S,L} \to \pi^0\gamma\gamma$: a first observation [4] of the decays $K_S \to \pi^0\gamma\gamma$ has been obtained yielding $BR(K_S \to \pi^0\gamma\gamma)_{(z>0.2)} = (4.9 \pm 1.8) \times 10^{-8}$ in agreement with the χPT prediction [5]. The chiral structure of the vertex need more statistics to be proved. The branching ratio for the K_L is measured to be $BR(K_L \to \pi^0\gamma\gamma) = (1.36 \pm 0.05) \times 10^{-6}$ [6]. The corresponding VMD coupling found is $a_v = -0.46 \pm 0.1$.

- $K_S \to \pi^0e^+e^-$: NA48 observed 7 events publishing [7] the result $BR(K_S \to \pi^0e^+e^-) = 5.8^{+2.8}_{-2.3} \times 10^{-9}$ in a remarkable agreement with the prediction [8]

References

[1] J.R.Batley et al., Phys.Lett. B544 (2002) 97.
[2] A. Lai et al., Eur.Phys.J.C30:33-49,2003.
[3] A. lai et al.,Phys.Lett.B551:7-15,2003.
[4] hep-ex/0309022 , submitted to Phys.Lett.B.
[5] G.Ecker et al., Phys.Lett.B189 (1987) 363.
[6] A. Lai et al.,Phys.Lett.B536:229-240,2002
[7] hep-ex/0309075 , Submitted to Phys.Lett.B
[8] L.M.Sehgal, Nucl. Phys. B19 (1970) 445 ; Nucl. Phys. B291 (1987) 692.

$K \to 2\pi$, ϵ'/ϵ and Isospin Violation

Gerhard Ecker

Institut für Theoretische Physik, Universität Wien
Boltzmanngasse 5, A-1090 Vienna, Austria

This talk is based on joint work with Vincenzo Cirigliano, Helmut Neufeld and Toni Pich [1].

We have performed a complete analysis of isospin breaking in $K \to 2\pi$ amplitudes in chiral perturbation theory, including both strong isospin violation ($m_u \neq m_d$) and electromagnetic corrections to next-to-leading order in the low-energy expansion. The unknown chiral couplings are estimated at leading order in the $1/N_c$ expansion, except for the basic weak couplings G_8, G_{27} that are fitted to the experimental branching ratios. For these couplings we find small deviations from the isospin-limit case.

The main results are:

- Isospin breaking reduces the $\Delta I = 1/2$ ratio $Re\, A_0/Re\, A_2$ by about 10 %.
- Isospin violation in the phases leads to a difference of about 13° between the isospin-limit values extracted from $K \to \pi\pi$ branching ratios and from $\pi\pi$ scattering data [2]. This difference cannot easily be accommodated by the usual uncertainties of large-N_c determinations of low-energy constants.
- Direct CP violation in $K^0 \to \pi\pi$ decays is parametrized by the parameter ϵ'. We locate three different sources of isospin violation for ϵ'. For the overall measure of isospin violation we find

$$\Omega_{\rm eff} = \begin{cases} 0.16 \pm 0.05 & \text{strong IB} \\ 0.06 \pm 0.08 & \text{strong + elm. IB} \end{cases},$$

  depending on whether or not electromagnetic corrections are included in the decay amplitudes.

References

[1] V. Cirigliano, G. Ecker, H. Neufeld and A. Pich, JHEP 0306 (2003) 012; Phys. Rev. Lett. 91 (2003) 162001; Isospin breaking in $K \to \pi\pi$ decays, hep-ph/0310351.

[2] G. Colangelo, J. Gasser and H. Leutwyler, Nucl. Phys. B 603 (2001) 125.

$K \to 3\pi$ in Chiral Perturbation Theory

Johan Bijnens[1], Fredrik Borg[1] (formerly Persson) and Pierre Dhonte[1]

[1]Dep. Theor. Phys. 2, Lund University,
Sölvegatan 14A, S22362 Lund, Sweden

We discuss the treatment of $K \to 3\pi$ in Chiral perturbation theory to one loop [1]. This has earlier been calculated and various relations discussed in [2]. The formulas from that calculation have unfortunately been lost prompting a full new calculation to take into account the new data. The $K \to 2\pi$ amplitudes had already been recalculated in [3] and were confirmed in [1] once more. We found a simple parametrization in terms of one parameter functions for the full amplitude and have compared our full expression to the data.

A full new fit of the Dalitz plot expansion parameters was also done. This fit has no indication of isospin breaking and has an acceptable χ^2. The fit of the ChPT expression to the data has a fairly large χ^2, mainly due to the fact that the various curvatures in the data are poorly fitted. It should be noted that there are large discrepancies between various experiments and that the new data from ISTRA+ have again significantly different curvatures [4]. A full new fit is required to see if the latter would solve some of the discrepancies.

We have embarked on a full calculation of all isospin breaking effects on presented preliminary results on the local isospin breaking, i.e. quark mass effects and local electromagnetic effects. The radiative corrections from photon loops are under further investigation. The indications are that the local isospin breaking effects do not solve the discrepancy with the curvatures, but basically only slightly change the fitted ChPT parameters [5].

The ChPT calculation of [1] has been fully confirmed by [6]. There also a study of the CP asymmetries can be found.

References

[1] J. Bijnens et al, Nucl. Phys. B **648** (2003) 317 [arXiv:hep-ph/0205341].

[2] J. Kambor, J. Missimer and D. Wyler, Phys. Lett. B **261** (1991) 496; J. Kambor et al, Phys. Rev. Lett. **68** (1992) 1818.

[3] J. Bijnens, E. Pallante and J. Prades, Nucl. Phys. B **521** (1998) 305 [arXiv:hep-ph/9801326].

[4] V. Obraztsov, these proceedings.

[5] J. Bijnens and F. Borg, in preparation.

[6] E. Gamiz, J. Prades and I. Scimemi, arXiv:hep-ph/0309172.

E865 Results on K_{e3}

A.A. Poblaguev for the E865 Collaboration

Physics Department, Yale University, New Haven, CT 06511, USA and
Institute for Nuclear Research of Russian Academy of Sciences, Moscow, Russia

E865 [1] at the Brookhaven National Laboratory AGS collected about 70,000 $K^+ \to \pi^0 e^+ \nu$ (K^+_{e3}) decays in flight with the purpose of measuring the K^+_{e3} branching ratio relative to the observed $K^+ \to \pi^+\pi^0$ ($K_{\pi 2}$), $K^+ \to \pi^0\mu^+\nu$ ($K_{\mu 3}$), and $K^+ \to \pi^+\pi^0\pi^0$ ($K_{\pi 3}$) decays [2]. The π^0 in all the decays were identified using the e^+e^- pair from $\pi^0 \to e^+e^-\gamma$ decay and no photon detection was required. The branching ratio was measured to be

$$\mathrm{BR}(K^+_{e3(\gamma)})/[\mathrm{BR}(K_{\pi 2}) + \mathrm{BR}(K_{\mu 3}) + \mathrm{BR}(K_{\pi 3}) = 0.2002 \pm 0.0008_{\mathrm{stat}} \pm 0.0036_{\mathrm{sys}}],$$

where $K^+_{e3(\gamma)}$ includes the effects of virtual and real photons. Using the Particle Data Group (PDG) braching ratios for the normalization decays we obtain

$$\mathrm{BR}(K^+_{e3(\gamma)}) = (5.13 \pm 0.02_{\mathrm{stat}} \pm 0.09_{\mathrm{sys}} \pm 0.04_{\mathrm{norm}})\%$$

This result is about 2.3 standard deviations higher than the current PDG value.

Radiative corrections for the decays inside the K^+_{e3} Dalitz plot boundary were estimated to be -1.3% using the procedure of Ref. [3], while the $K^+_{e3\gamma}$ decays outside the Dalitz plot boundary led to a $+0.5\%$ correction. Using the total radiative correction of -0.8%, the PDG value for the form factor f_+ slope $\lambda_+ = 0.0282 \pm 0.0027$, and the short distance enhancement factor $S_{EW}(M_\rho, M_Z) = 1.0232$ [3], we obtain for the element V_{us} of CKM matrix $|V_{\mathrm{us}} f_+(0)| = 0.2242 \pm 0.0020_{\mathrm{rate}} \pm 0.0007_{\lambda_+}$ which leads to

$$|V_{\mathrm{us}}| = 0.2272 \pm 0.0020_{\mathrm{rate}} \pm 0.0007_{\lambda_+} \pm 0.0018_{f_+(0)}$$

if $f_+(0) = 0.9874 \pm 0.0084$ [3].

With this value of V_{us} and V_{ud} from superallowed nuclear Fermi beta decays we obtain $|V_{\mathrm{ud}}|^2 + |V_{\mathrm{us}}|^2 + |V_{\mathrm{ub}}|^2 = 0.9999 \pm 0.0016$.

While this result is in perfect agreement with CKM unitarity, our K^+_{e3} decay rate disagrees with previous experimental measurements and increases the discrepancy with V_{us} from K^0_{e3} decay if extracted under conventional theoretical assumptions about symmetry breaking.

References

[1] R. Appel et al., Nucl. Instr. Meth. A 479 (2002) 349.

[2] A. Sher et al., Phys. Rev. Lett., in print; hep-ex/0305042.

[3] V. Cirigliano et al., Eur. Phys. J. C 23 (2002) 121.

High Statistics Study of the $K^- \to e\nu\pi^0$ and $K^- \to \pi^-\pi^0\pi^0$ Decays with ISTRA+ Setup

Vladimir Obraztsov

Institute for High energy Physics,
142284 Protvino,Moscow region, Russia

ISTRA+ is a setup which has been operating in a negative unseparated secondary beam of the U-70 PS [1]. The main statistics of the experiment was accumulated during two runs in 2001. A simple trigger, which selected wide class of K^- decays was used. About 700M events were logged on tapes. A sequence of selection cuts results in 550K events of the $K^- \to e\nu\pi^0$ with remaining background of $\leq 1.5\%$ and 252K completely reconstructed events of the $K^- \to \pi^-\pi^0\pi^0$ decay with a background of $\leq 0.05\%$. This is the highest statistics for these decays in the world.

The Dalitz-plot analysis of the first decay was performed in terms of V-A $f_+(t)$ formfactor, which is assumed to be , at most, quadratic function of t: $f_+(t) = f_+(0)(1 + \lambda_+ t/m_\pi^2 + \lambda'_+ t^2/m_\pi^4)$ and exotic scalar (f_S) and tensor (f_T) formfactors which assumed to be constant. Different fits lead to following results:

$$\lambda_+ = 0.0286 \pm 0.0008(stat) \pm 0.0006(syst);\ \lambda''_+ = -0.002^{+0.0031}_{-0.0066};$$
$$f_T/f_+(0) = 0.021^{+0.064}_{-0.075}(stat) \pm 0.026(syst);$$
$$f_S/f_+(0) = 0.002^{+0.020}_{-0.022}(stat) \pm 0.003(syst)$$

Detailed description of the analysis and of the results can be found in [2].

The second decay is analyzed in terms of the "Dalitz plot slopes" g,h,k

$$|A(K^\pm \to 3\pi)|^2 \ \propto\ 1 + g\,Y + h\,Y^2 + k\,X^2 + \ldots\ , \qquad (1)$$

where $X = (s_1 - s_2)/m_\pi^2$ and $Y = (s_3 - s_0)/m_\pi^2$, $s_i = (p_K - p_i)^2$, $s_0 = \frac{1}{3}(s_1 + s_2 + s_3)$, p_K and p_i are the $K^\pm$ and π_i four-momenta (π_3 is the odd pion). The values obtained from the fit are:

$$g = 0.627 \pm 0.004(stat) \pm 0.010(syst),$$
$$h = 0.046 \pm 0.004(stat) \pm 0.012(syst),$$
$$k = 0.001 \pm 0.001(stat) \pm 0.002(syst).$$

The details can be found in [3].

References

[1] I.V.Ajinenko et al., Yad.Fiz **65**(2002),2125.

[2] I. Ajinenko et al., Phys. Lett. **B574**(2003),14.

[3] I. Ajinenko et al., Phys. Lett **B567**(2003),159.

Measurement of the K_{e3} Form Factors Using Stopped Positive Kaons

Valery Anisimovsky

Institute for Nuclear Research RAS, 60th October Revolution prosp., 7a, Moscow, 117312 Russia

(For KEK-PS E246 Collaboration)

The E246 set-up was designed to measure T-violating muon polarization in the $K_{\mu 3}$ decay and consisted of the following main elements: superconducting toroidal spectrometer, the CsI(Tl) photon calorimeter, the muon polarimeter, active target, charged particle detectors, and TOF system [1]. The set-up allowed us to extract K_{e3} events with background fraction $< 0.55\%$ and measure K_{e3} kinematics completely. We were able to extract the natural Dalitz density by deconvolving the convolution equation (radiative corrections were subtracted using formulae of Ref. [2]) and then we determined the form factors by maximizing the logarithmic likelihood between the MC and experimental Dalitz distributions.

We analyzed 102×10^3 K_{e3} events for toroidal field of 0.65T and obtained:

$$\begin{aligned} \lambda_+ &= 0.0278 \pm 0.0017(stat) \pm 0.0016(syst) \\ |f_S/f_+(0)| &= 0.0040 \pm 0.0160(stat) \pm 0.0078(syst) \\ |f_T/f_+(0)| &= 0.019 \pm 0.080(stat) \pm 0.040(syst) \end{aligned}$$

This result is published in [3]. The values of exotic form factors are consistent with zero predicted by the Standard Model and are in a good agreement with ISTRA+ results. The 90% C.L. limits for the form factors are $|f_S/f_+(0)| < 0.033$ and $|f_T/f_+(0)| < 0.166$. The limit for $|f_S/f_+(0)|$ allows us to put a constraint on the two Higgs doublet model [4]: $|\lambda_{us}\lambda_{e\nu}| < 5.2 \times 10^{-3}$. The obtained λ_+ value is consistent with the ChPT prediction 0.031 [5].

References

[1] M. Abe *et al.* [E246 KEK PS Collaboration], Nucl. Instrum. Meth. A **506**, 60 (2003)

[2] E. S. Ginsberg, Phys. Rev. **162**, 1570 (1967)

[3] A. S. Levchenko *et al.* [KEK-PS E246 Collaboration], Phys. Atom. Nucl. **65**, 2232 (2002) [Yad. Fiz. **65**, 2294 (2002)]

[4] J. J. Godina Nava, Phys. Rev. D **53**, 1703 (1996)

[5] L. Maiani, G. Pancheri and N. Paver, "The Second Daphne Physics Handbook. Vol. 1", p.115-190.

$K_{\ell 3}$ at Two Loops in ChPT

Johan Bijnens[1] and Pere Talavera[2]

[1]Dep. Theor. Phys. 2, Lund University,
Sölvegatan 14A, S22362 Lund, Sweden

[2]Departament de Física i Enginyeria Nuclear,
Universitat Politècnica de Catalunya,
Jordi Girona 1-3, E-08034 Barcelona, Spain

We have calculated the decays $K \to \pi\ell\nu$ ($K_{\ell 3}$) to order p^6 in Chiral Perturbation Theory[1]. This calculation has been performed in the isospin limit.

The full discussion can be found in [1] as well as in the two published talks [2]. There are two main conclusions, one experimental and the other theoretical.

The experimental conclusion is that in order to obtain V_{us} with a precision better than 1% it becomes necessary to take the effect of curvature in the form factor $f_+(t)$ into account. This curvature can be predicted from ChPT by using the pion electromagnetic form factor data [3]. This inclusion has also a rather strong effect on the measured slope λ_+, typically moving it outside the error bars quoted on λ_+ from the experiment when assuming a linear dependence on t only.

The main theoretical result relevant for V_{us} is that the p^6 contribution to $f_+(0)$ only depends on two new parameters. These can in turn be determined from the $K\pi$ scalar form factor slope and curvature, thus allowing for a model independent determination of $f_+(0)$. The curvature can be determined from other scalar form factors, as is discussed in [4].

Work on the p^6 isospin breaking contribution is under way.

References

[1] J. Bijnens and P. Talavera, Nucl. Phys. B **669** (2003) 341 [arXiv:hep-ph/0303103].

[2] J. Bijnens, arXiv:hep-ph/0307082, arXiv:hep-ph/0304284.

[3] J. Bijnens and P. Talavera, JHEP **0203** (2002) 046 [arXiv:hep-ph/0203049].

[4] J. Bijnens and P. Dhonte, "Scalar form factors in SU(3) chiral perturbation theory," LU-TP-03-32, arXiv:hep-ph/0307044, these proceedings.

Pion Beta Decay, V_{ud} and $\pi \to \ell\nu\gamma$

Emil Frlež, for the PIBETA Collaboration

Department of Physics, University of Virginia,
Charlottesville, VA 22904-4714, USA

We have used a non-magnetic pure CsI calorimeter (the PIBETA detector [1]) and stopped π^+ beam technique at PSI to collect, over three years, a high statistics data set of (rare) π^+ and μ^+ weak decays. ¿From the background-free data set of $\sim 60,000$ pion beta ($\pi\beta$) decay events we have extracted the preliminary $\pi^+ \to \pi^0 e^+ \nu_e$ branching ratio of

$$\Gamma_{\pi\beta} = (1.038 \pm 0.004(\text{stat}) \pm 0.007(\text{sys})) \times 10^{-8}, \tag{1}$$

by normalizing to the known rate of the $\pi^+ \to e^+\nu_e$ decays [2]. This result implies the quark mixing matrix element $V_{ud} = 0.9737(40)$, in the agreement with the most recent Particle Data Group average $V_{ud} = 0.9734(8)$ [3]. Our experiment tests for the first time the calculation of the $\pi\beta$ radiative corrections which stand at $\text{RC}_{\pi\beta} \sim (+3.3 \pm 0.1)\,\%$ [4].

In the framework of the Standard Model (SM) and the conserved vector current hypothesis (CVC) the radiative pion decay process $\pi^+ \to e^+\nu_e\gamma$ is described by the weak axial and vector form factors, F_A and F_V [5]. By fitting the absolute partial branching ratios deduced from the $\sim 42,000$ radiative events' Dalitz plot we find the parameter $\gamma \equiv F_A/F_V$ to be

$$\gamma_{\text{EXP}} = 0.475 \pm 0.015(\text{stat}) \pm 0.014(\text{sys}), \tag{2}$$

consistent with the present chiral symmetry phenomenology [6].

References

[1] PIBETA Collaboration Home Page, `http://pibeta.psi.ch/~pibeta/`.

[2] W. J. Marciano, Phys. Rev. Lett. 71 (1993) 3629; G. Czapek et al., Phys. Rev. Lett. 70 (1993) 17; D. I. Britton et al., Phys. Rev. D 49 (1994) 28.

[3] K. Hagiwara et al., Phys. Rev. D 66 (2002) 010001.

[4] W. J. Marciano and A. Sirlin, Phys. Rev. Lett. 56 (1986) 22; W. Jaus, Phys. Rev. D 63 (2001) 053009(19); V. Cirigliano, M. Knecht, H. Neufeld, and H. Pichl, Eur. Phys. J C 27 (2003) 255.

[5] D. A. Bryman, P. Depommier, and C. Leroy, Phys. Rep. 88 (1982) 151.

[6] B. R. Holstein, Phys. Rev. D 33 (1986) 3316.

A New Determination of the $\gamma\pi \to \pi\pi$ Anomalous Amplitude Via $\pi^- e \to \pi^- e\, \pi^0$ Data

I. Giller[1], T. Ebertshäuser[2], **M. A. Moinester**[1], S. Scherer[2]

[1] School of Physics and Astronomy, R. and B. Sackler Faculty of Exact Sciences, Tel Aviv University, 69978 Tel Aviv, Israel

[2] Institut für Kernphysik, Johannes Gutenberg-Universität, D-55099 Mainz, Germany

We discuss the reaction $\pi^- e \to \pi^- e \pi^0$ with the purpose of obtaining information on the $\gamma\pi \to \pi\pi$ anomalous amplitude $\mathcal{F}_{3\pi}$. We compare a full calculation at $\mathcal{O}(p^6)$ in chiral perturbation theory and various phenomenological predictions [1] with the existing data of Amendolia *et al* [2]. By integrating our theory results using Monte Carlo techniques and comparing them with the experimental cross section of $\sigma = (2.11 \pm 0.47)$ nb we obtain an experimental value $\mathcal{F}_{3\pi}^{(0)\mathrm{exp}} \approx 10.0 \pm 1.0\ GeV^{-3}$, in good agreement with theoretical expectations. We emphasize the need for new data to allow comparison of experimental and theoretical distributions, and to obtain $\mathcal{F}_{3\pi}^{(0)\mathrm{exp}}$ with smaller uncertainty.

References

[1] B. R. Holstein, Phys. Rev. D **53** (1996) 4099.

[2] S. R. Amendolia *et al.*, Phys. Lett. B **155** 457 (1985) 457.

The π^0 Lifetime Experiment and Future Plans at JLab

A. Gasparian (for the PrimEx Collaboration)

Department of Physics, NC A&T State University, Greensboro, NC USA

The three neutral light pseudoscalar mesons, π^0, η and η', represent one of the most interesting systems in low energy QCD. This system contains fundamental information about the effects of SU(3) and isospin breaking by the u, d, and s quark mass differences, leading to important mixing effects among the mesons, as well as two types of chiral anomalies. Therefore, precision measurements of the two-photon decays of the π^0, η and η' will allow one to determine the light quark mass ratios and a direct extraction of the η, η' mixing angle. Thus, providing a stringent test of the low energy limit of QCD in a relatively clean setting.

Because of the small mass of π^0 meson, the prediction of the chiral anomaly for the $\pi^0 \to \gamma\gamma$ decay width is more accurate and exact in the massless quark limit:

$$\Gamma(\pi^0 \to \gamma\gamma) = \frac{\alpha^2 m_\pi^3}{64\pi^3 F_\pi^2} = 7.725 \text{ eV}$$

However, the real world current-quark masses are of the order of 5-7 MeV. Calculations of the chiral corrections have been performed in the combined framework of chiral perturbation theory (ChPT) and the $1/N_c$ expansion up to $\mathcal{O}(p^6)$ and $\mathcal{O}(p^4 \times 1/N_c)$ in the decay amplitude. It was shown that the next-to-leading order corrections increase the pion decay width by about 4%, with an estimated uncertainty of less than 1%. The PrimEx collaboration at Jefferson Lab is planning to perform a precision measurement of the neutral pion lifetime using the small angle coherent photoproduction of π^0's in the Coulomb field of a nucleus, *i.e.*, the Primakoff effect. This measurement will be a state-of-the-art experimental determination of the lifetime with a precision of less than 1.5% [1]. Currently, we are in the final stages of the preparation of the experiment and expect to have the first data in the Spring of 2004.

With the planned 12 GeV CEBAF energy upgrade, we will be able to perform the second part of this experimental program: radiative width and electromagnetic transition form factor measurements of the more massive η and η' mesons [2]. This comprehensive experimental program will provide fundamental tests of both QCD and QCD inspired models.

(This project is supported under NSF grant PHY-0079840.)

References

[1] Jefferson Lab proposal E02-103 (url:www.jlab.org/primex).

[2] Jefferson Lab 12 GeV proposal (www.jlab.org/div_dept/physics_division/pCDR_public/pCDR_12-1/).

Renormalization Group Equations for Effective Field Theories

M. Büchler [1] and G. Colangelo [2]

[1] Department of Physics,
University of Washington, Seattle, WA 98195-1550, USA

[2] Institut für theoretische Physik,
Universität Bern, Sidlerstr. 5, 3012 Bern, Switzerland

We derive and discuss the renormalization group equations for an arbitrary non-renormalizable quantum field theory [1]. We show that with these equations one can get the structure of the leading divergences at any loop order in terms of sole one loop diagrams. In the framework of chiral perturbation theory, this allows to calculate the series of leading chiral logs by an iteration of one loop calculations. The renormalization group equations can also be used for the calculation of sub(sub,..)-leading divergences; these can be written in terms of one-loop,two-loop,(three-loop,..) diagrams.
Another important issue we discuss is the role of counterterms which vanish at the solution of the equation of motion. Due to the freedom of choosing the couplings of these terms, one can find a basis of counterterms for which all divergences are generated only by one-particle-irreducible diagrams. This simplifies loop calculations considerably.
Finally, we show that the renormalization group equations we obtained apply equally well to renormalizable theories.
With the help of these equations, we calculate the double chiral logs for the weak nonleptonic chiral lagrangian [2].

References

[1] M. Buchler and G. Colangelo, arXiv:hep-ph/0309049.

[2] M. Buchler and G. Colangelo, to be published.

Scalar Form Factors to $O(p^6)$ in SU(3) Chiral Perturbation Theory

Johan Bijnens[1] and **Pierre Dhonte**[1]

[1]Dep. Theor. Phys. 2, Lund University,
Sölvegatan 14A, S22362 Lund, Sweden

We present the results of a full $\mathcal{O}(p^6)$ calculation of all scalar form factors for the eight lightest pseudo-scalar mesons in the framework of $SU(3)$ Chiral Perturbation Theory. This work is part of a larger project aiming at developing and testing $SU(3)$ χPT up to two-loop.

We worked in the isospin limit and focused on the pion and kaon form factors. As a first result we find several loop corrections to be large, as expected from the Feynman-Hellman theorem (see [1]). We also compared these expressions with the behaviour extracted in the dispersive analysis of [2]. Using the procedure developed in [3] for the analysis of K_{l4} and in the limit where the contribution from the $\mathcal{O}(p^6)$ low energy constants (LEC) to the form factors at zero vanishes, the matching of the scalar radii could be used to study the $\mathcal{O}(p^4)$ constants L_4 and L_6 while comparison of the curvatures gave information on the $\mathcal{O}(p^6)$ constants C_{12} and C_{13}. C_{12} is of particular interest for the determination of V_{us} via the K_{l3} study performed in [4]. The precision obtained on this constant was somewhat limited, but we could still obtain new constraints for L_4 and L_6. Thirdly, we extended to the scalar case the relation established by Sirlin in [5] for the vector form factors. This relation is valid at all non-special momentum transfers.

References

[1] J. Bijnens and P.A. Dhonte, *to be published in JHEP* [hep-ph/0307044]

[2] B. Moussallam, Eur. Phys. J. C **14** (2000) 111 [hep-ph/9909292].

[3] G. Amorós, J. Bijnens and P. Talavera, Nucl. Phys. B **585** (2000) 293 [Erratum-ibid. B **598** (2001) 665] [hep-ph/0003258].

[4] J. Bijnens and P. Talavera, Nucl. Phys. B **669** (2003) 341 [arXiv:hep-ph/0303103].

[5] A. Sirlin, Ann. of Phys. 61 (1970) 294, Phys. Rev. Lett. **43** (1979) 904.

Hadronic Processes and Electromagnetic Corrections

Ignazio Scimemi,

ECM-University of Barcelona,
Diagonal 647, E-08028, Barcelona, Spain

The inclusion of electromagnetism in a low energy effective theory is worth further study in view of the present high precision experiments (muon $g-2$, $\pi_0 \to \gamma\gamma$, τ decays, etc. , see also many talks of this conference). In particular in many applications of chiral perturbation theory, one has to purify physical matrix elements from electromagnetic effects. The theoretical problems that I want to point out here are following: the splitting of a pure QCD and a pure electromagnetic part in a hadronic process is model dependent: is it possible to parametrise in a clear way this splitting? What kind of information (scale dependence, gauge dependence,..) is actually included in the parameters of the low energy effective theory? In the recent work of ref.[1] we attempt to answer these questions introducing a possible convention to perform the splitting between strong and electromagnetic parts in some examples. We conclude the following. The splitting that is proposed is done order by order in the loop expansion. The strong part of a quantity depends only on couplings defined in a theory with $e = 0$ (up to the desired perturbative order in e) and it has no running proportional to the electromagnetic coupling e (still, up to the perturbative order in e which is considered). In order to proceed correctly in the construction of an effective theory it is important to characterise the relevant scales of the problem: μ (the renormalisation scale of the underlying theory), μ_{eff} (the renormalisation scale of the effective theory) and μ_1 (the scale at which the strong part of a quantity is defined). The splitting ambiguities are parametrised by the scale μ_1. The uncertainty related to μ_1 can be of numerical relevance as in the pion form factor, $F(\mu_1 = 1\,\text{GeV}) = F(\mu_1 = 500\,\text{MeV}) - 0.1\,\text{MeV}$. The error induced on F by μ_1 is of the order of the PDG error [2]. Another advantage of the proposed splitting, is that in the effective Lagrangian the parameters in the strong sector are expressed through the ones of the underlying theory in its strong sector. This makes the matching between the underlying and the effective theory more transparent. Finally the LECs of the effective theory also contain all information about scale and gauge dependence of the Green functions in the underlying theory with e.m.

References

[1] J. Gasser, A. Rusetsky, I. Scimemi, hep-ph/0305260, to be published in EJP-C.

[2] K. Hagiwara *et al.* [Particle Data Group Collaboration], Phys. Rev. D **66**, 010001 (2002).

The Generating Functional of Chiral SU(3)[1]

Rene Unterdorfer

INFN, Laboratori Nazionali di Frascati,
P.O. Box 13, I 00044 Frascati, Italy.

The generating functional of chiral SU(3) for strong and nonleptonic weak interactions [1,2] was extended to include up to three propagators [3]. An application of the SU(2) version of the functional was the analysis of four-pion production where also resonance exchange was taken into account [4].

Another interesting example is the decay $K_L \to \mu^+\mu^-$ that proceeds through two distinct mechanisms: the long-distance contribution generated by the two-photon exchange and the short-distance part due to W and Z exchange. The long-distance contribution contains one combination of chiral couplings, called χ. Using the long-distance amplitude and the measured branching ratio one can extract information on the Cabibbo-Kobayashi-Maskawa matrix element V_{td} that occurs in the short-distance contribution.

Our aim was to calculate the coupling χ. We applied the leading order large–N_C expression of the $K_L \to \gamma^*\gamma^*$ form factor [5] where we included two terms, the ρ term and one additional term with an effective mass Λ_H. The constants were determined with the help of experimental data on $K_L \to \gamma\, l^+l^-$ and by matching of the large–N_C expression with perturbative QCD [6]. Our final result for the long-distance part of χ is:

$$\chi_{\gamma\gamma}(M_\rho) = 5.8 \pm 1.1$$

References

[1] J. Gasser and H. Leutwyler, Nucl. Phys. B250 (1985) 465.

[2] J. Kambor, J. Missimer and D. Wyler, Nucl. Phys. B346 (1990) 17.

[3] R. Unterdorfer, in preparation.

[4] G. Ecker and R. Unterdorfer, Eur. Phys. J. C24 (2002) 535.

[5] M. Knecht, S. Peris, M. Perottet, E. de Rafael, Phys. Rev. Lett. 83 (1999) 5230.

[6] G. Isidori and R. Unterdorfer, in preparation.

[1]Rene Unterdorfer acknowledges the financial support provided through the European Community's Human Potential Programm under contract HPRN-CT-2002-00311 (EURIDICE).

Hadronic Contributions to a_μ below 1 GeV

Gilberto Colangelo

Institut für Theoretische Physik der Universität Bern

Sidlerstr. 5, 3012 Bern

The hadronic vacuum polarization contribution to a_μ is dominated by the $\pi\pi$ contribution (see, e.g. [1]), which is given by the pion vector form factor. The Omnés representation expresses the latter in terms of its phase which, below 16 M_π^2 exactly, and up to 4 M_K^2 to a good approximation, coincides with the $\pi\pi$ P-wave phase shift. The latter is strongly constrained by analyticity, unitarity and chiral symmetry [2]. We call the Omnés function constructed with the $\pi\pi$ P-wave phase shift $G_1(s)$:

$$G_1(s) = \exp\left[\Delta(s)\right] \qquad \Delta(s) = \frac{s}{\pi}\int_{4M_\pi^2}^{\infty} dx\, \frac{\delta_1^1(x)}{x\,(x-s)} \tag{1}$$

and represent the physical vector pion form factor as a product of three terms:

$$F_\pi(s) = G_1(s)G_2(s)G_\omega(s) \tag{2}$$

where $G_2(s)$ accounts for inelastic effects (i.e. it is the Omnés function constructed with the difference between the phase of the form factor and the P-wave $\pi\pi$ phase shift), and $G_\omega(s)$ takes into account the contribution of the ω through its interference with the ρ. Both functions G_2 and G_ω can be described in terms of a small number of parameters which are fixed by fitting the data.

I have presented preliminary results for a calculation of the hadronic vacuum polarization contribution to a_μ below 1 GeV obtained on the basis of Eq. (2). The analysis is currently in progress and final results will be presented in a forthcoming publication [3].

References

[1] M. Davier, S. Eidelman, A. Hocker and Z. Zhang, arXiv:hep-ph/0308213, and Eur. Phys. J. C **27** (2003) 497 [arXiv:hep-ph/0208177].

[2] G. Colangelo, J. Gasser and H. Leutwyler, Nucl. Phys. B **603** (2001) 125 [arXiv:hep-ph/0103088].

[3] I. Caprini et al., in preparation.

The Measurement of the Hadronic Cross Section with KLOE Via the Radiative Return

Federico Nguyen* on the behalf of the KLOE Collaboration [1]

*Dipartimento di Fisica, Univ. Roma TRE and INFN, Sez. Roma III,
Via della Vasca Navale 84, I-00146 Roma, Italy

Estimates of the hadronic contributions to the magnetic anomaly of the muon, a_μ^{had}, need data from τ decays and the reaction $e^+\,e^- \to$ hadrons. At DAΦNE this latter measurement is done at $\sqrt{s} = M_\phi$ using Initial State Radiation (ISR) events [2]. We measured the differential cross section $d\sigma_{e^+e^-\to\pi^+\pi^-\gamma}/dM^2_{\pi\pi} = \sigma_{e^+e^-\to\pi^+\pi^-}\, H(M^2_{\pi\pi}, \theta^{min}_{\pi\pi})$, where H is the radiation function and $\theta_{\pi\pi}$ is the polar angle of the dipion system. The KLOE detector [3] consists in a large acceptance spectrometer and a fine grained calorimeter, able to provide very good momentum and vertex resolution for pions and electrons.

The measurement of $d\sigma_{e^+e^-\to\pi^+\pi^-\gamma}/dM^2_{\pi\pi}$ consists of the following steps:
1) the detection of two charged tracks with polar angle $50^\circ < \theta < 130^\circ$,
2) the requirement $\theta_{\pi\pi} < 15^\circ$ (or $\theta_{\pi\pi} > 165^\circ$) for enhancing ISR with respect to Final State Radiation (FSR) events,
3) the discrimination of pions from electrons by means of calorimetric variables and the subtraction of $\mu^+\,\mu^-\,\gamma$ and $\pi^+\,\pi^-\,\pi^0$ events by suitable kinematic cuts,
4) the luminosity measurement using Bhabha events ($55^\circ < \theta_{e^\pm} < 125^\circ$), with a systematic uncertainty of $\delta L/L = 0.5\%_{th} \oplus 0.4\%_{exp}$.

After the whole selection we have *ca.* 1.5×10^6 events out of 140.7 pb^{-1} of data collected in year 2001, the experimental systematic error is 1.2%. We derive the $\pi\pi$ contribution to the muon anomaly [2], $a_\mu^{\pi\pi}$, for $M^2_{\pi\pi} \in [0.65, 0.93]$ GeV^2:

$$10^{10} \times a_\mu^{\pi\pi} = 378.4 \pm 0.8_{stat} \pm 4.5_{syst} \pm 3.0_{th} \pm 3.8_{FSR}\ .$$

Our data are in agreement with those from CMD-2 [4], confirming the discrepancy in a_μ^{had} using $e^+\,e^-$ or τ data.

References

[1] A. Aloisio *et al.* [KLOE Collaboration], arXiv:hep-ex/0307051.

[2] S. Binner, J. H. Kühn and K. Melnikov, Phys. Lett. B **459** (1999) 279.

[3] M. Primavera, these proceedings.

[4] B. A. Shwartz, these proceedings.

[1] The full authors' list can be found in [1].

[2] These numbers supersede the results presented at this Workshop:
$10^{10} \times a_\mu^{\pi\pi} = 374.1 \pm 1.1_{stat} \pm 5.2_{syst} \pm 3.0_{th}\,^{+7.5}_{-0}|_{FSR}$.

Hadronic Vacuum Polarization: the μ Magnetic Moment

Michel Davier

Laboratoire de l'Accélérateur Linéaire,
IN2P3-CNRS et Université de Paris-Sud, 91898 Orsay, France

A new evaluation of the hadronic vacuum polarization contribution to the muon magnetic moment is presented [1,2]. We take into account the reanalysis of the low-energy e^+e^- annihilation cross section into hadrons by the CMD-2 Collaboration [3]. The agreement between e^+e^- and τ spectral functions in the $\pi\pi$ channel is found to be much improved. Nevertheless, significant discrepancies remain in the center-of-mass energy range between 0.85 and 1.0 GeV, so that we refrain from averaging the two data sets. The values found for the lowest-order hadronic vacuum polarization contributions are

$$a_\mu^{\rm had,LO} = \begin{cases} (696.3 \pm 6.2_{\rm exp} \pm 3.6_{\rm rad})\ 10^{-10} & [e^+e^-{\rm -based}]\ , \\ (711.0 \pm 5.0_{\rm exp} \pm 0.8_{\rm rad} \pm 2.8_{\rm SU(2)})\ 10^{-10} & [\tau{\rm -based}]\ , \end{cases}$$

where the errors have been separated according to their sources: experimental, missing radiative corrections in e^+e^- data, and isospin breaking. The corresponding Standard Model predictions for the muon magnetic anomaly read

$$a_\mu = \begin{cases} (11\,659\,180.9 \pm 7.2_{\rm had} \pm 3.5_{\rm LBL} \pm 0.4_{\rm QED+EW})\ 10^{-10} & [e^+e^-{\rm -based}]\ , \\ (11\,659\,195.6 \pm 5.8_{\rm had} \pm 3.5_{\rm LBL} \pm 0.4_{\rm QED+EW})\ 10^{-10} & [\tau{\rm -based}]\ , \end{cases}$$

where the errors account for the hadronic, light-by-light (LBL) scattering and electroweak contributions. The deviations from the measurement at BNL are found to be $(22.1 \pm 7.2 \pm 3.5 \pm 8.0)\ 10^{-10}$ (1.9 σ) and $(7.4 \pm 5.8 \pm 3.5 \pm 8.0)\ 10^{-10}$ (0.7 σ) for the e^+e^-- and τ-based estimates, respectively, where the second error is from the LBL contribution and the third one from the BNL measurement [4].

Preliminary results are given on the radiative return process $e^+e^- \to \gamma\, 2\pi^+2\pi^-$ measured with BABAR. They are shown to significantly improve the contribution of the 4π channel to the muon magnetic anomaly.

References

[1] M. Davier, S. Eidelman, A. Höcker, Z. Zhang, *Eur.Phys.J.* **C27** (2003) 497.

[2] M. Davier, S. Eidelman, A. Höcker, Z. Zhang, hep-ph/0308213, to appear in *Eur.Phys.J.*.

[3] R. Akhmetshin *et al.* (CMD-2 Collaboration), hep-ex/0308008 (2003).

[4] G. W. Bennett *et al.* (Muon g-2 Collaboration), *Phys. Rev. Lett.* **89** (2002) 101804; Erratum-ibid. **89** (2002) 129903.

CMD-2 Results on $e^+e^- \to$ hadrons

Boris Shwartz[1] for CMD-2 collaboration

[1] Budker Institute of Nuclear Physics SB RAS
630090 Novosibirsk, Russia

This report presents the results obtained by the CMD-2 collaboration in the years 2002-2003. The CMD-2 [1] detector had been collecting data from 1992 to 2000 at VEPP-2M collider in the energy range from 0.4 to 1.4 GeV. Although the data collection was finished more than two years ago, the data processing is continued providing the physics yield.

The new results were obtained for the several processes [2,3,4]. Here, we mention briefly the results which are not published yet.

The process $e^+e^- \to \pi^0 e^+e^-$ was studied around the $\omega-$meson peak. The measured value of $Br(\omega \to \pi^0 e^+e^-) = (8.7 \pm 0.9 \pm 0.5) \times 10^{-4}$ has the best accuracy and it's in agreement with the previous experimental result as well as with theoretical predictions.

Study of the $\phi \to \pi^+\pi^-\pi^0$ decay dynamic. The fit of the Dalitz-plot distribution with the probability density function including the $\rho\pi$ final state amplitude and three pion point like amplitude, $A_n a e^{i\varphi} + A_{\rho\pi}$, gives the values $a = 0.103 \pm 0.028$ in a good agreement with [5] and $\varphi = -2.0 \pm 0.3$ that is rather different from the mentioned paper.

Very important subject studied at CMD-2 is the total hadron cross section in the available energy range. Recently, some uncertainty in the code for the calculation of the radiative corrections for $e^+e^- \to e^+e^-$ was found. Since this process is used for the luminosity measurement, the total cross section changed by 2-3% after reanalysis [6]. The most intriguing value of the hadron vacuum polarization contribution to muon (g-2) value became closer to theoretical one, $a_\mu(exp) - a_\mu(th) = (22.1 \pm 11.3) \times 10^{-10} (1.9\sigma)$.

References

[1] G.A.Aksenov et al., Preprint Budker INP 85-118, Novosibirsk, 1985.
R.R.Akhmetshin et al., Preprint Budker INP 99-11, Novosibirsk, 1999.

[2] R.R. Akhmetshin, et al., Phys.Lett.**B551** (2003) 27.

[3] R.R. Akhmetshin et al., Phys.Lett. **B562** (2003) 173.

[4] R.R. Akhmetshin et al.,hep-ex/0310012.

[5] A.Aloisio et al., Phys.Lett. **B561** (2003) 55.

[6] R.R. Akhmetshin, et al., hep-ex/0308008.

Roy–Steiner Equations for πK Scattering

Paul Büttiker[1], Sebastien Descotes–Genon[2], Bachir Moussallam[3]

[1] HISKP, Universität Bonn, D–53115 Bonn, [2] LPT, Université Paris–Sud, F-91406 Orsay, [3] IPN, Université Paris–Sud, F-91406 Orsay

πK scattering is the most simple $SU(3)$–process involving strange quarks. It is therefore an ideal place to test chiral predictions with non–vanishing strangeness. While this process is interesting by itself, a detailed knowledge of πK scattering also contributes to the understanding of the flavour dependence of the order parameters of Chiral Perturbation Theory (ChPT) [1]. In our work we use Roy–Steiner equations to analyze the available experimental πK and $\pi\pi \to K\bar{K}$ data and to explore the low energy region where no such data are available [2].

Taking the higher partial waves and all S– and P–waves above ≈ 1 GeV as input, we have found unique solutions for the πK S–waves and the $I = 1/2$ P–wave in the energy region from threshold to ≈ 1 GeV. Our solutions for the low–energy region turn out to be in general poor agreement with the available experimental data. Hence, the mass for the $K^*(892)$ is shifted by ≈ 10 MeV from the published value to 905 MeV. The solutions of the Roy–Steiner equations also yield predictions for the scattering lengths $a_0^{1/2}$ and $a_0^{3/2}$. In contradiction to $\pi\pi$ scattering the allowed values are strongly constrained by the data, so that there is no universal band for the two scattering lengths. We find $m_\pi a_0^{1/2} = 0.224 \pm 0.022, \quad m_\pi a_0^{3/2} = -0.0448 \pm 0.0077$ Finally, the matching of the subthreshold parameters in the chiral and the dispersive framework yields the following estimates for the low–energy constants (at the scale $\mu = m_\rho$ in units of 10^{-3}): $L_1^r = 1.05 \pm 0.12, L_2^r = 1.32 \pm 0.03, L_3^r = -4.53 \pm 0.14, L_4^r = 0.53 \pm 0.39$ and $2L_6^r + L_8^r = 3.66 \pm 1.52$.



Fig: 1–σ error ellipse for the scattering lengths $a_0^{1/2}$ and $a_0^{3/2}$.

References

[1] J. Stern, these proceedings; S. Descotes–Genon, L. Girlanda and J. Stern, JHEP **0001** (2000) 041, Eur. Phys. J. C **27** (2003) 115

[2] B. Ananthanarayan and P. Büttiker, Eur.Phys.J. C**19** (2001) 517; B. Ananthanarayan, P. Büttiker and B. Moussallam, Eur.Phys.J. C**22** (2001) 133; P. Büttiker, S. Descotes–Genon, B. Moussallam,hep-ph/0310283

Comments on some Chiral-Dispersive Calculations of $\pi\pi$ Scattering

F. J. Ynduráin

Departamento de Física Teórica, C-XI
Universidad Autónoma de Madrid,
Canto Blanco,
E-28049, Madrid, Spain

In two recent papers, Ananthanarayan, Colangelo, Gasser and Leutwyler and Colangelo, Gasser and Leutwyler (to be referred to as, respectively, ACGL and CGL) have used experimental information, analyticity and unitarity (in the form of the Roy equations) and, in CGL, chiral perturbation theory, to construct the $\pi\pi$ scattering amplitude at low energy, $s^{1/2} \leq 0.8$ GeV.

We will mainly discuss here the second paper, CGL, in which an outstanding precision (at the percent level) is claimed but which, unfortunately, presents a number of drawbacks. First of all, the input scattering amplitude at high energy ($s^{1/2} \geq 1.42$ GeV) which ACGL, CGL use is not physically acceptable; and it is also clear that the errors these authors take for some of their experimental input data (in particular, for the phase shift $\delta_0^{(0)}$ at 0.8 GeV) are excessively optimistic. What is more, the CGL threshold parameters, and the low energy S0, S2 and P waves below 0.8 GeV, are displaced from what one gets by direct fits to experimental data, disagree with what other authors (Descotes et al., and Kamiński, Leśniak and Loiseau) find using also Roy equations, and fail to pass a number of consistency tests. Among these last, and by 2 to 4 σ, we find failure of nonsubtracted forward dispersion relations for the amplitude with isospin 1 excahange (Olsson's sum rule), inconsistency of the D wave scattering length for $\pi^0\pi^0 \to \pi^+\pi^-$ scattering, and disagreement of the P wave effective range parameter, b_1 with results from the pion form factor. Moreover, we remark that some chiral parameters used by CGL are probably displaced from their correct values. As a consequence of this, we conclude that the error estimates of CGL are too optimistic.

The details of all this, based on a recent paper by the author and J. R. Peláez, may be found in the report by the author, "Comments on some chiral-dispersive calculations of $\pi\pi$ scattering", FTUAM 03-14 (hep-ph/0310206).

On the Precision of Theoretical Calculations for $\pi\pi$ Scattering

Gilberto Colangelo

Institut für Theoretische Physik der Universität Bern

Sidlerstr. 5, 3012 Bern

In a recent publication [1], Peláez and Ynduráin have raised objections against our work on Roy equations [2] and the subsequent matching between this representation and the chiral one, which led to remarkably sharp predictions for the $\pi\pi$ scattering amplitude at low energy [3]. In particular, they claim that the asymptotic representation we used is incorrect and that this affected all our results. This latter conclusion was based on an indirect reasoning: they have evaluated some of the low energy observables of $\pi\pi$ scattering and obtained flat disagreement with our earlier results. In their opinion this disagreement had to be attributed to the asymptotic representation for the $\pi\pi$ amplitude we had used. In this talk I have shown that:

1. if we take the asymptotic representation proposed by Peláez and Ynduráin the solution of the Roy equations barely changes;
2. the indirect reasoning that led them to citicize our results is flawed – there is a logical leap between the observed discrepancies among the two calculations and the conclusion that the asymptotics we used is incorrect;
3. the asymptotic representation proposed by Peláez and Ynduráin is in many respects worse than the one we used. In particular, it does not follow from factorization applied to data on high energy total cross sections of proton–proton and pion–proton.

A full account of our answer to the criticism in [1] can be found in [4].

References

[1] J. R. Pelaez and F. J. Yndurain, Phys. Rev. D **68** (2003) 074005 [arXiv:hep-ph/0304067].

[2] B. Ananthanarayan, G. Colangelo, J. Gasser and H. Leutwyler, Phys. Rept. **353** (2001) 207 [arXiv:hep-ph/0005297].

[3] G. Colangelo, J. Gasser and H. Leutwyler, Nucl. Phys. B **603** (2001) 125 [arXiv:hep-ph/0103088].

[4] I. Caprini, G. Colangelo, J. Gasser and H. Leutwyler, Phys. Rev. D **68** (2003) 074006 [arXiv:hep-ph/0306122].

From Two to Three Light Flavours

Jan Stern

Groupe de Physique Theorique, IPN, 91405 Orsay Cedex, France

In $N_f = 2$ ChPT one expands in powers of m_u and m_d keeping m_s fixed at its physical value, whereas $N_f = 3$ ChPT treats all three light quark masses m_u, m_d and m_s as expansion parameters. The corresponding $N_f = 2$ and $N_f = 3$ chiral limits are characterized by different order parameters: QCD suggests paramagnetic inequalities $\Sigma(3) < \Sigma(2)$ and $F^2(3) < F^2(2)$, where $\Sigma(N_f)$ denotes the $N_f = 2$ or $N_f = 3$ chiral limits of the condensate $-\langle \bar{u}u \rangle$ and $F^2(N_f)$ stands for the corresponding chiral limits of F_π^2. Hence, on general grounds, one expects a suppression of the two fundamental three-flavour order parameters relative to their two-flavour analogs. This effect is due to massive strange sea quark pairs which induce extra positive contributions to $\Sigma(2)$ and $F^2(2)$. The latter vanish for $m_s = 0$ i.e. they do not show up in $\Sigma(3)$ and $F^2(3)$. In practice,the resulting difference between $N_f = 2$ and $N_f = 3$ order parameters may be important since $m_s \sim \Lambda_{QCD}$ and furthermore, due to fluctuations of the lowest eigenvalues of Euclidean Dirac operator, QCD vacuum enjoys OZI-rule violating and large N_c suppressed correlations between scalar massive $\bar{s}s$ and massless non strange $\bar{u}u + \bar{d}d$ pairs [1] .In ChPT , this phenomenon is encoded in the values of the large N_c suppressed LEC's $L_6^r(\mu)$ and $L_4^r(\mu)$. Dominant effects of vacuum fluctuations of massive $\bar{s}s$ pairs are described by the scale independent combinations $\Delta L_6 = L_6^r(M_\rho) + 0.26 \times 10^{-3}$ and $\Delta L_4 = L_4^r(M_\rho) + 0.51 \times 10^{-3}$ which should remain positive due to the paramagnetic inequalities. They are usually multiplied by m_s and often appear with large numerical coefficients. Consequently, rather small positive values of ΔL_6 and/or ΔL_4 (close to their upper bounds of order 0.001 and 0.002 respectively which reflect the positivity of $\Sigma(3)$ and of $F^2(3)$) could be sufficient to produce an important suppression of the three-flavour condensate and/or of the decay constant and to destabilize the perturbative relation between order parameters and Goldstone boson observables [2] . Such instability need not affect the standard two-flavour ChPT characterized by a large condensate $\Sigma(2)$ and it need not reflect a too large value of m_s but rather an important OZI-rule violating fluctuation of vacuum $\bar{s}s$ pairs.

References

[1] S. Descotes-Genon, L. Girlanda and J. Stern, JHEP 0001 (2000) 041 [hep-ph/9910537]; Eur. Phys. J. C 27 (2003) 115 [hep-ph/0207337].

[2] S. Descotes-Genon, N.H. Fuchs, L. Girlanda and J. Stern [hep-ph/0311120]

Bayesian Approach to the Determination of N_f=3 Chiral Order Parameters

Sébastien Descotes-Genon

Laboratoire de Physique Théorique, 91405 Orsay Cedex, France.

Due to its light mass of order $\Lambda_{\rm QCD}$, the strange quark can play a special role in Chiral Symmetry Breaking (χSB). Differences in the pattern of χSB in the limits $N_f = 2$ ($m_u, m_d \to 0$, m_s physical) and $N_f = 3$ ($m_u, m_d, m_s \to 0$) may arise due to vacuum fluctuations of $s\bar{s}$ pairs in relation to the Zweig-rule violation in the scalar sector [1]. The latter is in particular described by the $O(p^4)$ low-energy constants L_4 and L_6 of the strong chiral lagrangian. Recent estimates of these constants [2,3] suggest sizeable fluctuations of $s\bar{s}$ pairs, which could lead to a significant suppression of the $N_f = 3$ quark condensate and pseudoscalar constant with respect to their $N_f = 2$ counterparts.

In the case of large fluctuations, the customary treatment of $SU(3) \times SU(3)$ chiral expansions generates instabilities upsetting their convergence [4]. We develop a systematic program to cure these instabilities by resumming nonperturbatively vacuum fluctuations of $s\bar{s}$ pairs, in order to extract information about χSB from experimental observations even in the presence of large fluctuations. The resulting Bayesian framework is exploited to analyse the low-energy $\pi\pi$ scattering amplitude [5] and to obtain a lower bound on the quark mass ratio $2m_s/(m_u + m_d) \geq 14$ at 95% confidence level. On the other hand, $\pi\pi$ scattering does not constrain the quark condensate or the pseudoscalar decay constant in the $N_f = 3$ chiral limit.

References

[1] S. Descotes-Genon, L. Girlanda and J. Stern, JHEP **0001** (2000) 041; Eur. Phys. J. C **27** (2003) 115.

[2] B. Moussallam, Eur. Phys. J. C **14** (2000) 111; JHEP **0008** (2000) 005.

S. Descotes-Genon, JHEP **0103** (2001) 002.

[3] B. Ananthanarayan, P. Büttiker and B. Moussallam, Eur. Phys. J. C **22** (2001) 133.

P. Büttiker, S. Descotes-Genon and B. Moussallam, hep-ph/0310283.

[4] S. Descotes-Genon, N. H. Fuchs, L. Girlanda and J. Stern, in preparation.

[5] S. Descotes-Genon, N. H. Fuchs, L. Girlanda and J. Stern, Eur. Phys. J. C **24** (2002) 469.

Hadronic Decays of η and η'

B. Borasoy

Physik Department, TU München,
85747 Garching, Germany

The hadronic decays $\eta \to \pi\pi\pi$, $\eta' \to \pi\pi\pi$ and $\eta' \to \eta\pi\pi$ are investigated within a $U(3)$ chiral unitary approach [1]. To this end, the most general chiral effective Lagrangian which describes the interactions between members of the lowest lying pseudoscalar nonet (π, K, η, η') up to fourth chiral order is utilized and the pertinent tree diagram amplitudes are derived. Final state interactions are included by deriving the effective s-wave potentials for meson meson scattering from the chiral effective Lagrangian and iterating them in a Bethe-Salpeter equation. With only a small set of parameters we are able to explain both rates and spectral shapes of these decays.

The decay $\eta \to \pi\pi\pi$ may be used to constrain the double quark mass ratio $Q^2 = \frac{m_s - \hat{m}}{m_d - m_u} \frac{m_s + \hat{m}}{m_d + m_u}$ with $\hat{m} = m_u + m_d$ from which we obtain $Q = 23.4 \pm 0.8$, a value which is in agreement with the requirement by Dashen's theorem and indicates that higher order corrections to this low-energy theorem may be small.

We can furthermore confirm the importance of the $a_0(980)$ resonance for $\eta' \to \eta\pi\pi$—as has been claimed in the literature—without including it explicitly in the theory.

The presented method can also be applied in the study of the decays $\eta/\eta' \to \gamma\gamma$ [2], $\eta/\eta' \to \pi^+\pi^-\gamma$ and $\eta/\eta' \to \pi^0\gamma\gamma$.

References

[1] N. Beisert and B. Borasoy, Nucl. Phys. **A716** (2003) 186.

[2] B. Borasoy and R. Nißler, hep-ph/0309011, EPJA (in print).

Φ Radiative Decay to Two Pseudoscalars

J. Palomar, L. Roca, **E. Oset** and M.J. Vicente Vacas

Department of Theoretical Physics and IFIC, University of Valencia, Spain

We study the radiative ϕ decay into $\pi^0\pi^0\gamma$ and $\pi^0\eta\gamma$ taking into account mechanisms in which there are two sequential vector-meson-pseudoscalar or axial-vector-vector-pseudoscalar steps followed by the coupling of a vector meson to the photon, considering the final state interaction of the two mesons [1]. There are other mechanisms in which two kaons are produced through the same sequential mechanisms or from ϕ decay into two kaons and then undergo final state interaction leading to the final pair of pions or $\pi^0\eta$. The latter mechanism is the leading one but the other mechanisms provide appreciable contributions. The results of the parameter free theory together with the theoretical uncertainties, are compared with the latest experimental KLOE results at Frascati [2] in the two figures below, which also show the theoretical errors. It was shown during the Conference that the data for $\pi^0\pi^0\gamma$ decay around 500 MeV have changed and are now in agreement with the theory.





References

[1] J. E. Palomar, L. Roca, E. Oset and M. J. Vicente Vacas, arXiv:hep-ph/0306249.

[2] A. Aloisio et al.[KLOE Collaboration] Phys. Lett. B 537 (2002) 21

η Decays Studies with KLOE

Biagio Di Micco† (on the behalf of the KLOE collaboration)[1]

†Dipartimento di Fisica, Università degi Studi di Roma Tre and INFN, Sez. Roma III, Via della Vasca Navale 84, I-00146 Roma, Italy

We present the status of the analyses on η meson decays with the KLOE detector at DAΦNE Φ factory[1]. The KLOE detector has collected about 20 milions of η events, produced by the $\Phi \to \eta\gamma$ decay process. Here we present the status of the analyses of the decay process: $\eta \to 3\gamma$, $\eta \to \pi^+\pi^-\pi^0$, $\eta \to 3\pi^0$, $\eta \to \pi^0\gamma\gamma$ and $\eta \to \pi^+\pi^-\gamma$.

The $\eta \to 3\gamma$ decay is a C violating decay. The SM prediction of the branching ratio is of the order of 10^{-12}[2], far below the present experimental limit (5×10^{-4} [3]), so any measurement of a larger branching ratio would be a clear signal of new physics. KLOE has analyzed 17×10^6 η decays [4]. No signal has been found, so we have evaluated an upper limit: $Br(\eta \to 3\gamma) < 1.6 \times 10^{-5}$ (95 % C.L.).

The $\eta \to \pi^0\gamma\gamma$ is interesting for testing ChPT to O(p^6) level [2]. We expect to have from 5000 to 14000 such decays in our data[3][2]. The main background is the $\eta \to 3\pi^0$ channel when there is a merging of the clusters in the calorimeter.

The Dalitz plot analysis of the $\eta \to \pi^+\pi^-\pi^0$ channel is interesting for C violation and for ChPT testing [2]. We have analized about 400000 $\eta \to 3\pi^0$ events, producing the following very preliminary results for the Dalitz plot parameters: (a,b,c) = $(-1.05 \pm 0.01, 0.20 \pm 0.03, -0.009 \pm 0.009)$ (only statistical error reported). The c parameter is compatible with 0, so no C violation is observed.

For the $\eta \to 3\pi^0$ the overall selection efficiency is 16%. So we expect to have a sample of 1 million events with the full KLOE statistic. The analysis reproduces well the Dalitz plot parameter put in the MC generator.

The $\eta \to \pi^+\pi^-\gamma$ is an interesting decay to check the box anomaly [2] studying the energy distribution of the γ. We have selected 18508 among 1 million of η events with a preliminary analysis.

References

[1] M.Primavera, these proceedings.

[2] J. Bijnens, G. Faldt and B. M. Nefkens, *Eta Phisics Handbook* Proceedings of workshop in Uppsala, 25-27 Oct, 2001, Physica Scripta T99 (2002)

[3] K. Hagiwara et al., Physical Review D**66**, 010001 (2002)

[4] The KLOE collaboration, hep-ex/0307042

[1] See [4] for the complete authors list

$\eta \to \pi^0\gamma\gamma$ Decay within a Chiral Unitary Approach

E. Oset[1], J. R. Peláez[2] and **L. Roca**[1]

[1]Departamento de Física Teórica and IFIC
Institutos de Investigación de Paterna, 22085, 46071, Valencia, Spain

[2] Departamento de Física Teórica II, Universidad Complutense.
28040 Madrid, Spain.

The $\eta \to \pi^0\gamma\gamma$ decay has attracted much theoretical attention since, in Chiral Perturbation Theory (ChPT), the calculation at $O(p^2)$ vanishes and at $O(p^4)$ is very small. The first sizeable contribution comes at $O(p^6)$. This makes this reaction to be, in principle, a good test of ChPT at $O(p^6)$ and higher. But the parameters involved in the ChPT Lagrangians at these higher orders are not easy to determine. On the other hand models using Vector Meson Dominance (VMD) have been used, but expanding the vector meson propagator to obtain the $O(p^6)$ chiral coefficients gives results $\sim 1/2$ of those obtained by keeping the full vector meson propagator which is supposed to contain higher orders. This implies that $O(p^8) > O(p^6)$. This makes us to conclude that a strict chiral counting has to be abandoned and we have to think in terms of 'relevant mechanisms' instead of 'diagrams contributing up to a certain order'.

At this point is where unitary extensions of ChPT (UChPT) can help to solve the problem. UChPT is based on the implementation of unitary techniques to resum loops and has allowed to extend the predictions of ChPT up to $\sim 1.2\,GeV$ even generating many resonances dynamically without including them explicitly.

Considering this background, we improve the calculations of the $\eta \to \pi^0\gamma\gamma$ decay within the context of meson chiral lagrangians. We use a chiral unitary approach for the meson-meson interaction, thus generating the $a_0(980)$ resonance and fixing the longstanding sign ambiguity on its contribution. This also allows us to calculate the loops with one vector meson exchange, thus removing a former source of uncertainty. In addition we ensure the consistency of the approach with other processes. First, by using vector meson dominance couplings normalized to agree with radiative vector meson decays. And, second, by checking the consistency of the calculations with the related $\gamma\gamma \to \pi^0\eta$ reaction. We find an $\eta \to \pi^0\gamma\gamma$ decay width of 0.47 ± 0.10 eV, in clear disagreement with published data but in remarkable agreement with the most recent measurement.

References

[1] E. Oset, J. R. Pelaez and L. Roca, Phys. Rev. D **67** (2003) 073013 [arXiv:hep-ph/0210282].

Working Group II: Meson-Baryon Dynamics

Convenors: Reinhard Beck and Thomas R. Hemmert

Nucleon Spin Structure / GDH

D. Drechsel — *GDH, Nucleon Spin Structure and Chiral Dynamics: Theory* — 96

A. Braghieri — *Helicity Dependence of Pion Photo-Production and the GDH Sum Rule* — 97

Real Compton Scattering / VCS

H. Fonvieille — *Virtual Compton Scattering at Low Energy and the Generalized Polarizabilities of the Nucleon* — 98

R. Hildebrandt — *Dynamical Polarizabilities from (Polarized) Nucleon Compton Scattering* — 99

L. Fil'kov — *Radiative Pion Photoproduction and Pion Polarizability* — 100

Strangeness Production / SU(3)

K.-H. Glander — *Strangeness Production from ELSA* — 101

B. Raue — *Strangeness Production at Jefferson Lab* — 102

E. Oset — *Unitarized Chiral Dynamics: SU(3) and Resonances* — 103

Pion (Electro-) Production and Chiral Dynamics

H. Merkel — *Threshold Pion Electroproduction: Update on Experiments* — 104

M. Kotulla — *Two π^0 Production at Threshold* — 105

S.N. Yang — *Threshold π^0 Photo- and Electro-Production* — 106

Hadronic Reactions / Sigma Term

R.A. Arndt — *Partial Wave Analysis of Pion-Nucleon Scattering below $T_{\rm LAB}$ = 2.1 GeV* — 107

P. Piirola — *Towards the Pion-Nucleon PWA* — 108

M.F.M. Lutz — *Meson-Baryon Scattering* — 109

M. Frink — *Order Parameters of Chiral Symmetry Breaking* — 110

Chiral Dynamics and Nucleon Resonances

A.M. Bernstein — *Hadron Deformation and Chiral Dynamics* — 111

V. Pascalutsa — *Chiral Dynamics in the $\Delta(1232)$ Region* — 112

Recent Formal Developments

R. Lewis — *Lattice Regularized Chiral Perturbation Theory* — 113

S. Scherer — *Renormalization and Power Counting in Manifestly Lorentz-invariant Baryon Chiral Perturbation Theory* — 114

J. Gegelia — *Baryon ChPT with Vector Mesons* — 115

P.J. Ellis — *Baryon ChPT with Infrared Regularization: Update* — 116

U.-G. Meißner — *Cutoff Schemes in Chiral Perturbation Theory and the Quark Mass Expansion of the Nucleon Mass* — 117

GDH, Nucleon Spin Structure and Chiral Dynamics: Theory

Dieter Drechsel

Institut für Kernphysik, Universität Mainz,
55099 Mainz, Germany

The Gerasimov-Drell-Hearn sum rule ist based on low-energy theorems and forward dispersion relations [1]. It connects the anomalous magnetic moment to an integral over the helicity difference $\sigma_{1/2} - \sigma_{3/2}$ of the photoabsorption cross sections weighted with the inverse power of the photon lab energy ν. This cross section difference was recently measured by the GDH Collaboration [2] at MAMI for 200 MeV $< \nu <$ 800 MeV and at ELSA for 700 MeV $< \nu <$ 2.8 GeV. The experiments confirmed the sum rule within a 10% deviation, and most of the still missing strength is expected at the higher energies on the basis of Regge models. In the mean time the collaboration has also taken data for the deuteron, which holds the promise to solve the "neutron puzzle": Theoretical analysis using photoproduction multipoles results in a 20-30% discrepancy between the sum rule and the dispersion integral for the neutron. At the same time the GDH experiments have provided a value for the forward spin polarizability [1,2] by weighting the cross section difference by ν^{-3}. In the case of virtual photons there appears an additional inclusive cross section σ_{LT} as a longitudinal-transverse interference, and all cross sections and related integrals are also functions of the four-momentum transfer $Q^2 > 0$. These integrals provide various generalizations of the GDH sum rule, the Burkhardt-Cottingham sum rule, and values for generalized polarizabilities. A wealth of new precision data has been obtained at the Jefferson Lab [3], which can be compared to the results of chiral theories [4] and dispersion analysis [1]. These ongoing activities hold the promise to fully determine the spin structure in the resonance region and thus to bridge the gap between low-energy coherent processes and deeply inelastic incoherent scattering.

References

[1] D. Drechsel et al., Physics Reports 378 (2003) 99

[2] B. Schoch, *Polarized total photoabsorption cross section*, these proceedings, A. Braghieri, *Real photon GDH, forward spin polarizability, and chiral dynamics: experiment*, these proceedings

[3] W. Korsch, *Spin structure of the nucleon - experiment*, these proceedings

[4] V. Bernard, *Spin structure of the nucleon - theory*, these proceedings

Helicity Dependence of Pion Photo-Production and the GDH Sum Rule

Alessandro Braghieri

INFN, Sezione di Pavia,
Via Bassi 6, 27100 Pavia, Italy

The helicity dependence of the total photo-absorption cross section on the proton was measured in the energy range $0.2 < E_\gamma < 3.1$ GeV. These data provide the first experimental check of the Gerasimov-Drell-Hearn sum sule [1] and measurement of the forward spin polarizability γ_0.

The experiment was carried out in two steps: at MAMI (Mainz) up to 800 MeV and at ELSA (Bonn) at higher energy. First results from Mainz have already been published [2] and preliminary results from Bonn are available [3]. An estimate of the GDH sum rule was deduced combining these measurements with model predictions in the unmeasured photon energy range and it was found in agreement with the theoretical value. A preliminary estimate of the forward spin polarizability gives $\gamma_0 = (104 \pm 8 \pm 10) \cdot 10^{-6}$ fm^4.

In addition, all partial channels up to E_γ=800 MeV were measured and the helicity dependence of both total and differential cross sections are now available. First results have been published: the single π^+ and π^0 production in the Δ region [4]; the double pion production ($\pi^+\pi^0$) [5]; the η production [6]; the π^0 production in the second resonance region [7].

These measurements provide a strong constraint for multipole analysis [8] of pion photo-production and new information on excitation of baryon resonances.

References

[1] S. B. Gerasimov , Sov. J. Nucl. Phys. 2, 430 (1966);
S. D. Drell and A. C. Hearn, Phys. Rev. Lett. 16, 430 (1966)

[2] J. Ahrens et al., Phys. Rev. Lett. 87, 022003 (2001)

[3] to be published

[4] J. Ahrens et al., Phys. Rev. Lett. 84: 5950-5954, 2000

[5] J. Ahrens et al., Phys. Lett. B551: 49-55, 2003

[6] J. Ahrens et al., Eur .Phys. J. A17: 241-244, 2003

[7] J. Ahrens et al., Phys. Rev. Lett. 88: 232002, 2002

[8] http://gwdac.phys.gwu.edu
http://www.kph.uni-mainz.de/MAID

Virtual Compton Scattering at Low Energy and the Generalized Polarizabilities of the Nucleon

Hélène Fonvieille

Laboratoire de Physique Corpusculaire IN2P3-CNRS
Université Blaise Pascal Clermont-II, 63170 Aubière Cedex, France

For the Jefferson Lab Hall A Collaboration
and the VCS Collaboration

Virtual Compton Scattering has opened a new field of investigation of nucleon structure. At low center-of-mass energies, the process $\gamma^* p \to \gamma p$ allows the determination of the Generalized Polarizabilities (GPs) of the proton [1]. These observables generalize the concept of nucleon polarizabilities to any photon virtuality Q^2. The GPs are predicted by many models at low Q^2, including Heavy Baryon Chiral Perturbation Theory [2]. A first generation of experiments studying photon electroproduction $ep \to ep\gamma$ have been performed at MAMI [3], Jefferson Lab [4] and Bates [5]. They measure the unpolarized VCS structure functions $P_{LL} - P_{TT}/\epsilon$ and P_{LT} which are linear combinations of the lowest order dipole GPs. Analysis methods are based on the Low Energy Theorem [1] or the Dispersion Relation formalism [6]. Results of the MAMI [3] and JLab [7] experiments are presented, together with the future prospects in the field.

References

[1] P. Guichon, G.Q. Liu and A. Thomas, Nucl. Phys. A 591 (1995) 606; P. Guichon and M. Vanderhaeghen, Prog. Part. Nucl. Phys. 41 (1998) 125.

[2] T. Hemmert *et al.*, Phys. Rev. D 62 (2000) 014013; C.W. Kao and M. Vanderhaeghen, Phys. Rev. Lett. 89 (2002) 272002.

[3] J. Roche *et al.*, Phys. Rev. Lett. 85 (2000) 708.

[4] P. Bertin, C. Hyde-Wright, P. Guichon *et al.*, CEBAF Proposal PR93050 (1993). http://hallaweb.jlab.org/experiment/E93-050/vcs.html

[5] J. Shaw, R. Miskimen *et al.*, MIT-Bates Proposal E97-03 (1997).

[6] B. Pasquini *et al.*, Eur. Phys. J. A 11 (2001) 185; D. Drechsel, B. Pasquini and M. Vanderhaeghen, Phys. Rept 378 (2003) 99.

[7] H. Fonvieille, JLab E93-050 preliminary results in Proc. Baryons 2002 Conf. (World Scientific, 2003), hep-ex/0206035; S. Jaminion *et al.*, in preparation; G. Laveissière *et al.*, in preparation.

Dynamical Polarizabilities from (Polarized) Nucleon Compton Scattering

Robert P. Hildebrandt[1], Harald W. Grießhammer[1] and Thomas R. Hemmert[1]

[1]Theoretische Physik T39, Physik-Department, TU München, D-85747 Garching, Germany

Nucleon polarizabilities describe the response of the charged constituents of the nucleon to an external electromagnetic field. In [1] the concept of *dynamical*, i.e. energy-dependent polarizabilities, derived from Compton multipoles, has been introduced. These quantities gauge the response of the internal nucleonic degrees of freedom to an external, real photon field of definite multipolarity and non-zero energy. In this work we present a projector formalism to extract the nucleon polarizabilities from a multipole expansion of the structure part of the six Compton amplitudes. We calculate the dynamical polarizabilities up to leading-one-loop order in an Effective Field Theory with (Small Scale Expansion) and without (HBχPT) explicit $\Delta(1232)$ degrees of freedom and compare our result to dispersion relation calculations [2], finding good agreement in most multipole channels and the possibility to determine the static spin-independent dipole polarizabilities $\bar{\alpha}_{E1}$ and $\bar{\beta}_{M1}$ from low energy Compton data via the SSE-amplitudes. To include the Δ resonance as explicit degree of freedom is crucial for the resonant polarizabilities (*e.g.* $\beta_{M1}(\omega)$) and for a correct description of backward Compton scattering data [2]. We demonstrate that quadrupole polarizabilities give negligibly small contributions to Compton cross sections – spin-averaged as well as polarized ones – i.e. the multipole expansion converges very fast. This leaves us with only the six dipole polarizabilities as free functions of energy which one therefore may be able to fit to experimental Compton scattering data. This hope is confirmed, as we show that the dynamical spin polarizabilities give sizeable contributions even to spin-averaged Compton scattering and that they are absolutely dominant in certain polarization configurations [3]. Finally we present one possible, model-independent way to determine the dynamical spin dipole polarizabilities directly from experiment, proving the principle by fitting only two of them to spin-averaged Compton data.

References

[1] H.W. Grießhammer and T.R. Hemmert, Phys. Rev. C 65, 054207 (2002).

[2] R.P. Hildebrandt, H.W. Grießhammer, T.R. Hemmert and B. Pasquini, nucl-th/0307070, to be published in Eur. Phys. J. A.

[3] R.P. Hildebrandt, H.W. Grießhammer and T.R. Hemmert, nucl-th/0308054.

Radiative Pion Photoproduction and Pion Polarizability

Lev Fil'kov[1]
(for A2 and TAPS collaborations of MAMI)

[1]Lebedev Physical Institute, Leninsky Prospect 53, 119991 Moscow, Russia

The process of the radiative π^+ meson photoproduction from the proton ($\gamma p \to \gamma\pi^+ n$) is studied with the aim to determine the π^+ meson polarizabilities [1,2]. For this purpose an experiment on the process under study has been carried out at MAMI-B in the kinematical region 537 MeV$< E_\gamma <$817 MeV, $140^\circ \leq \theta_{\gamma\gamma}^{cm} \leq 180^\circ$ where $\theta_{\gamma\gamma}^{cm}$ is the photon emission angle in *cms* of the $\gamma\pi$ scattering. The π^+ meson polarizabilities were determined from the comparison of the experimental data with two different theoretical models (without [3] and with baryon resonances). In order to decrease the contribution of the baryon resonances, the differential cross section of the process under study has been integrated over $\theta_{\gamma\gamma}^{cm}$ from 140° to 180° and over φ from 0° to 360° (where φ is the Treiman-Yang angle between the planes formed by the momenta of the initial photon and the proton and the initial photon and the emitted pion in *cms* of the $\gamma\pi \to \gamma\pi$ reaction). To decrease a model dependence of the results obtained, the kinematical region of this process has been chosen where the difference between the predictions of two considered models does not exceed 3% when $(\alpha-\beta)_{\pi^+} = 0$.

The setup efficiency has been normalized by the comparison of the theoretical model predictions with the experimental data in the kinematical region where the pion polarizability contribution is negligible ($s_1 < 5\mu^2$).

In the region, where the pion polarizability contribution is essential ($5 < s_1/\mu^2 < 15,\ -12 < t/\mu^2 < -2$), the difference of the electric and magnetic π^+ meson polarizabilities has been determined from the comparison of the experimental data on the total cross section of the radiative pion photoproduction with the predictions of two theoretical models under consideration. As a very preliminary result we have obtained:

$$(\alpha - \beta)_{\pi^+} = (11.6 \pm 1.5 \pm 5.1) \times 10^{-4}\ \mathrm{fm}^3$$

where the errors are statistical and systematic ones including the difference between the results found with help of the considered models, respectively.

References

[1] D. Drechsel and L.V. Fil'kov, Z. Phys. **A349** (1994) 177.

[2] J. Ahrens *et al.*, Few-Body Syst. Suppl. **9** (1995) 449.

[3] Ch. Unkmeir, PhD Thesis, Mainz University (2000).

Strangeness Production from ELSA

K.-H. Glander

Physikalisches Insitut, Universiät Bonn

Nußallee 12, D-53115 Bonn, Germany

The reactions $\gamma p \to K^+\Lambda$ and $\gamma p \to K+\Sigma^0$ were measured in the energy range from threshold up to a photon energy of 2.6 GeV [1]. The data were taken with the SAPHIR detector at the electron stretcher facility, ELSA. Results on cross sections and hyperon polarizations are presented as a function of kaon production angle and photon energy. The total cross section for Λ production rises steeply with energy close to threshold, whereas the Σ^0 cross section rises slowly to a maximum at about $E_\gamma = 1.45\,\mathrm{GeV}$. Cross sections together with their angular decompositions into Legendre polynomials suggest contributions from resonance production for both reactions. In general, the induced polarization of the Λ has negative values in the kaon forward direction and positive values in the backward direction. The magnitude varies with energy. The polarization of the Σ^0 follows a similar angular and energy dependence as that of the Λ, but with opposite sign. A detailed discussion of the data, their analysis and the results can be found in Ref. [1].

References

[1] K.-H. Glander *et al.*, [SAPHIR collaboration] arXiv:nucl-ex/0308025.

Strangeness Production at Jefferson Lab

Brian Raue

Florida International University,
Miami, FL, USA.

The Thomas Jefferson National Accelerator Facility has an extensive program of studying the electromagnetic production of strange particles. One of the main components of this program has been the study of both photo- and electroproduction of $K^+ + \Lambda^0$ and $K^+ + \Sigma^0$ final states. Experiments are being, or have been conducted in all three of Jefferson Lab's experimental halls measuring a wide range of observables at kinematics from threshold up to $W \approx 3.0$ GeV and Q^2 from 0.4 up to 5 $(\mathrm{GeV}/c)^2$.

The largest effort in this endeavor is taking place in Hall B using the CEBAF Large Acceptance Spectrometer (CLAS). Data have been taken at about ten different polarized electron beam energies and are currently being analyzed.

Preliminary results [1] for one beam energy exist wherein the unpolarized cross section has been separated into three components: $\sigma_T + \epsilon_L \sigma_L$, σ_{TT}, and σ_{LT}. The data indicate a t-channel dominance for the Λ^0 production and a strong s-channel dominance for Σ^0 production. The large σ_{LT} structure function for Λ^0 suggests a large coupling to the longitudinal component of the virtual photon. The W dependence of the structure functions for both final states indicate the likely influence of various intermediate resonances. Precision measurements of the photoproduction cross section have been made using CLAS and may also indicate the presence of previously unobserved intermediate resonances [2].

These results will soon be complemented with a full analysis of all CLAS data wherein σ_T and σ_L will be measured as well as the fifth structure function $\sigma_{LT'}$ [3]. Measurements of the transferred Λ^0 polarization [4] along with high-precision L-T separations from Hall C [5] and Hall A [6] will also provide additional constraints for understanding the underlying strangeness electroproduction process.

References

[1] Robert Feuerbach, Ph.D. Dissertation, Carnegie Mellon University, 2002.

[2] John McNabb, Ph.D. Dissertation, Carnegie Mellon University, 2002.

[3] D. S. Carman *et al.*, JLab experiment E00-112.

[4] D. S. Carman *et al.*, **Phys. Rev. Lett. 90**, 131804 (2003).

[5] R. Mohring *et al.*, **Phys. Rev. C67**, 055205 (2003).

[6] P. Markowitz *et al.*, JLab experiment E98-108.

Unitarized Chiral Dynamics: SU(3) and Resonances

E. Oset, A. Ramos, C. Bennhold, D. Jido, J. Antonio Oller and U.-G. Meißner

Valencia, Barcelona, Washington, Osaka, Murcia and Bonn Collaboration

Using a chiral unitary approach for the meson–baryon interactions, we show that two octets of $J^\pi = 1/2^-$ baryon states, which are degenerate in the limit of exact SU(3) symmetry, and a singlet are generated dynamically. The SU(3) breaking produces the splitting of the two octets, resulting in the case of strangeness $S = -1$ in two poles of the scattering matrix close to the nominal $\Lambda(1405)$ resonance. These poles are combinations of the singlet state and the octets. We show how actual experiments see just one effective resonance shape, but with properties which change from one reaction to another. Suggestions of experiments are made to populate either of the two $\Lambda(1405)$ resonances [1]. Members of these two octets are also the I=0 $\Lambda(1670)$ state and I=1 $\Sigma(1620)$ state with strangeness S=-1 [2], plus the $\Xi(1620)$ state with S=-2 [3]. Some partial evidence is seen for another I=1 state around the KN threshold and hints are given that the differences in the $\pi^+\Sigma^-$ and $\pi^-\Sigma^+$ distributions in $\Lambda(1405)$ photoproduction in a recent experiment at Spring8/Osaka could unveil the existence of this I=1 state [4].

References

[1] D. Jido, J. A. Oller, E. Oset, A. Ramos and U. G. Meißner, Nucl. Phys. A **725** (2003) 181 [arXiv:nucl-th/0303062].

[2] E. Oset, A. Ramos and C. Bennhold, Phys. Lett. B **527** (2002) 99 [Erratum-ibid. B **530** (2002) 260] [arXiv:nucl-th/0109006].

[3] A. Ramos, E. Oset and C. Bennhold, Phys. Rev. Lett. **89** (2002) 252001 [arXiv:nucl-th/0204044].

[4] T. Nakano and J.K. Ahn, Contribution to the PANIC 2002 Conference, Osaka, October 2002, Nucl. Phys. A 721 (2003) 715c.

Threshold Pion Electroproduction: Update on Experiments

Harald Merkel

for the A1 Collaboration

Institut für Kernphysik, Johannes Gutenberg-Universität Mainz
Becherweg 45, 55099 Mainz

While in threshold pion photoproduction experiment and theory is in good agreement, the current published data[1] on threshold pion electroproduction show considerable disagreement with predictions in the framework of Chiral Perturbation Theory[2].

To further investigate this disagreement, an experiment with extended momentum acceptance was performed at MAMI to increase the sensitivity to the small p wave contributions. Simultanously, a measurement with polarized electron beam and out of plane acceptance was performed to investigate the effects of the opening $n\pi^+$ channel on the $p\pi^0$ s wave amplitude via the TL' interference.

The new dataset[3] covers the energy range from threshold up to a center of mass energy of 40 MeV above threshold at a photon virtuality of $Q^2 = 0.05\,\mathrm{GeV}^2/c^2$. The extracted cross sections $\sigma_0 = \sigma_T + \epsilon\sigma_L$, σ_{LT}, and σ_{TT} are below the predictions of ChPT, but in good agreement with the phenomenological models MAID and DMT [4]. The unitary cusp in the the asymmetry $A_{TL'}$ is only described by the DMT model.

References

[1] H. B. van den Brink *et al.*, Phys. Rev. Lett. **74** (1995) 3561;
M. O. Distler *et al.*, Phys. Rev. Lett. **80** (1998) 2294.
H. Merkel *et al.*, Phys. Rev. Lett. **88**, 012301 (2002)

[2] V. Bernard *et al.*, Nucl. Phys. A 607 (1996) 379-401

[3] M. Weis, Doctorate Thesis, Mainz University 2003.

[4] MAID: D. Drechsel et al., Nucl. Phys. A645 (1999) 145-174;
DMT: S.S. Kamalov et al., Phys. Rev. Lett. 83, (1999) 4494, Phys. Rev. C 64 (2001) 032201

Two π^0 Production at Threshold

Martin Kotulla[1] for the TAPS/A2 Collaborations

[1]Department of Physics and Astronomy,
University of Basel, CH-4056 Basel (Switzerland)

The reaction $\gamma p \to \pi^0\pi^0 p$ has been measured using the TAPS BaF_2 calorimeter at the tagged photon facility of the Mainz Microtron accelerator. Close to threshold, chiral perturbation theory (ChPT) predicts that this channel is significantly enhanced compared to double pion final states with charged pions [1]. A ChPT calculation of the $2\pi^0$ channel up to order M_π^2 attributes the total strength dominantly to pion loops at order q^3 [2] - a finding that opens new prospects for the test of ChPT.



Figure 1: Total cross section for the reaction $\gamma p \to \pi^0\pi^0 p$ (full squares) at threshold in comparison with [5] (open circles). The prediction of the ChPT calculation [2] is shown (solid curve) and the prediction of Ref. [4] (dotted curve). The upper limit of the ChPT calculation stems from the uncertainty of the coupling $N^*N\pi\pi$ of the $P_{11}(1440)$ resonance (dashed curve).

Our measurement of the cross section at threshold (see Fig 1) is the first which is sensitive enough for a conclusive comparison with the ChPT calculation [2] and is in agreement with its prediction [3]. The data are also in agreement with a calculation in the unitary chiral approach [4].

References

[1] V. Bernard et al.: Nucl. Phys. A 580 (1994) 475–499.
[2] V. Bernard et al.: Phys. Lett. B 382 (1996) 19–23.
[3] M. Kotulla et al.: Phys. Lett. B accepted.
[4] L. Roca et al.: Phys. Lett. B 541 (2002) 77–86.
[5] M. Wolf et al.: Eur. Phys. J. A 9 (2000) 5-8 .

Threshold π^0 Photo- and Electro-Production

S. N. Yang[1], G.-Y. Chen [1], D. Drechsel[2], S. S. Kamalov[2] and L. Tiator[2]

[1]Dep. of Phys., National Taiwan University, Taipei 10617, Taiwan

[2]Institut für Kernphysik, Universität Mainz, 55099 Mainz, Germany

In this talk we present the predictions of the Dubna-Mainz-Taipei (DMT) dynamical model, based on meson-exchange picture, which we recently developed in Ref. [1] for the threshold electromagnetic pion production and compare them with the results of ChPT[2]. In the DMT model for the pion photo- and electro-production the t-matrix is given as

$$t_{\gamma\pi}(E) = v_{\gamma\pi} + v_{\gamma\pi} g_0(E)\, t_{\pi N}(E) \tag{1}$$

where $v_{\gamma\pi}$ is the $\gamma\pi$ transition potential, g_0 and $t_{\pi N}$ are the πN free propagator and $t-$matrix, respectively, and E is the total energy in the c.m. frame. In the present study, $t_{\pi N}$ is obtained in a meson-exchange πN model [3]. Both $v_{\pi N}$ and $v_{\gamma\pi}$ are derived from an effective Lagrangian containing Born terms as well as ρ- and ω-exchange in the t-channel.

In such approach $t_{\pi N}$ contains the effect of πN rescattering to all orders. However, we have found that only the first order rescattering contribution, i.e. the 1-loop diagram, is important. In Table 1, the results obtained up to tree, 1-loop, and 2-loop approximations for all four possible pion photoproduction channels are listed and compared to the experiments and ChPT results[2]. We see that for π^0 production from both proton and neutron, it is necessary to include one-loop contribution while tree approximation is sufficient for the charged pion productions.

Table 1. Threshold values of E_{0+} (in units $10^{-3}/m_\pi$).

	Tree	1-loop	2-loop	Full	ChPT	Exp
$\pi^0 p$	-2.26	-1.06	-1.01	-1.00	-1.1	-1.33 ± 0.11
$\pi^+ n$	27.72	28.62	28.82	28.85	28.2 ± 0.6	28.3 ± 0.3
$\pi^0 n$	0.46	2.09	2.15	2.18	2.13	
$\pi^- p$	-31.65	-32.98	-33.27	-33.31	-32.7 ± 0.6	-31.8 ± 1.9

References

[1] S. S. Kamalov and S. N. Yang, Phys. Rev. Lett. 83 (1999) 4494; S. S. Kamalov *et al.,* Phys. Rev. C64 (2001) 032201 (R).

[2] V. Bernard, J. Gasser, N. Kaiser, and Ulf-G. Meißner, Phys. Lett. B268 (1991) 291.

[3] C.T. Hung *et al.,* J. Phys. G20 (1994) 1531; Phys. Rev. C64 (2001) 034309.

Partial Wave Analysis of Pion-Nucleon Scattering below $T_{LAB} = 2.1$ GeV

R.A. Arndt[1], W.J. Briscoe[1], M.M. Pavan[2],
I.I. Strakovsky[1] and R.L. Workman[1]

[1]Dept. of Physics, The George Washington University, Washington, D.C., USA

[2]TRIUMF, Vancouver, B.C., Canada

We present results from a partial-wave analysis of pion-nucleon elastic and charge-exchange data, covering the region from threshold to a lab pion kinetic energy of 2.1 GeV, employing a coupled-channel formalism to simultaneously fit $\pi^- p \to \eta n$ data to 0.8 GeV. The main result, fit FA02, has been constrained by a complete set of dispersion-relation constraints, from 20 MeV to 1 GeV, and from t = 0 to -0.4 $(GeV/c)^2$. This represents an improvement, in the number and range of constraints, as compared to our previous SM95 analysis[1].

We have applied a number tests to our solution, including fits with fewer or no dispersion-relation constraints, fits excluding the charge-exchange data base (to gauge the size of charge-symmetry violation), and fits using a different Coulomb correction schemes. A modified form of the Nordita method has been used in the final FA02 result. We have also examined the sensitivity of the resonance extraction to a method commonly used to determine Breit-Wigner parameters. We now fit data directly in determining these BW parameters, as opposed to fitting single-energy solutions, as was done for SM95.

Our results for the pion-nucleon coupling constant and sigma term have not changed significantly with the addition of further dispersion-relation constraints or the inclusion of eta-production data. Somewhat surprisingly, the S_{11} partial-wave is also quite similar to that seen in SM95. However, the revised fitting strategy has resulted in a width for the $S_{11}(1535)$ which is now more in line with the PDG average.

References

[1] R.A. Arndt, I.I. Strakovsky, R.L. Workman, and M.M. Pavan Phys. Rev. C52 (1995) 2120.

Towards the Pion-Nucleon PWA

Pekko Piirola[1] and Mikko E. Sainio[1,2]

[1]Dep. of Physical Sciences, University of Helsinki,
P.O.Box 64, 00014 University of Helsinki, Finland

[2]Helsinki Institute of Physics,
P.O.Box 64, 00014 University of Helsinki, Finland

The last full Karlsruhe partial wave analysis of πN scattering was finished ca. 20 years ago [1,2]. After that a huge pile of new accurate πN scattering measurements has emerged with which the old analysis is not completely compatible. We are making a highly modernized version of the old analysis to all (recent *and* old) data, aiming to pay particular attention to the discrepancies between different data sets and to the handling of the experimental errors as well as to the electromagnetic corrections and the effects of isospin breaking [3].

The most essential feature of our analysis is the use of fixed-t dispersion relations as constraints. Instead of working with dispersion integrals directly, special series for the invariant B- and C-amplitudes are used:

$$C(\nu,t) = C_N(\nu,t) + H(Z,t)\sum_{i=0}^{n} c_i Z^i(\nu,t) \ .$$

Here the base functions $Z^i(\nu,t)$ are conformal mappings with the proper analytic structure [4], and $H(Z,t)$ is a factor describing the expected asymptotic behaviour. When fitting the coefficients c_i, convergence and smoothness of the fit are achieved using the convergence test function method, *i.e.* minimizing not only the pure χ^2, but the sum $\chi^2 + \Phi$, where Φ is the penalty function of the CTF method [4].

At the time of writing (*i.e.* October 2003) we are testing the machinery with the old KA85 solution [2] excluding all of the experimental input, in order to find out the possible internal inconsistencies of our implementation.

References

[1] R. Koch and E. Pietarinen, Nucl. Phys. **A336**, (1980) 331.

[2] R. Koch, Z. Phys. **C29**, (1985) 597.

[3] P. Piirola and M.E. Sainio, Progress in Particle and Nuclear Physics 50 (2003) 367.

[4] E. Pietarinen, Nuovo Cimento **12A**, (1972) 522; E. Pietarinen, Nucl. Phys. **B55**, (1973) 541.

Meson-Baryon Scattering

Matthias F.M. Lutz

Gesellschaft für Schwerionenforschung (GSI)
Planck Str. 1, D-64291 Darmstadt, Germany

We study $J^P = \frac{3}{2}^-$ baryon resonances as generated by chiral coupled-channel dynamics in the $\chi-$BS(3) approach [1,2,3]. Parameter-free results are obtained in terms of the Weinberg-Tomozawa term predicting the leading s-wave interaction strength of Goldstone bosons with baryon-decuplet states [4]. In the 'heavy' SU(3) limit with $m_\pi = m_K \sim 500$ MeV the resonances turn into bound states forming a decuplet and octet representation of the SU(3) group. Using physical masses the mass splitting are remarkably close to the empirical pattern.

Our effective field theory is based on the assumption that at subthreshold energies the scattering amplitudes can be evaluated in standard chiral perturbation theory. Once the available energy is sufficiently high to permit elastic two-body scattering a further typical dimensionless parameter $m_K^2/(8\,\pi f^2) \sim 1$ arises if strangeness is considered explicitly. Since this ratio is uniquely linked to two-particle reducible diagrams it is sufficient to sum those diagrams keeping the perturbative expansion of all irreducible diagrams. We suggest to glue s- and u-channel unitarized scattering amplitudes together at subthreshold energies [3,4]. This construction reflects our basic assumption that diagrams showing a s- or u-channel unitarity cut need to be summed to all orders at least at energies close to where the diagrams develop their imaginary part. By construction, a glued scattering amplitude satisfies crossing symmetry exactly at energies where the scattering process takes place. At subthreshold energies crossing symmetry is implemented approximatively only, however, to higher and higher accuracy when more chiral correction terms are considered. Insisting that the scattering amplitude reproduces the interaction kernel at subthreshold energies guarantees that subthreshold amplitudes match smoothly and therefore the final glued amplitudes comply with the crossing-symmetry constraint to high accuracy.

References

[1] M.F.M. Lutz and E.E. Kolomeitsev, Proc. of Int. Workshop XXVIII on Gross Properties of Nuclei and Nuclear Excitations, Hirschegg, Austria, January 16-22, 2000.

[2] M.F.M. Lutz und E.E. Kolomeitsev, Found. Phys. 31 (2001) 1671.

[3] M.F.M. Lutz and E.E. Kolomeitsev, Nucl. Phys. A 700 (2002) 193.

[4] E.E. Kolomeitsev and M.F.M. Lutz, nucl-th/0305101.

Order Parameters of Chiral Symmetry Breaking from Meson – Baryon Dynamics

Matthias Frink

IKP, Forschungszentrum Jülich
D-52425 Jülich, Germany

In a theory with N_f light quark flavours spontaneous breaking of chiral symmetry $SU(N_f)_V \times SU(N_f)_A \to SU(N_f)_V$ entails $N_f^2 - 1$ (approximately) massless Goldstone bosons. Order parameters, i. e. vacuum expectation values of operators transforming non–trivially under the broken symmetry group, are a useful tool to examine symmetry breaking. Examples of chiral order parameters are the quark condensate and Goldstone boson decay constants.
Chiral order parameters O are conjectured to behave paramagnetically with an increasing number of light quark species, i. e. $O(N_{f+1}) < O(N_f)$ [1]. Fields of relevance and possibel applications are e. g. the determination of hadron masses, the applicability of Generalized ChPT and an approach to OZI rule violation in the scalar sector [2].
Descotes and Stern have analyzed the role of chiral order parameters in the mesonic sector [1], [2]. A step towards a further investigation is the inclusion of meson–baryon dynamics into the analysis.
Basic input for such calculations is the complete and minimal three–flavour meson–baryon Lagrangian. Its third–order portion has been formulated by Krause [3]. Building blocks for the Lagrangian to fourth order are besides baryon and meson fields external source terms and Clifford–algebra elements. Various reduction mechanisms can be used in the process of minimization, e. g. relations between the fields, trace relations for $SU(3)$, identities among Clifford–algebra elements, and equation of motion eliminations. Work is in progress here.
As a first application we plan to calculate octet baryon masses to fourth order in the chiral expansion, using the infrared regularization scheme. The corresponding third–order contributions are given in [4]. Further applications comprise e. g. scalar baryon form factors and their low–energy relation with scattering amplitudes as well as sigma–term and strangeness content determinations.

References

[1] S. Descotes, L. Girlanda, and J. Stern, hep-ph/9910537.

[2] S. Descotes, hep-ph/0012221.

[3] A. Krause, Helv. Phys. Acta 63 (1990) 3.

[4] P. J. Ellis and K. Torikoshi, nucl-th/9904017.

Hadron Deformation and Chiral Dynamics

A.M.Bernstein
Physics Dept. and Laboratory for Nuclear Science,M.I.T.
Cambridge, Mass., U.S.A.

The much conjectured deviation of the proton shape from spherical symmetry cannot be directly observed since it has spin 1/2 and therefore no quadrupole moment. This has led to an active field of measurements of the electromagnetic transition of the proton to its first excited state, the Δ with spin 3/2. We have taken accurate data at $Q^2 = 0.126 GeV^2$ at the MIT/Bates accelerator with the out of plane spectrometer system (OOPS)[1]. These and other photo- and elctropion production experiments[1,2] have observed significant non-zero electric and Coulomb quadrupole matrix elements which signify deviations from spherical symmetry in the proton and/or the Δ. These observed quadrupole amplitudes are generally an order of magnitude greater then predicted by the quark model[2]. As has been pointed out for years in terms of the cloudy bag model, this should not be surprising since quark models do not properly handle chiral symmetry or the pion. More recently two dynamical models of the $\gamma^* p \rightarrow \pi N$ reaction[3] have shown that the contribution of the pion cloud is the dominant one and that this maximizes in the vicinity of our Bates data[1] .

In my view the dominance of pionic effects follows naturally from the spontaneous breaking of chiral symmetry. The coupling of a pion (Goldstone Boson) to a nucleon is given $L_{\pi N} = g\vec{\sigma} \cdot \vec{k}$ where g is the $\pi - N$ coupling constant(predicted by chiral symmetry), σ is the nucleon spin, and k is the pion momentum. This interaction vanishes in the s wave and is strong in the p wave. This is the basis of the deviation from spherical symmetry in the nucleon since the pion cloud is emitted and absorbed in the p wave. Furthermore this is the same physical effect as is observed in the well known, long range, tensor force between nucleons.$V_{NN} = g^2 \ \vec{\sigma_1} \cdot \vec{k} \ \frac{1}{k^2 + m_\pi^2} \ \vec{\sigma_2} \cdot \vec{k}$.

It is an open problem to calculate the deformation in ways more strongly linked to QCD such as ChPT or the lattice, and to perform more measurements to determine the precise magnitude of the proton and Δ deformations.

References

[1] C.N.Papanicolas, International Conference on Quark Nuclear Physics, Julich, Germany, 2002, to be published in Europ. Jour. A C. Kunz et. al.Phys. Lett. B564,21(2003). N.F.Sparveris et al., to be published(2003).

[2] A.M. Bernstein, Eur.Phys.J.A17:349-355(2003).
Enhanced version hep-ex/0212032, and references therein.

[3] T. Sato and T.-S. H. Lee, Phys. Rev. Phys.Rev.**C54**, 2660(1996),
S.S. Kamalov and S.N. Yang, Phys. Rev. Lett. **83**, 4494 (1999).

Chiral Dynamics in the $\Delta(1232)$ Region

Vladimir Pascalutsa[1,2] and Daniel R. Phillips[2]

[1]Department of Physics, College of William & Mary, Williamsburg, VA 23185
and Theory Group, JLab, 12000 Jefferson Ave, Newport News, VA 23606

[2]Department of Physics and Astronomy, Ohio University, Athens, OH 45701

The limit of applicability of ChPT in the baryon sector is set by the excitation energy of the first resonance: $\Delta \equiv M_\Delta - M_N \approx 293$ MeV. We have recently proposed a systematic scheme that extends ChPT into the resonance region in a natural way [1] (see also [2] for a more concise explanation).

Our power counting catches the most crucial feature of the resonance physics: the need for the resummation of the self-energy insertions which result in dynamical generation of the width. This is achieved by distinguishing the resonance-excitation energy from the other low-energy hadronic scales, such as m_π. In terms of small expansion parameter δ, we assume that $\Delta = O(\delta)$ while the lower-energy scales are of higher order, e.g., $m_\pi = O(\delta^2)$. The power-counting then depends on kinematics, namely on whether we are in the low-energy regime $p \sim m_\pi$ or in the resonance region $p \sim \Delta$, where p is a typical relative momentum. It shows that at low energies the Δ-isobar contributions are suppressed with respect to the nucleon ones by a power of the small parameter. One the other hand, in the resonance region one class of graphs — the one-Δ-reducible graphs — needs to be resummed, since each graph in this class goes as some power of $1/(p-\Delta)$ and hence does not have a well-defined power counting for $p \sim \Delta$.

After the resummation, the one-Δ-reducible graphs obtain a definite power-counting index, which is determined by the index of the self-energy. The expansion of the Δ self-energy begins with the one πN-loop, which goes of course as p^3. Since the Δ-propagator in a *resummed* one-Δ-reducible graph is $1/(p-\Delta-\Sigma)$, it counts as $1/\delta^3$ in the resonance regime. Thus, the one-Δ-reducible graphs are enhanced in the resonance region by two powers of the small parameter.

The mechanisms of Δ excitation and its dynamical width generation are of course present in many phenomenological models. Our goal was to incorporate them into a systematic power-counting scheme which could extend ChPT.

In this talk we have also discussed several applications. Most notably, the extraction of the magnetic moment of the $\Delta(1232)$, where both $p \sim m_\pi$ and $p \sim \Delta$ regimes are at play simultaneously!

References

[1] V. Pascalutsa and D. R. Phillips, Phys. Rev. C **67**, 055202 (2003).

[2] V. Pascalutsa and D. R. Phillips, arXiv:nucl-th/0308065.

Lattice Regularized Chiral Perturbation Theory

Buğra Borasoy[1], **Randy Lewis**[2] and Pierre-Philippe A. Ouimet[2]

[1]Physik Department, Technische Universität München, D-85747 Garching, Germany

[2]Department of Physics, University of Regina, Regina, SK, S4S 0A2, Canada

The spacetime lattice is a regularization technique for quantum field theories. Its application to chiral perturbation theory (ChPT) can be of interest to lattice QCD studies since the effective theory, lattice ChPT, is then defined in the same discrete spacetime as the underlying theory, lattice QCD.[1] Lattice ChPT can also be viewed as yet another way to implement a cut-off regularization scheme, somewhat comparable to Ref. [2], and with the advantage that chiral symmetry and gauge invariance are automatically perserved throughout any lattice ChPT calculation. Since lattice regularization does not rely on perturbation theory, it might be valuable for multi-nucleon effective field theories as well.[3]

The simplest lattice ChPT Lagrangian is obtained by writing the standard ChPT Lagrangian in Euclidean spacetime and converting derivatives to finite differences in such a way as to preserve the desired symmetries without introducing doublers or ghost states. In the continuum limit, ChPT observables are independent of regularization scheme and continuum observables agree exactly with dimensional regularization. For more details, see Ref. [4].

References

[1] S. Myint and C. Rebbi, Nucl. Phys. B421, 241 (1994); A. R. Levi, V. Lubicz and C. Rebbi, Phys. Rev. D56, 1101 (1997); I. A. Shushpanov and A. V. Smilga, Phys. Rev. D59, 054013 (1999); R. Lewis and P.-P. A. Ouimet, Phys. Rev. D64, 034005 (2001); B. Borasoy, R. Lewis and P.-P. A. Ouimet, Phys. Rev. D65, 114023 (2002).

[2] J. F. Donoghue and B. R. Holstein, Phys. Lett. B436, 331 (1998); J. F. Donoghue, B. R. Holstein and B. Borasoy, Phys. Rev. D59, 036002 (1999); B. Borasoy, B. R. Holstein, R. Lewis and P.-P. A. Ouimet, Phys. Rev. D66, 094020 (2002). R. D. Young, D. B. Leinweber and A. W. Thomas, Prog. Nucl. Part. Phys. 50, 399 (2003); V. Bernard, T. R. Hemmert and U.-G. Meißner, hep-ph/0307115.

[3] S. Chandrasekharan, M. Pepe, F. D. Steffen and U. J. Wiese, hep-lat/0306020.

[4] B. Borasoy, R. Lewis and P.-P. A. Ouimet, hep-lat/0310054.

Renormalization and Power Counting in Manifestly Lorentz-invariant Baryon Chiral Perturbation Theory

Thomas Fuchs[1], Jambul Gegelia[1,2], George Japaridze[3] and **Stefan Scherer**[1]

[1]Institut für Kernphysik, Johannes Gutenberg-Universität, Mainz, Germany

[2] High Energy Physics Institute, Tbilisi State University, 380086 Tbilisi, Georgia

[3] CTSPS, Clark Atlanta University, Atlanta, Georgia 30314, USA

The implementation of a successful effective field theory program requires two main ingredients, namely: (a) a knowledge of the most general effective Lagrangian and (b) an expansion scheme for observables in terms of a consistent power counting method. It was shown by Gasser, Sainio, and Švarc [1] that, in the baryonic sector, the application of dimensional regularization in combination with the modified minimal subtraction scheme of chiral perturbation theory (ChPT) does not suffice to produce a simple correspondence between the loop expansion and the chiral expansion. A solution to this problem was first obtained in the framework of the heavy-baryon formulation of ChPT [2]. More recently, the problem was also attacked in a manifestly Lorentz-invariant form [3,4]. In Ref. [4] we have proposed a new renormalization scheme which makes use of finite subtractions of dimensionally regularized diagrams beyond the standard $\overline{\rm MS}$ scheme of ChPT to remove contributions violating the power counting. Such subtractions can be realized in terms of local counterterms in the most general effective Lagrangian. Our approach may be used in an iterative procedure to renormalize higher-order loop diagrams and also allows for implementing a consistent power counting when vector (and axial-vector) mesons are explicitly included [5].

References

[1] J. Gasser, M. E. Sainio, and A. Švarc, Nucl. Phys. **B307** (1988) 779.

[2] E. Jenkins and A. V. Manohar, Phys. Lett. B **255** (1991) 558; **259** (1991) 353; V. Bernard, N. Kaiser, J. Kambor, and U.-G. Meißner, Nucl. Phys. **B388** (1992) 315.

[3] H. Tang, hep-ph/9607436; P. J. Ellis and H. B. Tang, Phys. Rev. C **57** (1998) 3356; T. Becher and H. Leutwyler, Eur. Phys. J. C **9** (1999) 643; M. F. Lutz and E. E. Kolomeitsev, Nucl. Phys. **A700** (2002) 193.

[4] T. Fuchs, J. Gegelia, G. Japaridze, and S. Scherer, Phys. Rev. D **68** (2003) 056005; T. Fuchs, J. Gegelia, and S. Scherer, hep-ph/0309234.

[5] T. Fuchs, M. R. Schindler, J. Gegelia, and S. Scherer, hep-ph/0308006, to appear in Phys. Lett. B.

Baryon ChPT with Vector Mesons

T. Fuchs[1], M. R. Schindler[1], **J. Gegelia**[1,2] and S. Scherer[1]

[1]Institut für Kernphysik, Johannes Gutenberg-Universität, Mainz, Germany

[2]High Energy Physics Institute, Tbilisi State University, 380086 Tbilisi, Georgia

By considering characteristic self-energy and vertex Feynman diagrams [1] we have demonstrated that the inclusion of explicit virtual (axial) vector particles in manifestly Lorentz-invariant baryon chiral perturbation theory does not violate the power counting if one uses the extended on-mass-shell (EOMS) renormalization scheme of Ref. [2]. This renormalization procedure can also be (iteratively) applied to multiloop diagrams leading to a consistent power counting.

We have reformulated the IR regularization of Becher und Leutwyler [3] in a form analogous to the EOMS renormalization scheme [4]. Within this new formulation the subtraction terms are found by expanding the integrands of loop integrals in powers of small parameters (small masses and Lorentz-invariant combinations of external momenta and large masses) with subsequent exchange of integration and summation. Within this new formulation of IR regularization one does not necessarily need to use dimensional regularization. The renormalized results are regularization scheme independent.

The IR regularization in its new formulation can be easily applied to diagrams with an arbitrary number of propagators with various masses (e.g. resonances) and/or diagrams with several fermion lines as well as to multiloop diagrams.

Applying IR regularization in our formulation to EFT with explicit vector mesons in the antisymmetric tensor field representation and analyzing the diagrams contributing to the electromagnetic form factors of the nucleon to $\mathcal{O}(q^4)$, we observe that in Ref. [5] all relevant loop integrals have actually been taken into account. This is due to the fact that those loop integrals involving vector mesons which formally contribute at a given order vanish in IR regularization.

References

[1] T. Fuchs, M. R. Schindler, J. Gegelia and S. Scherer, arXiv:hep-ph/0308006, to appear in Phys. Lett. B.

[2] T. Fuchs, J. Gegelia, G. Japaridze and S. Scherer, Phys. Rev. D **68** (2003) 056005.

[3] T. Becher and H. Leutwyler, Eur. Phys. J. C **9** (1999) 643.

[4] M. R. Schindler, J. Gegelia and S. Scherer, arXiv:hep-ph/0309005.

[5] B. Kubis and U. G. Meißner, Nucl. Phys. A **679** (2001) 698.

Baryon ChPT with Infrared Regularization: Update

Paul J. Ellis

School of Physics & Astronomy, University of Minnesota
Minneapolis, MN 55455, USA

In order to preserve power counting, loop integrals which involve both baryon and meson propagators are often evaluated using the heavy baryon (HB) approximation. Infrared regularization (IR) is an alternative method proposed by Becher and Leutwyler [1] which retains only the infrared singular part of the integral; the remainder can be expanded in a Taylor series and absorbed in the low energy constants. The IR part includes the corresponding HB result, thus preserving power counting, and also sums an infinite series of (baryon mass)$^{-n}$ corrections which appear in higher orders in the HB scheme. This gives the correct analytic behavior in threshold regions which is not always true in the HB case.

HB results for the baryon masses in SU(3) give third order contributions of similar magnitude to second order, whereas using IR they are smaller by roughly a factor of a half so that the convergence is much improved [2]. The convergence of the chiral expansion for electromagnetic form factors through fourth order [3] is also improved when IR is employed. Including vector mesons explicitly, IR yields a good account of the data. In $\pi - N$ scattering the IR representation does not allow a sufficiently accurate extrapolation [4] from the threshold region to the unphysical Cheng-Dashen point. Further in the physical region [5] IR allows a fit to the phase shifts only over a much more limited region than the HB scheme and even then the parameters are unreasonable. An IR study of the baryon octet axial vector currents shows poor convergence even when the decuplet is included [6]. (For discussion of the spin structure of the nucleon see V. Bernard, Plenary Session 2). Overall the picture is mixed at the moment.

References

[1] T. Becher and H. Leutwyler, Eur. Phys. J. C **9** (1999) 643.

[2] P.J. Ellis and K. Torikoshi, Phys. Rev. C **61** (2000) 015205; see also M. Frink, Working Group 2.

[3] B. Kubis and Ulf-G. Meißner, Nucl. Phys. **A679** (2001) 698; Eur. Phys. J. C **18** (2001) 747.

[4] T. Becher and H. Leutwyler, J. High Energy Phys. **6** (2001) 017.

[5] K. Torikoshi and P.J. Ellis, Phys. Rev. C **67** (2003) 015208.

[6] S-L. Zhu, G. Sacco and M.J. Ramsey-Musolf, Phys. Rev. D **66** (2002) 034021.

Cutoff Schemes in Chiral Perturbation Theory and the Quark Mass Expansion of the Nucleon Mass

Véronique Bernard[1], Thomas R. Hemmert[2] and **Ulf-G. Meißner**[3]

[1]Laboratoire de Physique Théorique, ULP, F-67084 Strasbourg, France
[2]Physik Department T39, TU München, D-85747 München, Germany
[3]HISKP (TH), Universität Bonn, D-53115 Bonn, Germany

We discuss the use of cutoff methods in chiral perturbation theory [1]. We develop a cutoff scheme based on the operator structure of the effective field theory that allows to suppress high momentum contributions in Goldstone boson loop integrals and by construction is free of the problems traditional cutoff schemes have with gauge invariance or chiral symmetries. As an example, we discuss the chiral expansion of the nucleon mass. Contrary to other claims in the literature we show that the mass of a nucleon in heavy baryon chiral perturbation theory has a well behaved chiral expansion up to effective Goldstone boson masses of 400 MeV when one utilizes standard dimensional regularization techniques. With the help of the here developed cutoff scheme we can demonstrate a well-behaved chiral expansion for the nucleon mass up to 600 MeV of effective Goldstone Boson masses. We also discuss in detail the prize, in numbers of additional short distance operators involved, that has to be paid for this extended range of applicability of chiral perturbation theory with cutoff regularization, which is usually not paid attention to. We also compare the fourth order result for the chiral expansion of the nucleon mass with lattice results and draw some conclusions about chiral extrapolations based on such type of representation. In particular, the recent CP-PACS data [2] can be well represented by the fourth order dimensionally regularized nucleon mass expression with LECs consistent with values obtained from the analysis of pion-nucleon scattering, see e.g. [3,4].

References

[1] V. Bernard, T. R. Hemmert and U.-G. Meißner, arXiv:hep-ph/0307115.

[2] A. Ali Khan *et al.* [CP-PACS Collaboration], Phys. Rev. D **65**, 054505 (2002) [Erratum-ibid. D **67**, 059901 (2003)] [arXiv:hep-lat/0105015].

[3] P. Büttiker and U.-G. Meißner, Nucl. Phys. A **668**, 97 (2000) [arXiv:hep-ph/9908247].

[4] N. Fettes and U.-G. Meißner, Nucl. Phys. A **676**, 311 (2000) [arXiv:hep-ph/0002162].

Working Group III: Few-Body Dynamics and Hadronic Atoms

Convenors: Paulo Bedaque, Nasser Kalantar-Nayestanaki and Akaki Rusetsky

Few-Body Dynamics I

U. van Kolck	*Charge Symmetry Breaking in Pion Production*	119
E. Stephenson	*Observation of the Charge Symmetry Breaking $d+d \to {}^4He+\pi^0$ Reaction near Threshold*	120
A. Nogga	*Probing Chiral Interactions in Light Nuclei*	121
F. Myhrer	*Solar Neutrino Reactions and Effective Field Theory*	122
S. Ando	*Solar-Neutrino Reactions on Deuteron in EFT* and Radiative Corrections of Neutron Beta Decay*	123
M.C. Birse	*EFT's for the Inverse-Square Potential and Three-Body Problem*	124

Hadronic Atoms

M. Cargnelli	*DEAR - Kaonic Hydrogen: First Results*	125
J. Gasser	*Comments on $\bar{K}N$ Scattering*	126
A.N. Ivanov	*On Pionic and Kaonic Hydrogen*	127
P. Zemp	*Deser-type Formula for Pionic Hydrogen*	128
L. Nemenov	*Atoms Consisting of $\pi^+\pi^-$ and πK Mesons*	129
L. Girlanda	*Deeply Bound Pionic Atoms: Optical Potential at $O(p^5)$ in ChPT*	130

Few-Body Dynamics II

H. Grießhammer	*All-Order Low-Energy Expansion in the 3-Body System*	131
T. Black	*New Results in the Three and Four Nucleon Systems*	132
H. Schieck	*Nd Scattering at Low Energies*	133
W. Tornow	*Nucleon-Deuteron Scattering at Low Energies*	134
S. Kistryn	*Three-Nucleon System Dynamics Studied in the dp-Breakup at 130 MeV*	135
J. Messchendorp	*Few-Body Studies at KVI*	136
H. Sakai	*dp Elastic Scattering Experiments*	137

Few-Body Dynamics III

J.F. Donoghue	*Nuclear Binding Energies and Quark Masses*	138
F. Rathmann	*Proton-Proton Scattering Experiments*	139
C. Hanhart	*Systematic Approaches to Meson Production in NN and dd Collisions*	140
D. Cabrera	*Vector Meson Properties in Nuclear Matter*	141
A. Schwenk	*A Renormalization Group Method for Ground State Properties of Finite Nuclei*	142
L. Platter	*The Four-Body System in EFT*	143
H. Krebs	*Electroproduction of Neutral Pions Off the Deuteron Near Threshold*	144

Charge Symmetry Breaking in Pion Production [1]

U. van Kolck[1,2]

[1]Department of Physics, University of Arizona,
Tucson, AZ 85721, USA

[2]RIKEN BNL Research Center, Brookhaven National Laboratory,
Upton, NY 11973, USA

Charge symmetry is a rotation of π around the y axis in isospin space. One of the most important low-energy consequences of charge-symmetry breaking is the proton-neutron mass splitting $\Delta m_N = m_p - m_n = -1.3$ MeV, which receives contributions from both the quark-mass difference —denoted δm_N— and electromagnetism —denoted $\bar{\delta} m_N$. At least one other suitable CSB observable is needed to separate these two low-energy constants.

The approximate chiral symmetry of QCD implies [1] that there are two sets of pion-nucleon interactions with strengths fixed by δm_N and $\bar{\delta} m_N$, respectively. Assuming $\bar{\delta} m_N > 0$ as in most models, it was predicted in Ref. [2] that the corresponding contributions to the front-back asymmetry in the reaction $pn \to d\pi^0$ near threshold would be 1–2 times larger, but in the opposite direction, than the sum of other, more conventional mechanisms. This prediction has recently been confirmed at TRIUMF [3].

In addition, an unambiguous signal was seen at IUCF for the CSB reaction $dd \to \alpha\pi^0$ near threshold, resulting in a cross-section of about 10 pb [4]. A group of theorists is now working on the evaluation of the effects from these (and other) interactions [5].

The corresponding two-pion exchange two-nucleon potential has also been calculated [6], but its effects on CSB observables are generically small.

References

[1] U. van Kolck, Texas Thesis (1993); Few-Body Syst. Suppl. 9 (1995) 444.

[2] U. van Kolck, J.A. Niskanen, and G.A. Miller, Phys. Lett. B 493 (2000) 65.

[3] A.K. Opper et al, `nucl-ex/0306027`.

[4] E.J. Stephenson et al, Phys. Rev. Lett. 91 (2003) 142302.

[5] A. Gårdestig et al, in preparation.

[6] J.L. Friar, U. van Kolck, G.L. Payne, and S.A. Coon, Phys. Rev. C 68 (2003) 024003.

[1]Research supported in part by the US DOE and the Alfred P. Sloan Foundation.

Observation of the Charge Symmetry Breaking $d + d \rightarrow {}^4He + \pi^0$ Reaction near Threshold

E.J. Stephenson, for the CSB Collaboration

Indiana University Cyclotron Facility,
Bloomington, IN 47408 USA.

We report the first observation of the isospin violating and charge symmetry breaking (CSB) $d + d \rightarrow {}^4He + \pi^0$ reaction near threshold. The measurements were made with the Indiana University electron-cooled storage ring. A magnetic channel was constructed just downstream of a 6° bending magnet to separate the forward-going ^{4}He recoil nuclei from the circulating beam. A deuterium gas jet target was placed upstream of the bend and surrounded by two arrays of Pb-glass detectors to signal the presence of photons from the decay of the π^0. The formation of a CSB π^0 was separated from the continuum of isospin-allowed double radiative capture ${}^4He + \gamma + \gamma$ events by measuring the ^{4}He recoil momentum and reconstructing the pion missing mass. The 4-momentum measurement depended upon the ^{4}He time of flight in the channel and scattering angle. A clean sample of candidate events was produced by requiring the correct energy loss in the channel scintillators and the presence of two coincident 70-MeV photons in the Pb-glass arrays. The CSB π^0s showed a clear peak above the double radiative capture continuum.

Measurements were made at two energies to verify the algorithms for pion reconstruction. The total CSB cross section was found to be 12.7 ± 2.2 pb at 228.5 MeV and 15.1 ± 3.1 pb at 231.8 MeV [1]. The systematic errors of 6.6% were considerably less than the statistical errors from background subtraction. These measurements are consistent with S-wave pion production and are proportional to $\eta = p_\pi / m_\pi$ with a slope of $\sigma_{\rm TOT}/\eta = 80 \pm 11$ pb. The integral of the double radiative capture process for the 2 MeV range just below the kinematic limit in missing mass was 6.9 ± 0.9 pb at 228.5 MeV and 9.5 ± 1.4 pb at 231.8 MeV.

Charge symmetry breaking effects arise fundamentally from the differences in the masses of the down and up quarks and differences in their electromagnetic interactions. These two effects give rise to the neutron-proton mass difference. A detailed analysis of the $d + d \rightarrow {}^4He + \pi^0$ reaction is underway [2]. It will include entrance channel distortions, modern wavefunctions, and the contributions to CSB from meson mixing ($\pi^0 - \eta$ and $\rho^0 - \omega$).

References

[1] E.J. Stephenson *et al.*, Phys. Rev. Lett. **91**, 142302 (2003).

[2] A. Gårdestig *et al.*, to be published.

Probing Chiral Interactions in Light Nuclei

Andreas Nogga

Institute for Nuclear Theory, University of Washington, Box 351550, Seattle, WA 98195-1550, USA

Recently, chiral perturbation theory was used to obtain a consistent model of NN and 3N interactions (see [1]). The model is able to describe nd scattering up to at least 65 MeV reasonably well. The chiral 3N force has two *a priori* unknown constants, which were determined using the 3N binding energy and the doublet nd scattering length. The resulting 3N force could also describe the ^{4}He binding energy.

In Ref. [2] it was found that the binding energy of the p-shell nuclei are sensitive to the 3NF structure. A simple 3NF model, which is able to describe the 3N and 4N systems, at least at low energies, does not provide the correct binding and excitation energies for p-shell nuclei. This can be resolved by additional 3NF terms. The aim of this study is to understand the effects of chiral NN and 3N interactions on these spectra. The Schrödinger equation for p-shell nuclei is solved using the no-core shell model approach [3]. Here we concentrate on ^{6}Li.

We need to establish the convergence with respect to the chiral expansion. The calculation shows that the p-shell binding energies are affected by NNLO contributions, which is the highest order in the chiral expansion, we take into account. We also find variations of the order of 1.5 MeV varying the cut-off of our interactions. This indicates that our binding energies can not be predicted to a higher accuracy at this order. The excitation energy is also affected by NNLO contributions. The cut-off dependence indicates that the error of the NNLO prediction is 200 keV. The experimental value is within the predicted range. We have also established that the 3NF terms can change the excitation energy and binding energy of ^{6}Li, even if the ^{3}H and ^{4}He binding energies remain constant. The error estimates hint to visible N3LO contributions, which will be subject of a forthcoming study. More details of this work can be found in [4].

The work was supported in parts by grants from the NSF and DOE. The numerical calculations have been made at NERSC.

References

[1] E. Epelbaum *et al.*, Phys. Rev. C **66**, 064001 (2002).

[2] S. C. Pieper *et al.*, Phys. Rev. C **64**, 014001 (2001).

[3] P. Navrátil *et al.*, Phys. Rev. C **62**, 054311 (2000).

[4] A. Nogga *et al.*, to be published in Nucl. Phys. A.

Solar Neutrino Reactions and Effective Field Theory

Fred Myhrer

Univ. South Carolina, Columbia, SC 29208, USA .

Many astrophysical phenomena are governed by low-energy nuclear weak-interaction processes, and effective field theory (EFT) is believed to be a useful framework for describing them. I present here a comparison of EFT with the standard nuclear physics approach (SNPA), focusing on the νd reactionsrelevant for the SNO experiments. The low-energy νd reactions are dominated by the spatial component of the axial current, $\vec{A} = \vec{A}_{one} + \vec{A}_{two}$, a sum of 1-body and 2-body terms. $\vec{A}_{one}$ being well known, one's task is to pin down $\vec{A}_{two}$. In SNPA, $\vec{A}_{two}$ is derived from pion, rho and A_1 meson exchange diagrams [1], devised to fulfill the low-energy theorems, current algebra, etc. SNPA contains some unsatisfactory aspects: (1) There is no obvious link between SNPA and QCD; (2) It does not offer a systematic way to estimate theoretical uncertainties; (3) Low-energy physics ($E \ll m_\pi$) and high-energy physics ($E \geq m_\pi$) are intermingled. EFT can resolve these problems.

The terms in the EFT lagrangian, $\mathcal{L}_\chi$, form a perturbation series in powers of (Q/Λ_χ), where Q is a typical energy of the process in question. In [1] we used EFT* wherein the operators are derived from $\mathcal{L}_\chi$ while the nuclear wave functions are generated from a realistic NN potential. The $\vec{A}_{two}$ in [1] involves a pion-exchange and a four-nucleon contact term with one LEC. Since this LEC is determined from the triton β-decay rate, EFT* allows a model-independent calculation of the νd cross sections[1]. The results agree with those obtained in SNPA to 1 % . Butler et al.'s EFT calculation[2] has one unknown LEC, L_{1A}, which was adjusted to reproduce the ν-d cross sections calculated in [1]. The attempts to constrain L_{1A} from experimental values of observables are not yet very stringent.

References

[1] S. Nakamura et al., Nucl. Phys. A 707 (2002) 561; S. Ando et al., Phys. Lett. B, 555 (2003) 49; S. Ando, contribution to the present proceedings.

[2] M. Butler et al., Nucl. Phys. A 675 (2000) 575; Phys. Rev. C 63 (2001) 035501.

Solar-Neutrino Reactions on Deuteron in EFT* and Radiative Corrections of Neutron Beta Decay

Shung-ichi Ando

TRIUMF, 4004 Wesbrook Mall, Vancouver, B.C. V6T 2A3, Canada

The cross sections for low-energy neutrino-deuteron reactions are calculated within heavy-baryon chiral perturbation theory employing a cut-off regularization scheme[1]. The transition operators are derived up to next-to-next-to-next-to-leading order in accord with the Weinberg counting rule, while the nuclear matrix elements are evaluated using wave functions generated by a high-quality phenomenological NN potential. Only one unknown low-energy constant appears in our calculation, the axial-current-four-nucleon coupling constant, which is fixed by using data from tritium beta decay[2]. Our results exhibit a high degree of stability against different choices of the cutoff parameter, a feature which indicates that, apart from radiative corrections, the uncertainties in the calculated cross sections are less than 1%. We also discuss the feasibility of fixing the low energy constant through the accurate measurement of the ordinary muon capture rate on the deuteron being planned at PSI, which would avoid the complication of the three-body current in tritium beta decay[3].

The lifetime and angular correlation coefficients of neutron beta decay are also evaluated up to next-to-leading order in effective field theory[4], where pions are integrated out because of the small typical energy. Up to this order the nucleon recoil corrections, including weak-magnetism, and the radiative corrections are calculated. Our results agree with those of the long-range and model-independent part of previous calculations. Moreover, in the effective theory the model dependent "inner" radiative corrections are replaced by a well-defined low energy constant. Based on the counting rule of the effective theory, we estimate the accuracy of our results to be of the order of 10^{-3}.

References

[1] S. Ando, Y.H. Song, T.-S. Park, H. W. Fearing, K. Kubodera, Phys. Lett B555(2003)49: S. Nakamura *et al.*, Nucl. Phys. A707(2002)561.

[2] T.-S. Park *et al.*, Phys. Rev. C67(2003)055206.

[3] S. Ando, T.-S. Park, K. Kubodera, F. Myhrer, Phys. Lett. B533(2002)25.

[4] S. Ando, F. Myhrer, V. Gudkov, T. Sato, H. W. Fearing, S. Nakamura, K. Kubodera, *"Neutron beta decay in effective field theory"*, in preparation.

EFT's for the Inverse-Square Potential and Three-Body Problem

Michael C. Birse and Thomas Barford

Department of Physics and Astronomy, University of Manchester, Manchester, M13 9PL, UK

Compared with EFT's for the corresponding two-body systems, it has proved much more complicated to determine the power counting for three-body systems with short-range forces and large two-body scattering lengths. This is particularly true of attractive systems such as three bosons or the triton.

As shown by Efimov [1], the wave functions at short distances satisfy a 2D Schrödinger equation with an attractive $1/r^2$ potential. These wave functions are ill-defined without a boundary condition to ensure that no flux is lost at the centre and to fix the phase of the wave functions for small r (a self-adjoint extension). The resulting wave functions have the form $\sin(s_0 \ln r/R_0)$ as $r \to 0$.

Using the distorted-wave RG [2], we apply a cut-off at $E = \pm\Lambda^2/M$ on the distorted waves and bound states of this potential, and follow the RG flow of the three-body forces as $\Lambda \to 0$. The oscillatory behaviour of the wave functions means that the potential tends to a limit cycle. Since the amplitude of these oscillations is constant as $r \to 0$, the power counting for perturbations is the same as that for a 2D system with zero angular momentum. This implies that the leading, energy-independent interaction is marginal [3].

We find the following power counting for these systems. At LO (Λ^0) there is the marginal three-body force. This corresponds to the RG limit cycle and it determines the phase of the wave functions as $r \to 0$. At NLO (Λ^1) the two-body effective range appears as a perturbation. Then at NNLO (Λ^2) we have an energy-dependent three-body force, along with the two-body effective range at second order. This agrees with the counting obtained by Bedaque et al. [4] using the Skorniakov–Ter-Martirosian equation with a momentum cut-off.

References

[1] V. Efimov, Sov. J. Nucl. Phys. **29** (1971) 589.

[2] T. Barford and M. C. Birse, Phys. Rev. **C67** (2003) 064006 [hep-ph/0206146].

[3] T. Barford and M. C. Birse, Few Body Syst. Suppl. **14** (2003) 123 [nucl-th/0210084]; and in preparation.

[4] P. F. Bedaque, H. W. Griesshammer, H.-W. Hammer and G. Rupak, Nucl. Phys. **A714** (2003) 589 [nucl-th/0207034].

DEAR - Kaonic Hydrogen: First Results

M. Cargnelli[d], G. Beer[i], A.M. Bragadireanu[a,e], C. Curceanu (Petrascu)[a,e], J.-P. Egger[b,c], H. Fuhrmann[d], C. Guaraldo[a], M. Iliescu[a], T. Ishiwatari[d], K. Itahashi[g], M. Iwasaki[f], P. Kienle[d], B. Lauss[h], V. Lucherini[a], L. Ludhova[b], J. Marton[d], F. Mulhauser[b], T. Ponta[a,e], L.A. Schaller[b], R. Seki[j,k], D. Sirghi[a], F. Sirghi[a], P. Strasser[f] , J. Zmeskal[d]

[a]INFN - Laboratori Nazionali di Frascati; [b]Universite de Fribourg; [c]Universite de Neuchâtel; [d]Institute for Medium Energy Physics, Vienna; [e]Institute of Physics and Nuclear Engineering, Bucharest; [f]RIKEN, Saitama; [g]Tokyo Institute of Technology; [h]University of California and Berkeley; [i] University of Victoria; [j]California Institute of Technology; [k]California State University

The DEAR[1] experiment [1] measures the energy of X-rays emitted in the transitions to the ground states of kaonic hydrogen. The shift ϵ and the width Γ of the 1s state are related to the real and imaginary parts of the complex S-wave scattering length by the Deser Trueman formula.



Figure 2: Background subtracted energy spectrum of kaonic hydrogen. For the first time K_β, K_γ, K_{high} are clearly resolved.

The preliminary results are: ϵ= - 183 $\pm$ 62 eV and Γ= 213 $\pm$ 138 eV. Both values are smaller then those from the previous experiment [2] and consistent with recent theoretical studies [3].

Part of the work was supported by "Transnational access to Research Infrastructure" (TARI) Contract No. HPRI-CT-1999-00088.

References

[1] S. Bianco et al., Rivista del Nuovo Cimento 22, No.11, (1999), 1.

[2] M. Iwasaki et at., Phys. Rev. Lett. 78 (1997), 3067.

[3] A. Ivanov et al., submitted to Eur. Phys. J. A, nucl-th/0310081.

[1]DAΦNE Exotic Atom Research, conducted at the Frascati electron positron collider

Comments on $\bar{K}N$ Scattering

Jürg Gasser

Institute for Theoretical Physics, University of Bern
Sidlerstrasse 5, CH-3012 Bern, Switzerland

The DEAR collaboration has investigated the ground state of $\bar{K}p$ atoms, in particular, the energy level shift and width ϵ_{1s} due to strong interactions [1]. The quantity ϵ_{1s} is related to the elastic threshold amplitude $T^{\rm th}_{\bar{K}p}$ [2],

$$\epsilon_{1s} = -2\alpha^3 \mu_r^2 T^{\rm th}_{\bar{K}p}(1+X)\,. \tag{1}$$

Here, $T^{\rm th}_{\bar{K}p}$ denotes the amplitude in pure QCD in the isospin symmetry limit $m_u = m_d$, and the quantity X stands for isospin symmetry breaking corrections. These are presently evaluated [3] in the framework of effective non relativistic quantum field theories [4]. To work out the implication of a precise measurement of ϵ_{1s} on the kaonic sigma terms [5] requires a complex theoretical analysis. I have suggested in my talk a more modest aim as an intermediate step: Calculate the amplitude $T^{\rm th}_{\bar{K}p}$ as precisely as possible, based on a framework that relies on QCD (incorporating chiral symmetry at least), and compare with (1). Available predictions [6,7] differ considerably - it would be instructive to investigate how these calculations can be made more precise.

References

[1] See the contribution of M. Cargnelli at this workshop.

[2] S. Deser, M. L. Goldberger, K. Baumann and W. Thirring, Phys. Rev. **96** (1954) 774; T. L. Trueman, Nucl. Phys. **26** (1961) 57.

[3] Ulf-G. Meißner, U. Raha and A. Rusetsky, work in progress.

[4] See the contributions of J. Schweizer and P. Zemp at this workshop.

[5] S. Bianco *et al.* [DEAR Collaboration], Riv. Nuovo Cim. **22N11** (1999) 1.

[6] N. Kaiser, P. B. Siegel and W. Weise, Nucl. Phys. A **594** (1995) 325 [arXiv:nucl-th/9505043].

[7] J. A. Oller and U. G. Meißner, Phys. Lett. B **500** (2001) 263 [arXiv:hep-ph/0011146].

On Pionic and Kaonic Hydrogen

A. N. Ivanov, M. Cargnelli, M. Faber, A. Hirtl, J. Marton, N. I. Troitskaya, and J. Zmeskal

Atominstitute, Vienna University of Technology and Institute of Medium Energy Physics, Austrian Academy of Sciences,, Vienna, Austria

In [1] the energy level displacement of the ground state of hadronic hydrogen A_{hp}, where $h = \pi$ or K, has been expressed in terms of the amplitude $M(h^-p \to h^-p)$ of h^-p scattering, weighted with the wave function Φ_{1s} of hadronic hydrogen in the ground state

$$\begin{aligned} -\epsilon_{1s} + i\,\frac{\Gamma_{1s}}{2} &= \frac{1}{4m_{h^-}m_p}\int\frac{d^3k}{(2\pi)^3}\int\frac{d^3q}{(2\pi)^3}\sqrt{\frac{m_{h^-}m_p}{E_{h^-}(\vec{k}\,)E_p(\vec{k}\,)}}\sqrt{\frac{m_{h^-}m_p}{E_{h^-}(\vec{q})E_p(\vec{q})}} \\ &\times \Phi^\dagger_{1s}(\vec{k}\,)\,M(h^-(\vec{k}\,)p(-\vec{k},\sigma_p) \to h^-(\vec{q})p(-\vec{q},\sigma_p))\,\Phi_{1s}(\vec{q}). \qquad (1) \end{aligned}$$

The knowledge of the amplitude of h^-p scattering for arbitrary relative momenta should give the possibility to calculate explicitly the energy level displacement. The low–energy limit $k, q \to 0$, justified by the wave functions $\Phi^\dagger_{1s}(\vec{k}\,)$ and $\Phi_{1s}(\vec{q})$, reduces (1) to the DGBT formula [2]. In [1] (nucl–th/0310027) formula (1) has been generalized to arbitrary excited $n\ell$ states of the hadronic atom A_{hp}. In [1] (nucl–th/0310081) we have suggested a model of low–energy K^-p scattering near threshold of the K^-p pair. We have assumed that the amplitude of K^-p scattering near threshold is fully defined by the contribution of three resonances $\Lambda(1405)$, $\Lambda(1800)$ and $\Sigma(1750)$ and a smooth elastic background. We predict the energy level displacement of the ground state of kaonic hydrogen [1] $-\epsilon_{1s} + i\Gamma_{1s}/2 = (-203 \pm 15) + i\,(113 \pm 8)$ eV fitting well preliminary experimental data by the DEAR Collaboration $-\epsilon^{\rm exp}_{1s} + i\Gamma^{\rm exp}_{1s}/2 = (-183 \pm 62) + i\,(106 \pm 69)$ eV [3].

References

[1] A. N. Ivanov, M. Faber, A. Hirtl, J. Marton, and N. I. Troitskaya, nucl–th/0306047 (to appear in EPJA), nucl–th/0310027, (to appear in EPJA); A. N. Ivanov, M. Cargnelli, M. Faber, J. Marton, N. I. Troitskaya, and J. Zmeskal, nucl–th/0310081.

[2] S. Deser, M. L. Goldberger, K. Baumann, and W. Thirring, Phys. Rev. **96**, 774 (1954).

[3] M. Cargnelli (the DEAR Collaboration), Plenary talk at Workshop on *Chiral Dynamics*, Bonn, Germany, 8–13 September 2003.

Deser-type Formula for Pionic Hydrogen

Peter Zemp

Institute for theoretical Physics, University of Bern,
Sidlerstrasse 5, 3012 Bern, Switzerland

The decay width of the ground state of pionic hydrogen is dominated by the two decay channels $(\pi^- p)_{1s} \to \pi^0 n$ and $(\pi^- p)_{1s} \to n\gamma$. Using δ as a common counting for the light quark mass difference $m_u - m_d$ and for the fine-structure constant α, the leading term of the former (latter) decay channel is of order $\delta^{7/2}$ (δ^4). The decay width through these two channels can be expressed by a Deser-type formula

$$\Gamma_{1s} = 4\,\alpha^3 M_r^2 q_0 \left(1 + \frac{1}{P}\right) [a_{\pi^- p \to \pi^0 n}(1 + \delta_\Gamma)]^2 \,. \tag{1}$$

Since the Panofsky ratio $P = \sigma(\pi^- p \to \pi^0 n)/\sigma(\pi^- p \to n\gamma)$ is about 1.5, the $n\gamma$ channel amounts to a 60% correction. To the best of my knowledge, this formula was first established in the framework of potential models [1]. I want to verify its validity in QCD + QED, using the technique of non-relativistic effective field theories [2]. Constructing the effective Lagrangian and solving the master equation, one encounters the obstacle that the photon in the intermediate state carries a hard momentum of order M_π. To circumvent this problem, I do not incorporate $n\gamma$ intermediate states with effective fields. Instead I introduce a non-hermitian contact term [2] (cf. also [3]) in the Lagrangian with the coupling $d_1^R + i d_1^I$,

$$\mathcal{L}_{\rm I}^{n\gamma} = (d_1^R + i d_1^I)\; \psi^\dagger(x)\psi(x)\; \pi_-^\dagger(x)\pi_-(x) \,. \tag{2}$$

The imaginary part d_1^I replaces the imaginary part generated by the intermediate $n\gamma$ state. With this contact term, I can prove that formula (1) is valid at order δ^4. It remains to evaluate Γ_{1s} at order $\delta^{9/2}$ and to determine δ_Γ in equation (1) [4].

References

[1] G. Rasche and W. S. Woolcock, Helv. Phys. Acta **49** (1976) 557; D. Sigg *et al.*, Nucl. Phys. A **609** (1996) 269.

[2] W. E. Caswell and G. P. Lepage, Phys. Lett. B **167** (1986) 437; V. Antonelli, A. Gall, J. Gasser and A. Rusetsky, Annals Phys. **286** (2001) 108 [arXiv:hep-ph/0003118]; V. E. Lyubovitskij and A. Rusetsky, Phys. Lett. B **494** (2000) 9 [arXiv:hep-ph/0009206].

[3] J. Gasser, V. E. Lyubovitskij, A. Rusetsky and A. Gall, Phys. Rev. D **64** (2001) 016008 [arXiv:hep-ph/0103157].

[4] P. Zemp, work in progress.

Atoms Consisting of $\pi^+\pi^-$ and πK Mesons

L.Nemenov[1]

[1]JINR, Dubna

Using experience obtained in the DIRAC experiment at CERN [1] new experiments are proposed for CERN PS and J-PARC in Japan [2] to check precisely predictions of low energy QCD.

These experiments aim to measure the $\pi^+\pi^-$ atom ($A_{2\pi}$) lifetime with precision better than 6% and to determine the S-wave $\pi\pi$ scattering length difference $|a_0 - a_2|$ at 3%. Simultaneously with the $A_{2\pi}$ investigation DIRAC plans to observe πK atoms ($A_{\pi K}$) and to measure their lifetime at the 20% level in order to evaluate the S-wave πK scattering length difference $|a_{1/2} - a_{3/2}|$ at 10%.

Furthermore the observation of long-lived (metastable) $A_{2\pi}$ states is also envisaged to be done in the same setup. This allows to measure the energy difference between nS and nP states and to determine in a model-independent way $2a_0 + a_2$.

Low energy QCD [3] predicts $\pi\pi$ scattering lengths at 2% [4,5] and πK scattering lengths at 10% [6,7]. These scattering lengths have never been verified experimentally with the same accuracy as theoretically predicted.

The theoretical results have been obtained assuming strong condensation of quark-antiquark pairs in the vacuum. Therefore the proposed experiments will test crucially low energy QCD predictions and accordingly our understanding of the nature of the QCD vacuum [8].

References

[1] B. Adeva *et al.*, CERN/SPSLC 95-1, SPSLC/P 284, Geneva (1995).

[2] B. Adeva *et al.*, Lifetime measurement of $\pi^+\pi^-$ and $\pi^\pm K^\pm$ atoms to test low energy QCD (2002) Letter of Intent of the experiment at Japan Proton Accelerator Research Complex.

[3] S. Weinberg, Physica A **96** (1979) 327; J. Gasser and H. Leutwyler, Phys. Lett. B **125** (1983) 325; ibid Nucl. Phys. B **250** (1985) 465, 517, 539

[4] G. Colangelo, J. Gasser and H. Leutwyler, Phys. Lett. B **488** (2000) 261.

[5] G. Colangelo, J. Gasser and H. Leutwyler, Nucl. Phys. B **603** (2001) 125.

[6] V. Bernard, N. Kaiser and U.-G. Meißner, Phys. Rev. D **43** (1991) 2757. V. Bernard, N. Kaiser and U.-G. Meißner, Nucl. Phys. B **357** (1991) 129.

[7] A. Roessl, Nucl. Phys. B **555** (1999) 507; (hep-ph/9904230)

[8] M. Knecht *et al.*, Nucl. Phys. B **457** (1995) 513.

Deeply Bound Pionic Atoms: Optical Potential at $O(p^5)$ in ChPT

Luca Girlanda[1], Akaki Rusetsky[2,3] and Wolfram Weise[1,4]

[1]ECT*, Strada delle Tabarelle 286, I-38050 Villazzano (Trento), Italy
[2]HISKP (Theorie), University of Bonn, Nußallee 14-16, 53115 Bonn, Germany
[3]HEPI, Tbilisi State University, University St. 9, 380086 Tbilisi, Georgia
[4]Physik-Department, TU München, D-85747 Garching, Germany

Deeply bound pionic atoms can be considered as a laboratory to test the effects of baryon density on chiral symmetry breakdown [1]. The present effective field theory approach to this problem is provided by the in-medium ChPT [2], which treats the nucleus as an infinite and uniform Fermi-sea of protons and neutrons. The results are then expressed through a local Fermi momentum $\vec{k}(\vec{r})$ [3] with some (arbitrary) prescription. Such arbitrariness is worrisome especially for the case at hand, in which the bound state wave function is picked at the nuclear surface, due to a balance between Coulomb attraction and strong repulsion.

In Ref. [4] we give a consistent formulation of ChPT in a non-uniform fermionic background, corresponding to the (finite) nucleus. The nuclear structure information is encoded in a set of nuclear matrix elements of free-nucleon field operators. Chiral counting applied to the above matrix elements allows to reduce considerably the nuclear input needed. As an application we calculate the charged pion self-energy in the background of a heavy nucleus at $O(p^5)$ of the chiral expansion, including consistently all isospin-breaking effects arising at this order. Finite nuclear size effects in the pion-nucleus optical potential are identified unambiguously, without any ad-hoc prescription such as local density approximation.

However, in order to be relevant phenomenologically, this analysis has to be extended to $O(p^6)$, beyond the linear density approximation for the optical potential. We plan to address this issue in future publications.

References

[1] W. Weise, Nucl. Phys. A **690** (2001) 98; P. Kienle and T. Yamazaki, Phys. Lett. B **514** (2001) 1.

[2] N. Kaiser and W. Weise, Phys. Lett. B **512** (2001) 283; U. G. Meißner, J. A. Oller and A. Wirzba, Annals Phys. **297** (2002) 27; T. S. Park, H. Jung and D. P. Min, J. Korean Phys. Soc. **41** (2002) 195 (2002).

[3] E. E. Kolomeitsev, N. Kaiser and W. Weise, Phys. Rev. Lett. **90** (2003) 092501; Nucl. Phys. A **721** (2003) 835.

[4] L. Girlanda, A. Rusetsky and W. Weise, hep-ph/0311128.

All-Order Low-Energy Expansion in the 3-Body System

Harald W. Grießhammer[1]

[1] Institut für Theoretische Physik (T39), TU München, *and* ECT*, Trento

In "pion-less" Effective Field Theory, the power-counting in the 3-body system with a shallow two-body bound state was extended and systematised to all orders in the low-energy expansion in Ref. [1]. The analytical considerations rest on the tenet that a 3-body force (3BF) is included if and only if necessary to cancel cut-off dependences in observables at a given order. The one momentum-independent 3BF needed for cut-off independence up to NLO in the ${}^2\mathrm{S}_{\frac{1}{2}}$ (triton) channel is determined by the three-body scattering length. One and only one new parameter enters at NNLO, namely the Wigner-$SU(4)$-symmetric 3BF with two derivatives, fixed to reproduce the triton binding energy. Only the $SU(4)$-symmetric 3-body forces are systematically enhanced over their contributions found from a naïve dimensional estimate, with a $2n$-derivative 3BF entering at the $2n$th order. A new, computationally convenient and simple scheme to perform higher-order calculations in the three-body system iterates a kernel to all orders, which has been expanded to the desired order of accuracy. The *nd* scattering phase shifts agree well with phase shift analysis and modern potential model calculations.

In contradistinction, Wigner-symmetric 3BFs are suppressed compared to naïve dimensional analysis in all other partial waves [2]: In the lth partial wave of the spin doublet ($\lambda = 1$) and quartet ($\lambda = -\frac{1}{2}$) channels, the amplitude converges for large off-shell momenta p as p^{-s_0-1}, with s_0 the solution to

$$1 \;=\; (-)^l \,\frac{2^{1-l}\,\lambda}{\sqrt{3\pi}}\,\frac{\Gamma[\frac{l+1+s_0}{2}]\,\Gamma[\frac{l+1-s_0}{2}]}{\Gamma[l+\frac{3}{2}]}\;\;{}_2F_1\left[\frac{l+1+s_0}{2},\frac{l+1-s_0}{2};l+\frac{3}{2};\frac{1}{4}\right] \quad .$$

Except in the ${}^2\mathrm{S}_{\frac{1}{2}}$ channel, the asymptotics is always weaker than in the naïve analysis ($s_0 = 1$). Thus, effective range corrections in the two-body system containing n external momenta need 3BFs with m derivatives to absorb UV divergences only when $n \geq \mathrm{Re}[m+2s_0]$. For example, the first Wigner-symmetric 3BF in the ${}^4\mathrm{S}_{\frac{3}{2}}$ channel is demoted from N^{4}LO to N^{7}LO, suggesting that two-body physics rules this channel to very high accuracy.

Sponsored in part by DFG Sachbeihilfe GR 1887/2-1 and BMBF.

References

[1] P. F. Bedaque, G. Rupak, H. W. Grießhammer and H.-W. Hammer, Nucl. Phys. **A714** (2003), 589 [nucl-th/0207034].

[2] H. W. Grießhammer, *Three-Body Forces in the Higher Partial Waves of the Loosely Bound Three-Body System at Very Low Energies*; forthcoming.

New Results in the Three and Four Nucleon Systems

Timothy Black

Department of Physics, University of North Carolina at Wilmington
Wilmington, NC, USA

Recent measurements of the n-d and n-^{3}He bound coherent scattering lengths, conducted at the Neutron Interferometer and Optics Facility at NIST, are in significant disagreement with exact calculations of these observables using modern NN+3NF potentials[1,2]. The bound coherent n-d scattering length was measured to be $b_{nd} = (6.665 \pm 0.004)$ fm, which yields a new world average for this parameter of $b_{nd} = (6.669 \pm 0.003)$ fm[3,4]. The bound coherent scattering length is related to the free doublet and quartet scattering lengths by $b_{nd} = \frac{m_D+m_n}{m_D}\left(\frac{1}{3}a_2 + \frac{2}{3}a_4\right)$. Combining the new world average for b_{nd} with calculated values of ${}^4a_{nd}$ yields a value for the free doublet scattering length of ${}^2a_{nd} = (0.645 \pm 0.003[\text{expt}] \pm 0.007[\text{thry}])$ fm.

In the n-^{3}He system, the bound coherent scattering length was measured as $b_{n^3\text{He}} = (5.857 \pm 0.007)$ fm, which yields a new world average for this parameter of $b_{n^3\text{He}} = (5.854 \pm 0.007)$ fm. Combining these measurements with Zimmer's result for the bound incoherent scattering length of $b_i = (-2.365 \pm 0.020)$ fm[5], one obtains the free singlet and triplet scattering lengths for the n-^{3}He system; $a_0 = (7.456 \pm 0.026)$ fm, and $a_1 = (3.364 \pm 0.010)$ fm, respectively. The extreme precision of these new measurements in the n-d and n-^{3}He systems renders them suitable for use in high-order, contemporary EFT calculations.

References

[1] H. Witała, A. Nogga, H. Kamada, W. Glöckle, J. Golak, R. Skibiński, arXiv: (2003) nucl-th/0305028v1.

[2] H. M. Hofmann and G. M. Hale, Phys. Rev C68 (2003) 021002(R).

[3] T. C. Black, P. R. Huffman, D. L. Jacobson, W. M. Snow, K. Schoen, M. Arif, H. Kaiser, S. K. Lamoreaux, and S. A. Werner, Phys. Rev. Lett. 90:19, (2003) 192502.

[4] K. Schoen, D. L. Jacobson, M. Arif, P. R. Huffman, T. C. Black, W. M. Snow, S. K. Lamoreaux, H. Kaiser, and S. A. Werner, Phys. Rev. C 67:4 (2003) 044005.

[5] O. Zimmer, G. Ehlers, B. Farago, H. Humblot, W. Ketter, and R. Scherm, (2002) EPJdirect **A1**.

Nd Scattering at Low Energies

H. Paetz gen. Schieck

Institut für Kernphysik, Universität zu Köln,
Zülpicher Straße 77, D-50937 Köln, Germany

Despite the tremendous progress made by few-body theory — by numerically exact calculations using precision meson-exchange nucleon-nucleon potentials including different three-body forces and, recently, in applying realistic effective-field theory — a number of long-standing low-energy puzzles are still unsolved. These are: the d+N elastic cross section **(Sagara) anomaly**, the Nd elastic $\mathbf{A}_y$ **(and iT$_{11}$) puzzle**, the Nd breakup anomaly in the **space-star (SST)** configuration, and the new Nd breakup anomaly in the **quasi-free scattering (QFS)** configuration.

For the pd system the data base is much larger, with mostly better-quality data, than for the nd system. For a general survey see [1]. In order to compare low-energy pd data to realistic calculations inclusion of the Coulomb force is mandatory. This has been achieved for Faddeev calculations of elastic scattering observables (see e.g. Fig. 1) whereas for the pd breakup this is still eagerly awaited. Some breakup data (e.g. in the SST and QFS, but not in the final-state interaction configurations) exhibit large Coulomb effects [2]. Fig. 2 shows the low-energy systematics of nd vs. pd SST cross sections. The unsolved puzzles call for new theoretical efforts with different three-body forces or modifications of the two-nucleon input. Also measurements of new observables and in larger phase-space areas may be useful.



Fig. 1: Polarization transfer coefficients compared to realistic calculations, also including the UR IX three-body and the Coulomb force [3]. Fig. 2: Comparison of nd vs. pd breakup cross sections in the symmetric space-star configuration at various energies.

References

[1] W. Glöckle et al., Phys. Rep. 274 (1996) 107.
[2] H. Paetz gen. Schieck et al., Few-Body Systems 30 (2001) 81.
[3] A. Kievsky et al., Phys. Rev. C64 (2001) 041001(R) and priv. comm.

Nucleon-Deuteron Scattering at Low Energies

W. Tornow

Department of Physics, Duke University and
Triangle Universities Nuclear Laboratory, Durham, NC 27708, USA

We discuss a new twist to the long-standing nucleon-deuteron analyzing power puzzle. The new low-energy neutron-deuteron $A_y(\theta)$ data at $E_n = 1.2$ and 1.9 MeV [1] revealed a sizeable difference when compared to proton-deuteron data. This difference cannot be accounted for by the Coulomb interaction (in the case of *p*-*d* scattering) and it is orders of magnitude larger than expected from charge-symmetry breaking effects in the $^3\mathrm{P}_j$ NN interactions. We speculate that the experimentally observed effect is due to the different magnetic moment interactions between neutrons and protons and the deuteron target nuclei. Recent calculations support this conjecture [2,3]. The proton-deuteron angular distribution for $A_y(\theta)$ is considerably affected by the magnetic moment interaction, while for neutron-deuteron scattering the effect is restricted to very forward angles.

We present proton-deuteron phase-shift analysis results [4] that clearly show that the discrepancy between the experimental $^4\mathrm{P}_J$ proton-deuteron 3N phase shifts and the theoretically predicted ^4P_J phase shifts (i.e., the A_y puzzle) increases with increasing energy. This is in contrast to the observation that the discrepancy between $A_y(\theta)$ data and calculations decreases with increasing energy. This finding suggests that $A_y(\theta)$ is not dominated anymore at higher energies by the $^4\mathrm{P}_J$ 3N phase shifts. Sensitivity calculations for $A_y(\theta)$ with varying $^4\mathrm{P}_J$ phase shifts support this conjecture.

We argue that the 3N $A_y(\theta)$ puzzle cannot be solved by concentrating on the 3N system alone. In the 4N system the *p*-^{3}He elastic scattering $A_y(\theta)$ shows an even larger discrepancy (40%) between data and calculations than observed in the 3N system (25%). $A_y(\theta)$ data for *n*-^{3}He elastic scattering are needed to make further progress.

References

[1] E.M. Neidel *et al.*, Phys. Lett. B552 (2003) 29.

[2] H. Witała *et al.*, Phys. Rev. C67 (2003) 064002.

[3] A. Kievsky *et al.*, Phys. Rev. C, in press.

[4] Z. Chen *et al.*, submitted to Few-Body Systems.

Three-Nucleon System Dynamics Studied in the dp-Breakup at 130 MeV

R. Bieber[3], A. Biegun[2], K. Bodek[1], K. Ermisch[3], W. Glöckle[4], J. Golak[1], N. Kalantar-Nayestanaki[3], **St. Kistryn**[1], M. Kiš[3], J. Kuroś-Żołnierczuk[1], A. Micherdzińska[2], M. Mahjour-Shafiei[3], R. Skibiński[1], E. Stephan[2], H. Witała[1], J. Zejma[1] and W. Zipper[2]

[1]Institute of Physics, Jagellonian University, 30059 Kraków, Poland

[2]Institute of Physics, University of Silesia, 40007 Katowice, Poland

[3]Kernfysisch Versneller Instituut, 9747 AA Groningen, The Netherlands

[4]Institut für Theoretische Physik II, Ruhr Universität Bochum, 44780 Bochum, Germany

The measurement of exclusive deuteron-proton breakup, employing a detection system covering large fraction of the available phase space and using 130 MeV polarized deuteron beam has been performed at KVI Groningen. Theoretical calculations [1] predict large but generally model dependent three-nucleon force (3NF) effects in certain regions of the breakup phase space for different experimentally attainable observables.

In the first step of data analysis [2] high precision fivefold cross-section values in 38 kinematical configurations have been extracted, using for normalization the simultaneously measured elastic dp-scattering data. The results have been compared to predictions of modern nuclear forces. To this aim the three-nucleon (3N) Faddeev equations have been solved rigorously using realistic NN potentials AV18, CD Bonn, Nijm I and Nijm II alone, and combining them with the 2π-exchange Tucson-Melbourne (TM) 3NF and with its modified version TM99, more consistent with chiral symmetry. Global comparison of the measured cross sections to pairwise NN force predictions only and with 3NF's included clearly reveals the presence of 3NF effects.

All details of our study, showing the usefulness of the kinematically complete breakup reaction measured in the full phase space to search for 3NF effects, are presented in Ref. [3]. Further insight into details of the 3N interaction dynamics is expected from the undergoing analysis of the vector and tensor analyzing powers.

References

[1] J. Kuroś-Żołnierczuk *et al.*, Phys. Rev. C **66** (2002) 024003 and 024004.

[2] A. Micherdzińska, Ph.D. thesis, University of Silesia, Katowice 2003.

[3] St. Kistryn *et al.*, Phys. Rev. C **68** (2003) in print.

Few-Body Studies at KVI

J.G. Messchendorp[1]

[1]Kernfysisch Versneller Instituut, Zernikelaan 25,
9747 AA Groningen, The Netherlands

The KVI facility allows a detailed study of few-nucleon interactions below the pion-production threshold exploiting polarized proton and deuteron beams up to energies of 190 MeV. The experimental and theoretical studies of bremsstrahlung in the two-nucleon scattering processes have a long-standing tradition at KVI. The first experimental efforts in this field date from 1995 where the $pp \rightarrow pp+\gamma^{(*)}$ channel was measured with an incident-proton beam of 190 MeV using a coincidence setup consisting of the hadron detector SALAD [1] and the photon spectrometer TAPS [2]. The setup allowed to determine the real (γ) [3] and virtual ($\gamma^* \rightarrow e^+e^-$) [4] bremsstrahlung yields. Modern state-of-the-art microscopic calculations based on a fully-relativistic framework are not able to describe the measured bremsstrahlung data in the two-nucleon system.

The focus of the few-body program at KVI has recently shifted towards the understanding of the three-nucleon system by exploiting $p+d$ and $d+p$ reactions with polarized proton and deuteron beams. Different final states have been observed which includes the elastic, break-up, and radiative capture reactions. A systematic study of the elastic $p+d$ scattering reaction has been carried out in which high-precision cross sections and analyzing powers have been determined for several bombarding energies up to 190 MeV using the Big-Bite Spectrometer (BBS) [5]. The data show that effective models based on a well-understood two-nucleon interaction are not sufficient to describe the three-nucleon pd system [6]. Including the most modern type of three-nucleon potentials in the calculations resolves only a part of the discrepancies.

References

[1] N. Kalantar-Nayestanaki, Nucl. Inst. Meth. **A444** (2000) 591.

[2] A. Gabler et al., Nucl. Instr. and Meth. **A346** (1994) 168.

[3] H. Huisman et al., Phys. Rev. Lett. **83** (1999) 4017, Phys. Lett. **B519** (2001) 8, Phys. Rev. C **65** (2002) 031001.

[4] J.G. Messchendorp et al., Phys. Rev. Lett. **82** (1999) 2649, Phys. Rev. Lett. **83** (1999) 2530, Phys. Rev. **C61** (2000) 064007.

[5] A.M. van den Berg, Nucl. Instr. and Meth. **B99** (1995) 637.

[6] K. Ermisch et al., Phys. Rev. Lett. **86** (2001) 5862.

dp Elastic Scattering Experiments

Hide Sakai

Dep. Phys./CNS, University of Tokyo,
Hongo 7-3-1, Bunkyo, Tokyo, Japan

Recent experimental results to study the three-nucleon-force ($3NF$) effects in the dp elastic scattering at intermediate energies are reported.

One of the new measurement at RIKEN is the deuteron-to-proton polarization transfer (PT) coefficients for the dp elastic scattering at 135 MeV/u by Sekiguchi *et al.* [1]. Very precise data for $K_y^{y'}$, $K_{xx}^{y'}$, $K_{yy}^{y'}$ and $K_{xz}^{y'}$ are obtained for $\theta_{cm} = 90^\circ - 180^\circ$.

Including those new data, comparisons between the existing data [2,3] at 135 MeV/u and the Faddeev calculations are made.

- $d\sigma/d\Omega$ and iT_{11} are well reproduced by including $3NF$, which indicates the clear signature of $3NF$s.
- Spin observables (T_{20}, T_{21},T_{22}, $P^{y'}$ and PT coefficients) are not necessarily reproduced by including $3NF$, which shows deficiency of present day 2π exchange type $3NF$ models.
- Particularly the TM $3NF$ does a poor job while updated versions TM'(99) $3NF$s respecting chiral symmetry do much better jobs similar to UR-IX $3NF$.
- Comparisons are also made between data at 70 MeV/u [3] and recent calculations based on the chiral effective field theory [4], which does a reasonably good job for $d\sigma/d\Omega$ but not much for tensor analyzing powers.

Data for the $\vec{n}d$ [5] and $\vec{p}d$ [6] at 250 MeV/u at RCNP are also presented and compared with calculations.

References

[1] K. Sekiguchi *et al.*, to be published.

[2] H. Sakai *et al.*, Phys. Rev. Lett. 84 (2000) 5288.

[3] K. Sekiguchi *et al.*, Phys. Rev. C65 (2002) 034002.

[4] E. Epelbaum *et al.*, Phys. Rev. C66 (2002) 064001.

[5] Y. Maeda *et al.*, to be published.

[6] K. Hatanaka *et al.*, Phys. Rev. C66 (2002) 044002.

Nuclear Binding Energies and Quark Masses

John F. Donoghue

Dept. of Physics, University of Massachusetts,
Amherst, MA 01003, USA

In this talk, I described some of my recent work on the nuclear force and its dependence on quark mass [1]. My initial motivation was in connection with a project on the equivalence principle and dilaton couplings [2]. For a dilaton that couples in a way similar to a quark mass term, the least understood effects of its violation of the equivalence principle is the effects that comes from nuclear binding energies. These can be understood if one understands the quark mass effects on binding. However, the results obtained are interesting in their own right in the context of the nuclear force.

The main innovation was to use the Omnes function technique [3] in the dispersive integral that determines the scalar-isoscalar nuclear potential [4]. The potential is calculated in terms of its imaginary part, and the low energy contributions to the imaginary part are known in chiral perturbation theory. However, as one considers these at higher energies, the chiral results quickly blow up. Here the Omnes technique becomes useful. By matching the Omnes function to the chiral results at low energy one can provide an extension to higher energies that is well behaved. The matching is given by

$$\rho_S(\mu) = \rho_S^{NNLO} Re\Omega(\mu) + \rho_S^{NNNLO} |\Omega(\mu)|^2 \tag{1}$$

where Ω is the Omnes function. Interestingly this representation leads to a nuclear potential which is very similar to a conventional σ potential, and also emerges with about the right strength. This provides a novel understanding of this ill-understood component of the nuclear force. In addition, knowing these ingredients one can explore the quark mass dependence in a way that was not possible before.

References

[1] J.F. Donoghue (to appear).

[2] T. Damour and J.F. Donoghue (to appear)

[3] J. F. Donoghue, J. Gasser and H. Leutwyler, Nucl. Phys. B **343**, 341 (1990).

[4] The literature can be traced through E. Epelbaum, W. Glöckle and U. G. Meißner, arXiv:nucl-th/0308010.

Proton-Proton Scattering Experiments

Frank Rathmann

Institut für Kernphysik, Forschungszentrum Jülich, 52425 Jülich, Germany

The lack of high-quality elastic proton-proton scattering data at the beginning of the 1990's hampered both the further improvement of potential models of the NN interaction and the refining of partial wave analyses. A precise pp data base was also needed to provide polarimetry for the analysis of inelastic reaction channels. Through a series of measurements at the Indiana Cooler with stored polarized protons impinging on a polarized hydrogen storage cell target, a new method to measure spin correlation parameters could be established. The measurements yielded full angular distributions of analyzing powers A_y and spin correlation parameters A_{xx}, A_{yy} and A_{xz} at eight different energies between 197 and 449 MeV [1], the fourth spin correlation parameter A_{zz} was measured at 197 MeV only [2]. At higher energies, without a storage cell, the EDDA collaboration achieved high-quality data of pp excitation functions of cross sections, analyzing powers, and spin correlation parameters between 200 MeV and 2.5 GeV at COSY [3]. Our theoretical understanding of the NN system above 1 GeV is still unsatisfactory [4].

But understanding the NN interaction involves *both* isospin channels. While for the pp system the data situation today can be called fair, the pn database up to 3 GeV is almost empty. Although not ideally suited for polarization experiments, some pn observables can be measured with the ANKE dipole spectrometer at COSY. Through the detection of spectator protons from a polarized deuterium gas target that is bombarded by polarized protons, pn elastic scattering at cm angles below $\theta = 30^\circ$ is accessible. There, the pn data base is almost empty at all energies [5].

References

[1] W. Haeberli *et al.*, Phys. Rev. **C 55** (1997) 597; F. Rathmann *et al.*, Phys. Rev. **C 58** (1998) 658; B.v. Przewoski *et al.*, Phys. Rev. **C 58** (1998) 1897.

[2] B. Lorentz *et al.*, Phys. Rev. **C 61** (2000) 054002.

[3] D. Albers *et al.*, Phys. Rev. Lett. **78** (1997) 1652; M. Altmeier *et al.*, Phys. Rev. Lett. **85** (2000) 1819; F. Bauer *et al.*, Phys. Rev. Lett. **90** (2003) 142301.

[4] K. O. Eyser, R. Machleidt, W. Scobel, the EDDA Collaboration, nucl-th/0311002.

[5] Abundance plots of the pp and np databases, made available to us by Dr. Mart Rentmeester `mart.rentmeester@sci.kun.nl`, are available from the website `http://nn-online.sci.kun.nl/NN/nnaplot.html`.

Systematic Approaches to Meson Production in NN and dd Collisions

C. Hanhart

IKP, Forschungszentrum Jülich,
52428 Jülich, Germany

The goal of this presentation was to demonstrate that it is possible to extract interesting physics from NN and dd induced meson production reactions, regardless the large momenta involved. For illustration the techniques/ideas to extract various charge symmetry breaking (CSB) matrix elements was discussed.

One can not expect the same methods to be applicable in reactions at different energies. It was demonstrated recently that pion production can be analyzed within the framework of chiral perturbation theory, once the scheme is adjusted to the large momentum transfer [1]. The expansion parameter turned out to be $\chi = \sqrt{m_\pi/M_N}$. Within this framework an extraction of CSB matrix elements from data on $pn \to d\pi^0$ as well as is straight forward. The corresponding calculations are in progress [2].

As we go to higher energies the straight forward application of chiral perturbation theory can not work any more. Already at the η production threshold the corresponding expansion parameter is 0.7. Therefore at present the extraction of CSB matrix elements in a controlled way from the production of heavier mesons is only possible in a few special situation. In the talk ideas were presented on how is could be possible to extract the CSB amplitude $\eta\alpha \to \pi^0\alpha$ from $dd \to \alpha\pi^0$ near the η production threshold.

The situation is a lot more transparent when it comes to the extraction of the CSB $f_0 - a_0$ mixing matrix element from $pn \to d\pi^0\eta$ and $dd \to \alpha\pi^0\eta$, for in this case the mixing effect is automatically enhanced compared to other CSB effects as they might occur in the production operator, since a_0 and f_0 are rather narrow overlapping resonances [3].

References

[1] C. Hanhart, U. van Kolck and G. A. Miller, Phys. Rev. Lett. **85** (2000) 2905; C. Hanhart and N. Kaiser, production in nucleon nucleon collisions at threshold," Phys. Rev. C **66** (2002) 054005.

[2] U. van Kolck, these proceedings.

[3] C. Hanhart, arXiv:nucl-th/0306073 and references therein.

Vector Meson Properties in Nuclear Matter

Daniel Cabrera[1], E. Oset[1] and M.J. Vicente Vacas[1]

[1]Departamento de Física Teórica and IFIC, Univ. de Valencia-CSIC,
Inst. de Inv. de Paterna, Apdo. Correos 22085, E-46071 Valencia, Spain

In this talk, we report on the properties of vector meson resonances in the nuclear medium. First, the ϕ meson spectrum due to the $\bar{K}K$ channel is obtained in terms of the ϕ selfenergy, which in vacuum is provided by a chiral $SU(3)$ dynamics model. To consider the K and $\bar{K}$ in-medium properties we use the results of previous calculations which account for $S-$ and $P-$wave kaon selfenergies based on the lowest order meson-baryon chiral effective Lagrangian [1]. The $S-$wave kaon selfenergy is built from a selfconsistent coupled channel chiral unitary approach to the $\bar{K}N$ interaction. The $P-$wave kaon selfenergy is driven by the excitation of $\Lambda - h$, $\Sigma - h$ and $\Sigma^*(1385) - h$ components. In addition, a set of vertex corrections is evaluated as demanded by gauge invariance. Within this scheme the mass shift and decay width of the ϕ meson in nuclear matter are studied. We find an increase of the ϕ decay width to around 30 MeV at normal nuclear density and a small mass shift to lower energies which stands below 1 % of the size of the ϕ mass in free space [2].

Second, the ρ meson is studied by looking at the pion-pion scattering amplitude in the vector-isovector channel. We start from a chiral unitary approach to meson-meson scattering, based on the tree level contributions from the lowest order χPT Lagrangian including explicit resonance fields. Low energy chiral constraints are considered by matching our expressions to those of one loop χPT, and unitarity is fulfilled exactly in the frame of the N/D method. To account for the medium corrections, the pion propagators are modified with a $P-$wave selfenergy, driven by the excitation of $p - h$ and $\Delta - h$ components. The terms where the ρ couples to the hadrons in the $p - h$ or $\Delta - h$ excitations are also considered as requested by gauge invariance. In addition, the ρ is allowed to couple to baryonic resonances, particularly to the $S-$wave $N^*(1520)$. The main visible effect of the nuclear medium is an enhancement of the ρ width as well as a small shift of the peak to higher energies as the nuclear density increases. We observe that an important source of strength appears in the ρ spectral function at low energies from the direct coupling to the $N^*(1520) - h$ components [3].

References

[1] E. Oset and A. Ramos, Nucl. Phys. A **679** (2001) 616.

[2] D. Cabrera and M. J. Vicente Vacas, Phys. Rev. C **67** (2003) 045203.

[3] D. Cabrera, E. Oset and M. J. Vicente Vacas, Nucl. Phys. A **705** (2002) 90.

A Renormalization Group Method for Ground State Properties of Finite Nuclei

Janos Polonyi[1,2] and **Achim Schwenk**[3]

[1]Institute for Theoretical Physics, Louis Pasteur University, Strasbourg, France
[2]Department of Atomic Physics, Lorand Eötvös University, Budapest, Hungary

[3]Department of Physics, The Ohio State University, Columbus, OH 43210, USA

The renormalization group (RG) is a powerful method to study complex physical systems with a large number of scales. In nuclear physics, the RG is an ideal tool for constructing effective interactions both in the two-nucleon system (leading to a model-independent NN interaction at low momenta, called $V_{\mathrm{low}\,k}$ [1]) and in many-body systems, such as for Fermi liquids [2,3].

In this talk, we propose a novel method for microscopic calculations of nuclear ground state (gs) properties within the framework of density functional theory. Starting from non-interacting nucleons in a harmonic oscillator background potential $(1-\lambda)V_\Omega$ with a control parameter λ, we use the RG to gradually turn on the two-body interaction $\lambda V_{\mathrm{low}\,k}$, while the background potential is removed. The RG evolution for $0 \leqslant \lambda \leqslant 1$ is constructed to follow the minimum of the density functional $\Gamma_\lambda[\rho]$. After separating off the background and Hartree contributions to the density functional, $\Gamma_\lambda[\rho] = (1-\lambda)V_\Omega \cdot \rho + \frac{\lambda}{2}\,\rho \cdot V_{\mathrm{low}\,k} \cdot \rho + \tilde{\Gamma}_\lambda[\rho]$, the RG equation for the exchange-correlation functional $\tilde{\Gamma}_\lambda[\rho]$ is given by [4]

$$\partial_\lambda \tilde{\Gamma}_\lambda[\rho] = \frac{1}{2}\,\mathrm{Tr}\left[V_{\mathrm{low}\,k} \cdot \left(\frac{\delta^2 \tilde{\Gamma}_\lambda[\rho]}{\delta\rho\,\delta\rho} + \lambda\, V_{\mathrm{low}\,k}\right)^{-1}\right], \qquad (1)$$

where the solution to the RG equation yields the gs density $\rho_{\mathrm{gs};\lambda}$ and gs energy $E_{\mathrm{gs};\lambda} \sim \Gamma_\lambda[\rho_{\mathrm{gs};\lambda}]$. The method can be improved systematically, e.g., to include three-body forces, pairing correlations, external currents and a projection on center-of-mass momentum. We expect it to be applicable over a wide region of the nuclear chart.

References

[1] S.K. Bogner, T.T.S. Kuo and A. Schwenk, Phys. Rep. **386** (2003) 1; and Phys. Lett. **B** (2003) in press, nucl-th/0108041.

[2] A. Schwenk, B. Friman and G.E. Brown, Nucl. Phys. **A713** (2003) 191.
A. Schwenk, G.E. Brown and B. Friman, Nucl. Phys. **A703** (2002) 745.

[3] A. Schwenk and B. Friman, nucl-th/0307089.

[4] J. Polonyi and A. Schwenk, in preparation.

The Four-Body System in EFT

Lucas Platter

IKP, Forschungszentrum Jülich
D-52425 Jülich, Germany
Email: l.platter@fz-juelich.de .

The contact effective field theory provides an ideal framework to compute observables for low-energy nuclear and atomic systems to very high precision. The theory has been applied successfully to the three-body system with large two-body scattering length [1], where it was shown that a three-body force at leading order is necessary to renormalize the problem. The obvious next step is to apply this EFT to the four-body system. Answering the question of the relevance of a four-body force in this system would shed some light on the open problem of a general power-counting for many-body forces in the contact theory. Furthermore, there is a vast amount of four-body reactions where a high-precision theory would be very useful .
We have set up the integral equations for the four-body system but instead of solving these rather complicated integral equations directly, we restricted ourselves so far to solve a simplified version of these equations to get some insight into the problem. We found that in this case no four-body force is needed and speculate that this finding might hold for the full problem.
In parallel, we are investigating a second approach. We make use of the fact that we can write an effective potential which is equivalent to the contact theory and use the Yakubovski-equations to solve for the four-body binding energies. By varying the relevant parameters we hope to find a quick answer to the to the question of the importance of a four-body force at leading order[3].

This work is done in collaboration with H.-W. Hammer and U.-G. Meißner.

References

[1] P.F. Bedaque, U. van Kolck, H.-W. Hammer, Nucl. Phys. A **646** (1999) 444.

[2] W. Glöckle, H. Kamada, Nucl. Phys. A **560** (1993) 541, and references therein.

[3] L. Platter, H.-W. Hammer, U.-G. Meißner, in progress.

Electroproduction of Neutral Pions Off the Deuteron Near Threshold

Hermann Krebs

Universität Bonn
Helmholtz–Institut für Strahlen- und Kernphysik (Theorie)
Nussallee 14–16, D–53115 Bonn, Germany

In the last twenty years chiral perturbation theory (CHPT) became a standard tool for the description of low energy physics. Based only on the symmetries of QCD, especially on chiral symmetry, this theory gives a systematic and model–independent description of low energy processes. Nowadays CHPT is not only formulated for the light meson sector but also for interactions of Goldstone bosons with one or more nucleons and even nuclei [1].

In this talk I presented our analysis of electroproduction of neutral pions on the deuteron near the threshold using the framework of heavy baryon CHPT [2]. I firstly discussed the results of the third order calculation, where two unknown low energy constants appear. They are related to π^0–electroproduction on the neutron, which contributes to the single nucleon part (these are diagrams with the scattering only on one nucleon. The second nucleon remains untouched during the scattering process.). They are fixed from the fit to the total cross section data obtained at MAMI [3]. Finally I discussed our new fourth order calculation of the three body part (these are diagrams, where both nucleons are involved in the scattering process). No new unknown low energy constants appear at this order. I demonstrated the improvement of the third order results for the prediction of the differential cross sections and the S–wave amplitudes and concluded that our partial fourth order calculation is in fair agreement with the experiment [3].

References

[1] S. Weinberg, Phys. Lett. B 295 (1992) 114

[2] H. Krebs, V. Bernard, Ulf–G. Meißner, Nucl. Phys. A 713 (2003) 405

[3] I. Ewald et al., Phys. Lett. B 499 (2001) 238

Working Group IV: Chiral Lattice Dynamics

Convenors: Elisabetta Pallante and Martin Savage

Formal Lattice Issues

I. Montvay	*Partially Quenched Chiral Perturbation Theory and Numerical Simulations*	146
C. Aubin	*Pion and Kaon Properties in Staggered Chiral Perturbation Theory*	147
R. Mawhinney	*Recent Developments in Domain Wall Fermions*	n.r.
S. Sint	*Twisted Mass QCD and Lattice Approaches to the $\Delta I = 1/2$ Rule*	148
C. Hoelbling	*Overlap Phenomenology*	149

Mesons Properties from the Lattice

N. Shoresh	*Lattice QCD and Chiral Effective Lagrangians*	150
D. Bećirević	*Chiral Perturbation Theory, Heavy-light Mesons and Lattice QCD*	151
S. Dürr	*The Pion Mass in a Finite Volume*	152
G. Villadoro	*The Lattice Determination of $K \to \pi\pi$ in the $I=0$ Channel*	153
T. Chiarappa	*Investigation of the ϵ Regime of ChPT: Meson Correlation Functions*	154
S. Peris	*ϵ'/ϵ and the $\Delta I=1/2$ Rule in the $1/N_c$ Expansion*	155

Nucleon Properties from the Lattice

M. Göckeler	*Nucleon Form Factors from the QCDSF Collaboration*	156
S.R. Beane	*N and NN Properties in PQQCD*	157
D. Arndt	*Partially Quenched Chiral Perturbation Theory for the Baryon and Meson Charge Radii*	158
R. Lewis	*Strange Matrix Elements in the Nucleon*	159
M. Procura	*Nucleon Mass and Nucleon Axial Charge from Lattice QCD*	160

Future Directions and Focus

M. Golterman	*What should be addressed in the next five years*	161

Partially Quenched Chiral Perturbation Theory and Numerical Simulations

István Montvay

Deutsches Elektronen-Synchrotron DESY
Notkestr. 85, D-22603 Hamburg, Germany

The Gasser-Leutwyler constants [1] of the low energy Chiral Lagrangian in QCD can be computed in lattice QCD by numerical simulations. This is facilitated by the fact that, besides the possibility of changing momenta, on the lattice one can also change the masses of the quarks. Chiral Perturbation Theory (ChPT) can also be extended by changing the *valence quark masses* in quark propagators independently from the *sea quark masses* in virtual quark loops. In this way one arrives at Partially Quenched Chiral Perturbation Theory (PQChPT) [2]. PQChPT can be directly formulated on the lattice at non-zero lattice spacing [3] and this can be used for correcting leading lattice artifacts in the results [4].

In recent papers of the qq+q Collaboration [5,6,7,8] we reported on numerical simulations with two degenerate light quark flavours and investigated the dependence of the pseudoscalar meson masses and decay constants on sea and valence quark masses. The present simulations at small quark masses but on relatively coarse lattices give first estimates for the values of the Gasser-Leutwyler constants L_4, L_5, L_6 and L_8 which are within the expected range.

References

[1] J. Gasser and H. Leutwyler, Annals Phys. **158** (1984) 142.

[2] C.W. Bernard, M.F.L. Golterman, Phys. Rev. **D49** (1994) 486.

[3] S.R. Sharpe, R. Singleton, Jr., Phys. Rev. **D58** (1998) 074501.

[4] G. Rupak, N. Shoresh, Phys. Rev. **D66** (2002) 054503.

[5] qq+q Collaboration, F. Farchioni, C. Gebert, I. Montvay and L. Scorzato, Eur. Phys. J. **C26** (2002) 237.

[6] qq+q Collaboration, F. Farchioni, C. Gebert, I. Montvay, E. Scholz and L. Scorzato, Phys. Lett. **B561** (2003) 102.

[7] qq+q Collaboration, F. Farchioni, I. Montvay, E. Scholz and L. Scorzato, Eur. Phys. J. DOI: 10.1140/epjc/s2003-01321-3; hep-lat/0307002.

[8] qq+q Collaboration, F. Farchioni, C. Gebert, I. Montvay, E. Scholz and L. Scorzato, to appear in the Proceedings of the International Lattice Conference, 2003, Tsukuba; hep-lat/0309094.

Pion and Kaon Properties in Staggered Chiral Perturbation Theory

C. Aubin and C. Bernard
Washington University, St. Louis, MO 63130 USA

Extracting physical quantities from lattice simulations requires an extrapolation to physical quark masses as well as the continuum limit. The first limit can be performed using Chiral Perturbation Theory (χPT), while taking the continuum limit has been uncontrolled until recently. Using dynamical staggered fermions, scaling violations arising from taste symmetry breaking at $O(a^2)$ are not negligible. We must incorporate these taste violations into a staggered χPT (SχPT) to take the continuum limit systematically.

This has been done for a single flavor (one staggered field with four tastes) by Lee and Sharpe [1]. In order to apply this to lattice data, we need to generalize the Lee-Sharpe Lagrangian from Ref. [1] to multiple flavors. Additionally, after performing the one-loop calculations, we must remove the effects of the extra staggered tastes. The Lee-Sharpe Lagrangian for multiple flavors adds to the continuum chiral Lagrangian a term $a^2\mathcal{V}$, where $\mathcal{V}$ is the taste-symmetry breaking potential, written in full for multiple flavors in Ref. [2].

Using SχPT, we can calculate the masses and decay constants of pions and kaons, shown in Refs. [2,3]. Here we show the specific case of the kaon mass in the $m_u = m_d \equiv m_l \neq m_s$ limit, including all analytic terms. Writing $(m_{K_5^+}^{1-\mathrm{loop}})^2 = \mu(m_l + m_s)\,(1 + \delta_{m^2})$, and denoting the chiral logarithm as $\ell(m^2) = m^2 \ln m^2 +$ *finite volume corrections*:

$$\delta_{m^2} = \frac{1}{16\pi^2 f^2}\left(\frac{2a^2\delta'_V\left(\ell(m^2_{\eta'_V}) - \ell(m^2_{\eta_V})\right)}{m^2_{\eta'_V} - m^2_{\eta_V}} + [V \to A] + \frac{2}{3}\ell(m^2_{\eta_I})\right) + \frac{16\mu}{f^2}\,(2L_8 - L_5)\,(m_l + m_s) + \frac{32\mu}{f^2}\,(2L_6 - L_4)\,(2m_l + m_s) + a^2 C. \quad (1)$$

Expressions for the pion and kaon properties can be used to extract physical numbers to be compared with experiment. For example, we find preliminary values for the pion decay constant: $f_\pi^{\mathrm{th}} = 130.7(1.0)_{\mathrm{stat}}(3.5)_{\mathrm{sys}}$MeV ($f_\pi^{\mathrm{exp}} = 130.7(0.4)$MeV) and for the kaon decay constant: $f_K^{\mathrm{th}} = 157.4(1.0)_{\mathrm{stat}}(3.7)_{\mathrm{sys}}$MeV ($f_K^{\mathrm{exp}} = 159.8(1.5)$MeV). Also of interest is the combination of low-energy constants $2L_8 - L_5 = -0.3(1)(^{+1}_{-2})\times 10^{-3}$, since this number is well outside the allowed range for $m_u = 0$ to be a solution of the strong CP problem.

References

[1] W. Lee and S. R. Sharpe, Phys. Rev. D **60**, 114503 (1999).
[2] C. Aubin *et al.*, Nucl. Phys. **B** (Proc. Suppl.) **119** (2003). C. Aubin and C. Bernard, Phys. Rev. D **68** (2003) 034014 and hep-lat/0306026.
[3] C. Aubin, *et al.*, hep-lat/0309088.

Twisted Mass QCD and Lattice Approaches to the $\Delta I = 1/2$ Rule

Carlos Pena[1], **Stefan Sint**[2] and Anastassios Vladikas[3]

[1]DESY Theory Group, Notkestrasse 85 D-20607 Hamburg, Germany,

[2]Departamento de Física Teórica C-XI, Universidad Autónoma de Madrid, E-28049 Cantoblanco, Madrid, Spain.

[3]INFN, Sezione di Roma II, Dipartimento di Fisica, Università di Roma "Tor Vergata", Via della Ricerca Scientifica 1, I-00133 Rome, Italy

Twisted mass lattice QCD (tmQCD) was originally designed to eliminate the problem of unphysical zero-modes in lattice QCD with Wilson-type quarks [1,2]. This is achieved by adding a chirally twisted quark mass term which sets a lower bound on the spectrum of the Wilson-Dirac operator. Renormalised tmQCD can be transformed back to standard QCD by a (non-singlet) chiral rotation of the fields, and the renormalized tmQCD correlation functions can thus be interpreted as linear combinations of the correlation functions in standard QCD. As the chiral field rotation is not a lattice symmetry, the renormalization of bare lattice operators can be quite different in the two cases. This has already been used for the computation of B_K where the usual renormalization problems with Wilson fermions are avoided [2,3]. Here we generalise tmQCD to four Wilson quark flavours, and study the computation of $K \to \pi$ matrix elements of the CP conserving weak Hamiltonian [4]. We show that, with an active charm quark, the renormalization of the $K \to \pi$ matrix elements requires at most the subtraction of a linearly divergent counterterm. In the quenched approximation, the parameters can be chosen such that only a finite counterterm needs to be subtracted. As a result one may be able to match to lowest order chiral perturbation theory and obtain the effective couplings g_8 and g_{27}, which determine the $\Delta I = 1/2$ rule in non-leptonic kaon decays.

References

[1] R. Frezzotti, P.A. Grassi, S. Sint and P. Weisz, Nucl. Phys. Proc. Suppl. 73 (2000) 941.

[2] R. Frezzotti, P.A. Grassi, S. Sint and P. Weisz, JHEP 08 (2001) 058.

[3] P. Dimopoulos et al. [ALPHA Collaboration], hep-lat/0309134.

[4] C. Pena, S. Sint and A. Vladikas, Nucl. Phys. Proc. Suppl. **119**, 368 (2003); talk by S. Sint at "Lattice 2003", July 15-19 2003, Tsukuba, Japan, to appear in the proceedings; in preparation.

Overlap Phenomenology

Christian Hoelbling

Centre de Physique Théorique de Marseille (CNRS)
163 Avenue de Luminy, Case 907 13288 Marseille cedex 9, France

Quenched overlap fermions [1] are used to study QCD phenomenology. With the simulation technique described in [2], we found the mass of the strange quark [3]

$$m_s^{\overline{MS}}(2\text{GeV}) = 102(6)(18)\text{MeV} \tag{1}$$

and the chiral condensate

$$\frac{1}{N_f}\langle\bar{\psi}\psi\rangle^{\overline{MS}}(2\text{GeV}) = (267(5)(15)\text{MeV})^3 \tag{2}$$

We determined B_K [4]

$$B_K^{RGI} = 0.87(8)^{+2+14}_{-1-14} \tag{3}$$

where the first error is statistical, the second is systematic and the third is an estimate of the quenching error.

The calculation of other weak matrix elements including B_7 and B_8, of baryon spectra and the charmed meson decay constant is underway and preliminary results are reported in [5].

References

[1] H. Neuberger, Phys. Lett. B **417** (1998) 141 [arXiv:hep-lat/9707022].

[2] L. Giusti, C. Hoelbling and C. Rebbi, Nucl. Phys. Proc. Suppl. **106** (2002) 739 [arXiv:hep-lat/0110184].

[3] L. Giusti, C. Hoelbling and C. Rebbi, Phys. Rev. D **64** (2001) 114508 [Erratum-ibid. D **65** (2002) 079903] [arXiv:hep-lat/0108007].

[4] N. Garron, L. Giusti, C. Hoelbling, L. Lellouch and C. Rebbi, arXiv:hep-ph/0306295.

[5] F. Berruto, N. Garron, C. Hoelbling, L. Lellouch, C. Rebbi and N. Shoresh, arXiv:hep-lat/0310006.

Lattice QCD and Chiral Effective Lagrangians

Oliver Bär[1], Gautam Rupak[2], and **Noam Shoresh**[3]

[1]Institute of Physics, University of Tsukuba,
Tsukuba, Ibaraki 305-8571, Japan

[2]Lawrence Berkeley National Laboratory,
Berkeley, CA 94720, U.S.A.

[3]Department of Physics, Boston University,
Boston, MA 02215,U.S.A.[0.3cm]

Chiral perturbation theory (χPT) is a necessary tool in extracting information about QCD from lattice simulations, which are done with relatively heavy quark masses. We consider several low energy effective theories – extensions of χPT – which capture the leading discretization artifacts for certain lattice actions. The strategy is a two-step process. First one constructs the local Symanzik action for a given lattice action. Then, in close analogy to the derivation of χPT from QCD, one constructs a low energy chiral effective action. The resulting chiral Lagrangian reduces to the familiar chiral Lagrangian in the limit $a \to 0$. When a is non-zero, new terms appear in the Lagrangian with corresponding unknown coefficients (analogous to the Gasser-Leutwyler coefficients).

This method has been applied to the Wilson lattice action in Ref. [1] to order a, and to order a^2 in Refs. [2,3]. The chiral Lagrangian for staggered fermions has been derived in Refs. [4,5]. Another type of lattice action which has been analyzed is one with Ginsparg-Wilson valence quarks and Wilson sea quarks [6,2].

References

[1] G. Rupak and N. Shoresh, Phys. Rev. **D66** (2002) 054503, [hep-lat/0201019].

[2] O. Bar, G. Rupak and N. Shoresh, [hep-lat/0306021].

[3] S. Aoki, Phys. Rev. D **68** (2003) 054508, [hep-lat/0306027].

[4] W.-J. Lee and S. R. Sharpe, Phys. Rev. **D60** (1999) 114503, [hep-lat/9905023].

[5] C. Aubin and C. Bernard, Phys. Rev. D **68** (2003) 034014, [hep-lat/0304014].

[6] O. Bar, G. Rupak and N. Shoresh, Phys. Rev. D **67** (2003) 114505, [hep-lat/0210050].

Chiral Perturbation Theory, Heavy-light Mesons and Lattice QCD

Damir Bećirević[1], Saša Prelovšek[2,3], Jure Zupan[2,4]

[1] Laboratoire de Physique Théorique (Bât 210) [1], Université de Paris Sud, Centre d'Orsay, 91405 Orsay-Cedex, France.

[2]J. Stefan Institute, Jamova 39, P.O. Box 3000, 1001 Ljubljana, Slovenia.

[3]Department of Physics, University of Ljubljana, Jadranska 19, 1000 Ljubljana, Slovenia.

[4]Department of Physics, Technion–Israel Institute of Technology, Technion City, 32000 Haifa, Israel.

We study the chiral properties of the $B \to \pi$ and $B \to K$ transition form factors in the static heavy quark limit by employing the quenched, partially quenched and full (unquenched) chiral perturbation theory (ChPT). From the expressions derived at NLO in all three versions of ChPT we were able to examine the size of systematic uncertainties which currently plague the accurate determination of the form factors by means of lattice QCD, namely: the effects of quenched approximation, the impact of the chiral log terms on the extrapolation toward the physical pion mass. The complete discussion with details of calculations is presented in refs. [1,2]. In the case of partially quenched theory with $N_f = 2$ degenerate dynamical flavours we propose a simple chiral extrapolation strategy by which one can avoid the effects of divergent quenched chiral logarithms [2].

We also discuss the ξ parameter which enters decisevely into the standard CKM unitarity triangle analyses [3]. In particular, we propose to combine the f_{B_s}/f_B and f_K/f_π in double ratio in which the large chiral logarithms cancel, thus reducing the otherwise huge uncertainties in the chiral extrapolation of f_{B_s}/f_B. From the double ratio computed on the lattice, and the experimentally determined f_K/f_π, one can predict f_{B_s}/f_B, with a very small uncertainty.

References

[1] D. Becirevic, S. Prelovsek and J. Zupan, Phys. Rev. D **67** (2003) 054010.

[2] D. Becirevic, S. Prelovsek and J. Zupan, Phys. Rev. D **68** (2003) 074003.

[3] D. Becirevic, S. Fajfer, S. Prelovsek and J. Zupan, Phys. Lett. B **563** (2003) 150.

[1]Unité mixte de Recherche du CNRS - UMR 8627.

The Pion Mass in Finite Volume

Stephan Dürr[1] and Gilberto Colangelo[2]

[1]DESY Zeuthen, 15738 Zeuthen, Germany
[2]Institute for Theoretical Physics, University of Bern, 3012 Bern, Switzerland

When studying QCD in a finite box $L^3 \times T$ with toroidal boundary conditions (as is standard in lattice calculations with $T \gg L$), properties of the QCD Hamiltonian will reflect the finite spatial extent. In particular, the relative shift

$$R_M(M_\pi, L) = \frac{M_\pi(L) - M_\pi}{M_\pi} \qquad \text{where} \quad M_\pi \equiv M_\pi(L=\infty) \tag{1}$$

is a systematic effect that one would like to have analytically predicted.

One way is due to Lüscher who noted that the finite-size effect in Euclidean space follows from the π-π forward scattering amplitude in Minkowski space [1]

$$R_M(M_\pi, L) = -\frac{3}{16\pi^2 M_\pi^2 L} \int_{-\infty}^{\infty} dy\; F(\mathrm{i}y)\, e^{-\sqrt{M_\pi^2+y^2}\,L} + O(e^{-\overline{M}L}) \tag{2}$$

where the generic bound $\overline{M} \geq \sqrt{3/2}M$ can be specified to $\overline{M} = \sqrt{2}M_\pi$. To LO in the chiral expansion the amplitude F is constant (in y), hence the integral in (2) can be done analytically [2]. Using NLO and NNLO chiral input, the resulting expressions for R_M become more involved [3,4], necessitating in the latter case a thorough discussion of the impact of the uncertainties of the NLO low-energy constants involved. The advantage is that one can explicitly watch the chiral convergence behavior at fixed M_π and L.

The other approach is to directly compute the pion mass in chiral perturbation theory in a finite volume. This has been done at 1-loop level in [5], resulting in

$$R_M(M_\pi, L) = \frac{1}{2N_\mathrm{f}} \xi\, \tilde{g}_1(\lambda) + O(\xi^2) \tag{3}$$

with $\xi = M_\pi^2/(4\pi F_\pi)^2$ and $\lambda = M_\pi L$. In [4] we provide a representation of $\tilde{g}_1$ which is handy for a numerical evaluation, and we include a table where the predictions of both approaches are compared for a reasonable range of M_π and L values.

References

[1] M. Lüscher, Commun. Math. Phys. **104**, 177 (1986).

[2] M. Fukugita, *et al.*, Phys. Rev. D **47**, 4739 (1993).

[3] G. Colangelo, S. Dürr and R. Sommer, Nucl. Phys. Proc. Suppl. **119**, 254 (2003) [hep-lat/0209110].

[4] G. Colangelo and S. Dürr, DESY 03-177

[5] J. Gasser and H. Leutwyler, Phys. Lett. B **184** 83 (1987).

The Lattice Determination of $K \to \pi\pi$ in the $I = 0$ Channel

Giovanni Villadoro

Dip. di Fisica, Univ. di Roma "La Sapienza" and INFN,
Sezione di Roma, P.le A.Moro 2, I-00185 Rome, Italy

In this talk, a brief review was presented about recent developments in understanding problems related to the extraction of $I = 0$ $K \to \pi\pi$ matrix elements from the lattice. In particular the talk focused on the interplay between finite volume corrections, final state interaction and the quenching approximation. After a short introduction about historical milestones on the subject, such as the Maiani-Testa no-go theorem and finite volume Lüscher formulae (see ref. [1]), the general strategy to extrapolate the amplitude to the physical point, through the use of χPT to NLO, was described [2]. Two recent procedures of implementing χPT in this type of calculations and the relevant formulae can be found in refs. [2,3]. Finite volume analysis has been discussed then in full, quenched (q) and partially quenched (pq) QCD by using χPT. In particular it was discussed how, in qQCD, the lack of unitarity prevents the extraction of matrix elements from correlators with multi-particle external states [4]. More recently, in ref. [5], it has been shown, however, that pqQCD can avoid this problem for particular choices of masses and momenta. More in details, the $I = 0$ $K \to \pi\pi$ matrix element can be extracted from pq simulations provided that sea quarks be degenerate with the valence ones and the c.o.m. energy be under threshold for production of quenched flavors. Outlook and consequences on actual numerical simulations were presented in the conclusion.

References

[1] L. Maiani and M. Testa, *Phys. Lett. B* **245** (1990) 585. M. Luscher, Quantum Field Theories. 2. Scattering States," *Commun. Math. Phys.* **105** (1986) 153. L. Lellouch and M. Luscher, *Commun. Math. Phys.* **219** (2001) 31. C. J. Lin, G. Martinelli, C. T. Sachrajda and M. Testa, *Nucl. Phys. B* **619** (2001) 467.

[2] P. Boucaud *et al. Nucl. Phys. Proc. Suppl.* **106** (2002) 323. C. J. Lin, G. Martinelli, E. Pallante, C. T. Sachrajda and G. Villadoro, *Nucl. Phys. B* **650** (2003) 301, *Nucl. Phys. Proc. Suppl.* **119** (2003) 383.

[3] J. Laiho and A. Soni, *Phys. Rev. D* **65** (2002) 114020 and `hep-lat/0306035`.

[4] C. J. Lin, G. Martinelli, E. Pallante, C. T. Sachrajda and G. Villadoro, *Phys. Lett. B* **553** (2003) 229, *Nucl. Phys. Proc. Suppl.* **119** (2003) 371.

[5] C. J. Lin, G. Martinelli, E. Pallante, C. T. Sachrajda and G. Villadoro, `hep-lat/0308014`.

Investigation of ϵ Regime of ChPT: Meson Correlation Functions

T. Chiarappa[1], W. Bietenholz[2], K. Jansen[1], K.-I. Nagai[1] and S. Shcheredin[2]

[1]NIC/Desy Zeuthen,
Platanenallee 6, D-15738 Zeuthen, Germany

[2] Institut für Physik, Humboldt Universität zu Berlin,
Newtonstr. 15, D-12489 Berlin, Germany

Surprisingly, physical information can be extracted from unphysical setups, like from a finite volume via a finite size analysis. This framework is highly suitable for lattice simulations.
One analytic tool to study such a setup is the ϵ-expansion of Chiral Perturbation Theory (ChPT) [1], which allows to extract the characteristic Low Energy Constants (LEC) by confronting its prediction with numerical data.
In contrast to the infinite volume case, in a finite box ($V = L^4$) gauge field of fixed topological sectors play a prominent rôle [2].
This turns out to be expecially important in the ϵ-regime of ChPT, where the Compton wavelength of the pion does not fit at all into the box ($Lm_\pi < 1$).
From a numerical point of view, the possibility to simulate chiral fermions on the lattice is one of the great achievement of the last years [3].
This new lattice world is very cost demanding, forcing us (so far) to restrict simulations to the quenched approximation.
Based on two recent analytical works [4], a numerical pilot study of the 2-point meson correlation functions in the ϵ-regime of quenched chiral perturbation theory was presented [5], using the overlap formalism with $\beta = 6.0$ and two lattices of physical volume 1.12 fm and 0.93 fm.
From the data relative to the axial correlation function, it was possible to extract the quenched bare pion decay constant to be $F_\pi^b = 86.7 \pm 4.0$ MeV.
The determination of other LEC is in progress.

References

[1] J. Gasser and H. Leutwyler, Phys. Lett. **B188**: 477 (1987)

[2] H. Leutwyler and A. Smilga, Phys. Rev. **D46**: 5607 (1992)

[3] H. Neuberger, Phys. Lett. **B417**: 141 (1998) ; Nucl. Phys. **B412**: 574 (1994)

[4] P. H. Damgaard et al., Nucl. Phys. **B629**: 445 (2002) ; Nucl. Phys. **B656**: 226 (2003)

[5] T. Chiarappa et al., hep-lat/0309083 ; K. Nagai et al., hep-lat/0309051

ϵ'/ϵ and the $\Delta I = 1/2$ Rule in the $1/N_c$ Expansion

S. Peris

Grup de Fisica Teorica and IFAE
Universitat Autonoma de Barcelona
08193 Barcelona, Spain

In this talk, after briefly discussing how to analytically compute unfactorized contributions to weak matrix elements in the large-N_c expansion [1], I reported on the new results for the matrix elements of the Q_6 and Q_4 penguin operators obtained in [2]. These results include unfactorized contributions of $\mathcal{O}(N_c^2 \times \frac{n_f}{N_c})$ and exhibit analytic matching between short– and long–distance scale dependence in the $\overline{\rm MS}$ scheme. It is found a numerically large and positive contribution to the $\Delta I = 1/2$ matrix element of Q_6 and hence to the direct CP-violation parameter ε'/ε.

The implications of these results for a quenched theory were analyzed in [3] where the conclusion was reached that the pure quenching artifact, α_q^{NS}, the quenched octet coupling $\alpha_{q1}^{(8,1)}$ and its counterpart in the unquenched theory, $\alpha_1^{(8,1)}$[4], obey the hierarchy $\alpha_q^{NS} \gg \alpha_{q1}^{(8,1)} \gg \alpha_1^{(8,1)}$. Since α_q^{NS} is currently neglected in lattice analyses of ε'/ε, the result just mentioned could help explain the present discrepancy between lattice and experimental results[5][6].

Results were also obtained for the $\Delta I = 1/2$ rule in $K \to \pi\pi$ amplitudes. Large "eye–diagram" contributions of $\mathcal{O}(N_c^2 \times \frac{1}{N_c})$ from the Q_2 operator were found which are related to the unfactorized contributions from Q_4 mentioned above. The results lead to an enhancement of the $\Delta I = 1/2$ effective coupling. A common origin for all the violation of factorization that we find was discussed in terms of the relevant scales in the problem.

I am very thankful to M. Golterman, T. Hambye and E. de Rafael for a most pleasant and fruitful collaboration.

References

[1] See, e.g., S. Peris, hep-ph/0204181; E. de Rafael, hep-ph/0210317.

[2] T. Hambye, S. Peris and E. de Rafael, JHEP **0305** (2003) 027.

[3] M. Golterman and S. Peris, hep-lat/0306028.

[4] M. Golterman and E. Pallante, JHEP **0110** (2001) 037.

[5] T. Blum *et al.*, hep-lat/0110075; J. I. Noaki *et al.*, Phys. Rev. D **68** (2003) 014501.

[6] See, e.g., A. Alavi-Harati *et al.*, Phys. Rev. D **67** (2003) 012005.

Nucleon Form Factors from the QCDSF Collaboration

M. Göckeler[1,2]

[1]Institut für Theoretische Physik, Universität Leipzig,
D-04109 Leipzig, Germany

[2]Institut für Theoretische Physik, Universität Regensburg,
D-93040 Regensburg, Germany

We have computed the electromagnetic form factors of the nucleon in quenched lattice QCD, using non-perturbatively improved Wilson fermions [1]. The analysis focusses on the isovector case as in this sector the quark-line disconnected contributions, which we have not evaluated, cancel. We want to compare the pion-mass dependence of the Monte Carlo results with predictions of chiral effective field theory. Since the pions in the simulations are relatively heavy ($m_\pi > 500\,\mathrm{MeV}$) quenching effects are expected to be small, and one can use ordinary chiral perturbation theory rather than quenched chiral perturbation theory. More specifically, we consider formulae obtained within the small scale expansion [2,3], which has explicit nucleon and Δ degrees of freedom. As the momenta in our simulations are too large for a comparison of the form factors themselves, we fit the Q^2 dependence of our form factors with a dipole ansatz and compare only the resulting radii as well as the anomalous magnetic moment. While a reasonable connection between the Monte Carlo data and the physical value is achieved for the Pauli radius and the anomalous magnetic moment, the Dirac radius is more problematic [1]. It seems that the leading one-loop calculation in the small scale expansion is not accurate enough to describe the quark-mass dependence of the isovector Dirac radius up to the masses used in present simulations. It remains to be seen whether additional terms in the expansion can solve this problem. On the other hand, the simulations will progress towards smaller quark masses so that eventually contact with chiral perturbation theory should be established.

References

[1] M. Göckeler, T.R. Hemmert, R. Horsley, D. Pleiter, P.E.L. Rakow, A. Schäfer, G. Schierholz, hep-lat/0303019.

[2] V. Bernard, H.W. Fearing, T.R. Hemmert and U.-G. Meißner, Nucl. Phys. A635 (1998) 121; 642 (1998) 563 (E).

[3] T.R. Hemmert and W. Weise, Eur. Phys. J. A15 (2002) 487.

N and NN Properties in PQQCD

Silas R. Beane[1,2]

[1]Department of Physics, University of New Hampshire, Durham, NH 03824

[2]Jefferson Laboratory, 12000 Jefferson Avenue, Newport News, VA 23606

Presently, unquenched lattice simulations with the physical values of the light-quark masses are prohibitively time-consuming, even on the fastest machines. Relatively recently it was realized that partially-quenched (PQ) simulations, in which the sea quarks are more massive than the valence quarks, provide a rigorous method to determine QCD observables and are much less time-consuming than their QCD counterparts. Technology has been developed to describe partially-quenched QCD (PQQCD) with PQχ-PT (Ref. [1] and references therein). It is hoped that future lattice simulations can be performed with sufficiently small quark masses where the chiral expansion is convergent, and can be used to extrapolate down to the quark masses of nature. Recently, meson and baryon properties have been studied extensively in PQχ-PT [2,3,4,5,6,7]. Two-nucleon systems have also been explored in PQQCD [8,9]. An important further motivation for PQQCD is the additional "knobs" that are provided by exploring the "plane" of sea and valence quark masses and which increase the consistency checks available to lattice simulations.

References

[1] S.R. Sharpe and N. Shoresh, Phys. Rev. **D62**, 094503 (2000).

[2] J.-W. Chen and M.J. Savage, Phys. Rev. **D65**, 094001 (2002).

[3] S.R. Beane and M.J. Savage, Nucl. Phys. **A709**, 319 (2002).

[4] S.R. Beane and M.J. Savage, *Nucl. Phys.* **B636**, 291 (2002),

[5] J.-W. Chen and M.J. Savage, *Phys. Rev.* **D66**, 074509 (2002),

[6] D.B. Leinweber, `hep-lat/0211017`.

[7] D. Arndt and B.C. Tiburzi, `hep-lat/0307003`, `hep-lat/0308001`, `hep-lat/0309013`.

[8] S.R. Beane and M.J. Savage, *Phys. Lett.* **B535**, 177 (2002); *Phys. Rev.* **D67**, 054502 (2003).

[9] D. Arndt, S.R. Beane and M.J. Savage, `nucl-th/0304004`.

Partially Quenched Chiral Perturbation Theory for the Baryon and Meson Charge Radii

Daniel Arndt and Brian C. Tiburzi

Department of Physics, Box 351560,
University of Washington, Seattle, WA 98195-1560, USA

The study of electromagnetic form factors of the hadrons at low momentum transfer provides important insight into the non-perturbative structure of QCD. A number of new lattice QCD calculations of these form factors have recently appeared [1]. These calculations use the quenched approximation of QCD (QQCD); partially quenched calculations (PQQCD) are expected in the near future. Since currently and foreseeably lattice calculations cannot be performed with the physical masses of the light quarks, it is crucial to know how to properly extrapolate the results of these calculations from the quark masses used on the lattice to those in nature. To aid in this extrapolation, low-energy effective theories—quenched and partially quenched chiral perturbation theory (QχPT, PQχPT)—have been developed [2].

We calculate [3] the electric charge radii of the $SU(3)$ meson and baryon octets in QχPT and PQχPT. We work in the isospin limit, up to next-to-leading order in the chiral expansion, and to leading order in the heavy baryon expansion. We find that, while the expansions about the chiral limit for QχPT and PQχPT are formally similar, $< r^2 > \sim \alpha + \beta \log m_Q + \ldots$, the QQCD result exhibits quenched oddities: for Σ^- and Ξ^-, for example, $\beta = 0$ and these radii are independent of the quark masses m_Q! Such behavior, also found for the mesons, illustrates how QQCD has no known connection to QCD. PQQCD, however, is smoothly connected to QCD and the low-energy constants of PQχPT are found to be the same as those of χPT. Hence, our PQχPT result will enable proper extrapolation of PQQCD lattice simulations to QCD.

References

[1] J. van der Heide et. al., Phys. Lett. B **566**, 131 (2003); M. Gockeler et. al (QCDSF), arXiv:hep-lat/0303019; A. Tang, W. Wilcox and R. Lewis, arXiv:hep-lat/0307006.

[2] C. W. Bernard and M. F. Golterman, Phys. Rev. D **46**, 853 (1992); J. N. Labrenz and S. R. Sharpe, Phys. Rev. D **54**, 4595 (1996); S. R. Sharpe and N. Shoresh, Phys. Rev. D **62**, 094503 (2000); J. W. Chen and M. J. Savage, Phys. Rev. D **65**, 094001 (2002).

[3] D. Arndt and B. C. Tiburzi, arXiv:hep-lat/0307003.

Strange Matrix Elements in the Nucleon

Randy Lewis[1], Walter Wilcox[2] and R. M. Woloshyn[3]

[1]Department of Physics, University of Regina, Regina, SK, S4S 0A2, Canada

[2]Department of Physics, Baylor University, Waco, TX, 76798-7316, USA

[3]TRIUMF, 4004 Wesbrook Mall, Vancouver, BC, V6T 2A3, Canada

The strange-quark current matrix elements of the nucleon probe non-valence degrees of freedom. This physics can be studied within quenched lattice QCD by invoking a stochastic method for the strange quark propagator[1]. Pioneering studies of the strangeness electromagnetic form factors demonstrated that a signal can be difficult to extract from lattice simulations[2], so our recent work[3] includes a study with many more configurations than previously used, as well as two different choices for the number of noises in the stochastic propagator. A clear signal is seen for the scalar matrix element, but the electric and magnetic matrix elements can only be bounded by an uncertainty consistent with zero.

Ref. [3] discusses the chiral extrapolations of these strangeness matrix elements through quenched SU(3) chiral perturbation theory (ChPT) calculations. Some pieces of the calculations can also be found in Ref. [4]. It is noted that the scalar, electric and magnetic cases share the same low energy constants (LEC's), so the established scalar signal from lattice data constrains the electric and magnetic extrapolations. There could be advantages to using SU(2) ChPT instead, and a calculation exists[5], but in SU(2) ChPT the scalar LEC's differ from the electromagnetic LEC's, so it seems that the application of SU(2) ChPT to electromagnetic matrix elements must wait for a clearer lattice QCD signal.

References

[1] S. J. Dong and K. F. Liu, Nucl. Phys. B (Proc. Suppl.) 26, 353 (1992); Phys. Lett. B328, 130 (1994).

[2] S. J. Dong, K. F. Liu and A. G. Williams, Phys. Rev. D58, 074504 (1998); N. Mathur and S. J. Dong, Nucl. Phys. B (Proc. Suppl.) 94, 311 (2001); W. Wilcox, Nucl. Phys. B (Proc. Suppl.) 94, 319 (2001).

[3] R. Lewis, W. Wilcox and R. M. Woloshyn, Phys. Rev. D67, 013003 (2003).

[4] B. Borasoy, Eur. Phys. J. C8, 121 (1999); T. R. Hemmert, B. Kubis and U.-G. Meißner, Phys. Rev. C60, 045501 (1999); S. J. Puglia, M. J. Ramsey-Musolf and Shi-Lin Zhu, Phys. Rev. D63, 034014 (2001); M. J. Savage, Nucl. Phys. A700, 359 (2002).

[5] J. W. Chen and M. J. Savage, Phys. Rev. D66, 074509 (2002).

Nucleon Mass and Nucleon Axial Charge from Lattice QCD

M. Procura[1,2], T.R. Hemmert[1] and W. Weise[1,2]

[1]Physik-Department, Theoretische Physik, TU München,
D-85747 Garching, Germany

[2]ECT*, Villa Tambosi, I-38050 Villazzano (Trento), Italy.

Lattice QCD on one side and Chiral Perturbation Theory (χPT) on the other, are progressively developing as important tools to deal with the non-perturbative aspects of hadron structure. At present, however, there is a gap between the relatively large quark masses accessible in fully dynamical lattice simulations of nucleon properties and the small quark masses relevant for comparison with physical observables. In our work we explore the consequences of a direct application of χPT in the range of presently accessible quark masses on the lattice and the feasibility of a systematic approach based on chiral effective Lagrangian for the extrapolation of nucleon properties from lattice QCD.

In [1] we focus on the quark mass dependence of the nucleon mass M_N and present an extrapolation of this observable from two-flavor lattice QCD results in the framework of relativistic baryon χPT up to order p^4 in infrared regularization scheme. Already at leading one-loop order we obtain a good chiral extrapolation function and the next-to-leading one-loop corrections turn out to be reasonably small. From the p^4 extrapolation we find $M_N = (0.91 \pm 0.07)\,\mathrm{GeV}$ together with the pion-nucleon sigma term $\sigma_N = (53 \pm 8)\,\mathrm{MeV}$, at the physical value of the pion mass. Including the empirical M_N as a constraint gives the nucleon mass in the chiral limit $M_0 \approx 0.89\,\mathrm{GeV}$, and $\sigma_N \approx 47\,\mathrm{MeV}$.

We study also the quark mass expansion of the axial-vector coupling constant g_A of the nucleon [2]. We compare two versions of non-relativistic chiral effective field theory: Heavy Baryon χPT and an extension of that (Small Scale Expansion) which incorporates in a systematic framework explicit $\Delta(1232)$ resonance degrees of freedom. It turns out that, in order to approach the physical value of g_A in a leading one-loop calculation, the inclusion of the explicit $\Delta(1232)$ degrees of freedom is crucial. With information on important higher order couplings constrained from analyses of the $\pi N \to \pi\pi N$ reaction, a chiral extrapolation function $g_A(m_\pi)$ is obtained, which works well from the chiral limit across the physical point into the region of present lattice data.

References

[1] M. Procura, T.R. Hemmert and W. Weise, [hep-lat/0309020].

[2] T.R. Hemmert, M. Procura and W. Weise, Nucl. Phys. **A721**, 938c (2003) and [hep-lat/0303002], Phys. Rev. D, in print.

"What should be addressed the next five years"

Maarten Golterman

Dept. of Physics and Astronomy, SFSU, San Francisco, CA 94044, USA

I collected and discussed a number of methodological issues I believe to be important for extracting hadronic physics from Lattice QCD, restricting myself to topics involving the use of effective field theory. Since almost all Lattice QCD computations are "unphysical" (wrong quark masses, unphysical momenta, small volume, *etc.*), EFT techniques such as ChPT are essential in extracting physics from Lattice QCD. An important new application of EFT is its use in parameterizing in a systematic way $O(a)$ and $O(a^2)$ corrections [1,2].

(Partially) quenched Lattice QCD with the number of sea quarks $N_f \neq 3$ leads to *uncontrolled* errors and unphysical effects. These can*not* be estimated/corrected for in ChPT, since the (partially) quenched low-energy constants can be different, and run differently. Since many Lattice QCD results at present are still obtained at $N_f = 0, 2$, it's useful pursuing quantitative *analytic* estimates of such differences; an example for the penguin Q_6 is in [3]. Because of the importance of $N_f = 3$ computations, one should consider using staggered fermions for sea quarks – even though there are issues to be understood (*e.g.* fractional powers of the fermion determinant and "mismatched" zero modes).

An important issue is the convergence of ChPT, both on the lattice (does it converge for typical lattice meson masses? – see *e.g.* [4] for examples of poor convergence) and in the real world (does it converge at M_K?). It does not help to stick LECs computed on the lattice into real-world ChPT if that does not converge (getting M_K right on the lattice, and extrapolating in $m_{u,d}$ [5] may help). I emphasized that LECs are interesting and easier than physical quantities ($g_{8,27}$ are easier than ε'/ε!), and phenomenological or model estimates are often available. Comparison with analytic approaches [6] should be interesting (example: $\Pi_{LR}(Q^2)$, for $\Pi_{EM}(Q^2)$ see [7]), because they have typically more difficulty outside the chiral limit, while the lattice has trouble getting to the chiral limit.

References

[1] N. Shoresh, contribution to this working group and refs. therein
[2] C. Aubin, contribution to this working group and refs. therein
[3] M. Golterman and S. Peris, Phys. Rev. D67 (2003) 096001
[4] RBC coll., hep-lat/0110075; Young *et al.*, Prog. Part. Nucl. Phys. 50 (2003) 399; Dong *et al.*, hep-lat/0304005; M. Procura, contr. to this working group
[5] see *e.g.* J. Chen and M. Savage, Phys. Rev. D66 (2002) 074509
[6] S. Peris, contribution to this working group and refs. therein
[7] T. Blum, PRL 91 (2003) 052001; M. Göckeler *et al.*, hep-lat/0310027